



US006518037B2

(12) **United States Patent**
Apicella et al.

(10) **Patent No.:** **US 6,518,037 B2**
(45) **Date of Patent:** ***Feb. 11, 2003**

(54) **TWO-COMPONENT SYSTEM THAT CONTROLS BACTERIAL MEMBRANE SYNTHESIS**

(75) Inventors: **Michael A. Apicella**, Solon, IA (US); **Andrew Preston**, Cambridge (GB)

(73) Assignee: **University of Iowa Research Foundation**, Iowa City, IA (US)

(*) Notice: This patent issued on a continued prosecution application filed under 37 CFR 1.53(d), and is subject to the twenty year patent term provisions of 35 U.S.C. 154(a)(2).

Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **09/439,226**

(22) Filed: **Nov. 12, 1999**

(65) **Prior Publication Data**

US 2001/0016349 A1 Aug. 23, 2001

(51) **Int. Cl.**⁷ **C12Q 1/18**; C12Q 1/68; C12Q 1/02; C12N 1/00; C12N 1/20

(52) **U.S. Cl.** **435/32**; 435/6; 435/7; 435/29; 435/32; 435/243; 435/871

(58) **Field of Search** 424/249.1, 250.1; 435/252.1, 29, 32, 6, 7, 243, 871

(56) **References Cited**

U.S. PATENT DOCUMENTS

6,180,111 B1 * 1/2001 Stein et al.

FOREIGN PATENT DOCUMENTS

WO 9817805 * 4/1998
WO 99/59625 11/1999

OTHER PUBLICATIONS

Pettit, R.K. et al, *Molecular Microbiology*, 6(6), 729-734. The interaction of naturally elaborated blebs from serum-susceptible and serum-resistant strains of *Neisseria gonorrhoeae* with normal human serum. 1992.*
Dorward, D.W. et al., *Journal of Bacteriology*, 171 (8), 4196-4201. DNA-binding proteins in cells and membrane blebs of *Neisseria gonorrhoeae*. 1989.*

Andersen, B.M. et al., *Acta Path. Microbiol. Scand. Sect. B*, 89: 271-278, 1981. Loss of endotoxin liberation in *Neisseria meningitidis*. 1981.*

Bulygin, V.V., et al., "Rotation of the ϵ Subunit during Catalysis by *Escherichia coli* F₀F₁-ATP Synthase", *J. Biol. Chem.*, 273, 31765-31769 (1998).

Meadow, P.M., et al., "The Effect of Lipopolysaccharide Composition on the Ultrastructure of *Pseudomonas aeruginosa*", *J. Gen. Microbiol.*, 105, 23-28 (1978).

Preston, A., et al., "A Two-Component Regulatory System Involved in LOS Microheterogeneity in *Neisseria gonorrhoeae*", *Abstracts Gen. Meet. Asm.*, 96, 208 (1996).

Pettit, R.K., et al., "Characterization of Naturally Elaborated Blebs from Serum-Susceptible and Serum-Resistant Strains of *Neisseria gonorrhoeae*", *Mol. Microbiol.*, 6, 723-728 (1992).

Ramirez-Arcos, S., et al., "Expression and Interaction of the minCDE Genes of *Neisseria gonorrhoeae*", *Abstracts Gen. Meet. Asm.*, 99, 352 (1999).

Suzuki, H., et al., "Murein-Lipoprotein of *Escherichia coli*: a protein involved in the stabilization of bacterial cell envelope", *Mol. Gen. Genet.*, 167, 1-9 (1978).

Watnick, R.S., et al., "*Escherichia coli* NusA is Required for Efficient RNA Binding by Phase HK022 Nun Protein", *PNAS*, 95, 1546-1551 (1998).

Zhou, D., et al., "Lipoooligosaccharide Biosynthesis in *Neisseria gonorrhoeae*: Cloning, Identification and Characterization of the α 1,5 Heptosyltransferase I gene (rfaC)", *Molecular Microbio.*, 14, 609-618 (1994).

* cited by examiner

Primary Examiner—Christopher R. Tate
Assistant Examiner—Michele C. Flood
(74) *Attorney, Agent, or Firm*—Schwegman, Lundberg, Woessner & Kluth, P.A.

(57) **ABSTRACT**

The present invention discloses a mutant *Neisseria* having extensive membrane blebbing, both an indicium and a cause of virulence in the gonococcus and meningococcus. Methods are disclosed for making and characterizing the mutant, bmrRS. Methods are disclosed for isolating bmrRS membranes for use as a vaccine. Methods are also disclosed for the use of the mutant for determining the virulence of clinical samples of *N. gonorrhoeae* and *N. meningitidis*. Methods are also disclosed for the screening of antibiotics targeted to virulent *Neisseria*.

5 Claims, 4 Drawing Sheets




Fig. 1

BASR

AAA AAQPYDLMLL
 NM: MRVLLVEDDA MIAEAVSASL KDGGYAVDWV KNGARLPLPS LMT*
 GC: MRVLLVEDDA MIA*AVSASL KDGGYAVDWV KNGAQVAAAA AAQPYDLMLL
 NC: MRVLLVEDDA MIAEAVSANL KDGGYAVDWV KNGAQVAAAA AAQPYDLMLL

DLGLLGRDGL DV

NM: MFVPL(BasS) . . .
 GC: DLGLPGRDGL DVLSEIRAAG CTVPVLIVTA RDDLYSRLNG LDGGADDYIV
 NC: DLGLPGRDGL DVLSEIRAAG CTVPVLIVTA RDDLYSRLNG LDGGADDYIV

GC: KPFDMAEFKA RMRAVLRGRS GQAQACLSNG ALSLNPATYQ VEIIAEGROV
 NC: KPFDMAELKA RMRAVLRGRS GQAQACLSNG ALSLNPATYQ VEIIAERROV

GC: ALSNQEFVSL QALLARPGVI LSRSDSEDKV YGWGGEVESN AVDFLIHGLC
 NC: ALSNKEFAVL QALLARPGVI LSRSDLEDKI YGWGAEVESN AVDFLIHALR

GC: KKLKESIQN VRGVGWLMPR QDAV*
 NC: KKLKENIQN VRGVGWLVPV AV*

BASS

NM: MFVPLAM LAGMFSYYET FHETEALQDD
 GC: MPDRFFKILK HSLQVRISLA LIWMFVPLAM LAGMFSYYET FHETEALQDD
 NC: MLNRLIRMLK QSLQVRICIA LILMFLPLAM VAGVFSYYDT FHEAEELQDD

NM: LLRQAALYVA PDSKPELPE GDGDTRIFVQ MPQQEDPVVS LPAHLADGLH
 GC: LLRQAALYVG PDSKSETLPE GDGDTRILVQ MPQQEDPVVS LPAHLADGLH
 NC: LLRQTALYVG PDYHPDALPE GDGDTRILVQ MPDQE.PIVS LPMHLKDGLH

NM: TLQADGDDDY YRVYIRTTEQ GRIAVMQENE YREDLAADA QSVLPLLAA
 GC: TLRADGDDDY YRAYIRTTEQ GRIAVMQENE YREDLAEDA QSVLPLLAA
 NC: TLRADEDDEY YRVYIRTTR GRIAVMQENE YREDLAADA QSVLPLLAA

NM: LPLMILLTVW ITHKAMPVR KLSQSLEQRR INDLALSVD NIPSEIRGFV
 GC: LPLMILLTVW ITHKAMPVR KLSQSLEQRR INGLPALSVD NIPSEIRGFV
 NC: LPLIILLTVW ITHQAMPVR ILSQNLQRR LDDLALNTD NIPSEIKGFV

NM: TAINLLKRA DEDIRHRQRF VADAAHELRT PMTALSLOAE RLNNMSLPPD
 GC: TAINLLKRV DEDIRRRQRF VADAAHELRT PMTALSLOAE RLNNMPLPPD
 NC: TAINLLKRV DEDIRRRQRF IADAAHELRT PMTALSLOAE RLNNMPLPPD

NM: AARQPAVLQQ SIRRNKHLE QLLALARSQS DETPLTKTTF GLQSRFRQVL
 GC: AARQSAVLQQ SIRRNKHLE QLLALARSQS DEPPLAKTTF GLQSRFRQVL
 NC: AARQSAVLQQ SIRRNKHLE QLLALARSQS DETPLTKTTF GLQSRFRRLV

NM: QELMPLALEK RQDIGVAVGG DVEVSADETE IYTLIKTFAD NAVRYTPPEG
 GC: QELMPLVLEK RQDIGVAVGG DVEVSADETE IYTLVKTTFAD NAVRYTPNGG
 NC: QELMPLALEK RQDIGVAVEG DFEIFADETE IYTLVKTTFVD NAIRYTPSEG

NM: RIDLGFTDEG KYLAVWVEDN GNGIPESERA RVLDPFYRIL GTEQQGTGLG
 GC: RIDLGFTDEG KYLAVWVEDN GNGIPESECA RVLDPFYRIL GTEQQGAGLG
 NC: RIDLGFTDEG KYLAVWVEDN GKGIPESER

NM: LSIADTLAKK YGGHLELTD RRFHGLLIR ALLDKETLK*
 GC: LSIADTLAKK YGGYLELTD RRFHGLLIR ALLDKETLK*

Fig. 2





Fig. 3

Fig. 4



TWO-COMPONENT SYSTEM THAT CONTROLS BACTERIAL MEMBRANE SYNTHESIS

The U.S. government sponsored the research leading to this invention under grants AI18384 and AI43924 from the National Institutes of Health. The Government retains certain rights in this invention.

BACKGROUND OF THE INVENTION

Neisseria gonorrhoeae ("gonococcus") and *N. meningitidis* ("meningococcus") are Gram-negative diplococci that are strictly human pathogens. *N. gonorrhoeae* causes primarily urethritis in males and pelvic inflammatory disease in females. *N. meningitidis* is the causative agent of middle ear infections and meningitis.

As is commonly found in other bacterial pathogens, strains of *Neisseria* sp. vary greatly in pathogenicity. Treatment and outcome of neisserial infections are dictated by pathogenicity. For example, *N. meningitidis* is frequently found in the throat of normal humans, where it can reside as a commensal without causing symptoms. However, virulent strains of *N. meningitidis* may cause a fulminating meningitis, resulting in brain damage or death before treatment can control the disease. Diagnostic methods to date have not been useful in distinguishing the mild, commensal *Neisseria meningitidis* from the virulent strains.

The standard method of diagnosing a bacterial disease is culturing, followed by identification by immunoreactivity, morphology, and biochemical reactions. In the case of neisserial pathogens, because of its fastidious growth requirements, the organism has often lost viability and will no longer grow in culture once isolated from the patient. U.S. Pat. No. 4,446,230 discloses a test method and bacterial strain for the laboratory diagnosis of gonorrhea. This strain can be maintained in a laboratory and will become transformed by exogenous neisserial DNA, even from a non-viable clinical sample. The transformation corrects an induced specific growth requirement, thereby permitting the strain to grow. However, this test does not distinguish virulent from non-virulent strains.

In vivo models of *N. gonorrhoeae* infection using male human volunteers have been done to elucidate the factors that contribute to bacterial virulence. However, such studies are costly and limited in scope. Recent tissue culture models of gonococcal infection have begun to define the interactions between the bacterium and the host. In brief, it is evident that the bacteria attach and invade human urethral epithelial cells, a niche that probably represents the primary site of infection during the course of urethritis. Gonococci reside and replicate within vacuoles within these cells. Infected cells can rupture or be shed into the urethral lumen, releasing gonococci to invade neighboring epithelial cells or to be excreted in the urine. Several gonococcal components, including pili, Opa proteins and lipooligosaccharide (LOS) have been implicated in the ability of gonococcus to attach to and invade host cells, and implicated in the pathogenesis of gonococcal infection.

Our current knowledge clearly suggests that the gonococcus exists within different environments during the course of infection, such as extra- and intra-cellular and vacuolar locations. It is also likely that the site of infection of *N. gonorrhoeae*, the genital tract, differs significantly between males and females. How *N. gonorrhoeae* adapts to these different conditions is not well understood. It has been observed that a gonococcus thought to be pathogenic

quickly converts to a less virulent type under the conditions of culture that are used to identify the bacterium and to test for antibiotic sensitivity.

Current treatment of neisserial infections is with broad spectrum antibiotics. However, treatment with broad spectrum antibiotics leads to the disturbance of the natural microflora, leaving the patient susceptible to infections with such opportunistic pathogens as *Candida albicans* and *Gardnerella*. An antibiotic targeted specifically at pathogenic, rather than commensal *Neisseria*, would avoid this complication. It is currently difficult to test virulent clinical isolates of *Neisseria* to determine each strain's sensitivity to targeted-specificity antibiotics because of the difficulty of maintaining virulence during growth of the strains in culture. Therefore, such an antimicrobial sensitivity test as is disclosed in U.S. Pat. No. 5,789,173 may not be useful in determining neisserial sensitivity to antibiotics.

If the factors that convert a commensal, mildly or non-pathogenic *Neisseria meningitidis* or *N. gonorrhoeae* into a pathogenic, invasive bacterium were known, it would be possible to use the identification of such factors as an aid for the diagnosis and therapy of neisserial disease. There is a need to determine these so that they can be used as a target in screening bacteriostatic or bacteriocidal drugs that are selectively effective against virulent *Neisseria*.

BRIEF SUMMARY OF THE INVENTION

The present invention provides a mutant *Neisseria* bacterium having excessive outer membrane. The bacterium may be *Neisseria gonorrhoeae* or *Neisseria meningitidis*.

The present invention also provides a method of identifying a pathogenic bacterium comprising culturing under identical conditions 1) a mutant *Neisseria* bacterium having excessive outer membrane, 2) a wild type *Neisseria* and 3) a clinical sample taken from a patient suspected of having a neisserial infection; examining the extent of blebbing in each cultured bacterium; and comparing the extent of blebbing of the clinical sample to that of the bacterium having excessive outer membrane and to that of the wild type *Neisseria* so as to determine the pathogenicity of the clinical sample.

The present invention further provides a method of identifying an inhibitor compound of virulent *Neisseria* providing a culture containing a bacterium having a mutation which results in blebbing of the outer membrane and the test inhibitor compound, and providing a culture containing a wild type bacterium having no blebbing and the test inhibitor; and comparing the growth of the mutant bacterium to that of the wild type in order to identify an inhibitor compound that inhibits the growth of the mutant to a greater degree than the inhibitor compound inhibits the growth of the wild type.

The present invention further provides individual, isolated proteins coded for or otherwise under the control of the two-component system, the presence of which quickly and easily determined and is indicative of virulence.

DESCRIPTION OF THE DRAWINGS

FIG. 1 shows a diagram of the *bmr* locus of *N. gonorrhoeae* (NG), *N. meningitidis* (NM) and *N. cinerea* (NC).

FIG. 2 shows the alignment of the amino acid sequences of the *BmrR* open reading frames (ORF) from *N. gonorrhoeae* (GC) (SEQ ID NO:2, SEQ ID NO:3), *N. meningitidis* (NM) (SEQ ID NO:1) and *N. cinerea* (NC) (SEQ ID NO:4); and the alignment of the amino acid sequences of the

BmrS ORFs from GC (SEQ ID NO:6), NM (SEQ ID NO:5), and NC (SEQ ID NO:7).

FIGS. 3A–F show scanning electron micrographs of wild type and mutant *N. gonorrhoeae*.

FIGS. 4A–D shows scanning electromicrographs of wild type and mutant *N. meningitidis*.

DETAILED DESCRIPTION OF THE INVENTION

Neisseria gonorrhoeae and *N. meningitidis* are strictly human pathogens. Current knowledge of the pathogenesis of these organisms suggests that these bacteria encounter different environments within the host during infection. However, little is known of how *Neisseria* sense and respond to these different environments or what systems contribute to virulence. Many bacteria respond to changes in their environment through changes in the expression level of particular genes. Two-component regulatory systems are now recognized as significant mediators of signaling in bacteria, relaying environmental signals that produce changes in gene expression patterns. During signal transduction by a typical two-component regulator, a membrane sensor protein detects a specific environmental stimulus presumably either through a direct interaction with a ligand or through conformational changes induced by changes in environmental conditions. Such a regulatory system might be useful as a diagnostic marker of virulence in *Neisseria*.

However, information regarding signaling within the gonococcus is limited. Previously a two-component regulator-like system was reported in pilus gene expression, although the environmental cue to which the system responds is unknown. Since membrane blebbing has been found to be associated with virulence, it was thought that a quantitative assay of the amount of membrane would be indicative of increased virulence. In order to further investigate a possible two-component regulator-mediated signaling pathway in *N. gonorrhoeae*, the polymerase chain reaction (PCR) with degenerate oligonucleotide primers was used to amplify a region from the DNA-binding component of a novel *N. gonorrhoeae* two-component regulator.

This regulatory system is known to be present in other *Neisseria* species and to have a role in outer-membrane blebbing. The term “blebbing” is used to mean the bubble- or blister-like extrusion of membrane from the surface of the bacterial cell. The term “high-blebbing” is used to describe bacteria showing such extensive outer membrane formation. “Blebs” are easily separated from the outer membrane by mild physical manipulation, such as differential centrifugation.

Outer membrane bleb formation by *Neisseria meningitidis* and *N. gonorrhoeae* has long been recognized as an important factor in *Neisseria* infection, particularly that due to *N. meningitidis*. Numerous membrane blebs were found at the tissue surface during ultrastructural analysis of cervical gonorrhoea (Evans et al. *J. Infect. Dis.* (1977) 136:248–255) and a meningococcus recovered from the plasma of an infected patient contained multiple, long membranous protrusions typical of blebbing (Brandtzaes et al. *J. Infect. Dis.* (1989) 155:195–204). Lipooligosaccharide (LOS), which is a component of the blebs shed by the gonococcus and meningococcus, has major physiological effects during sepsis and meningitis caused by *N. meningitidis* and the plasma LOS levels are closely correlated to prognosis. Furthermore, compartmentalization of LOS production correlates with the clinical presentation in meningococcal infection. LOS levels in patients defined as having septicaemia showed high levels

in plasma (median 3500 ng/ml) and low levels in cerebral spinal fluid, while with patients with meningitis, LOS was detectable in the plasma of three out of 19 patients and in the CSF in 18 of 19 patients, with median levels of 2500 ng/ml.

From these observations, it is possible that blebbing is both the indicium and the cause of pathogenicity in *Neisseria*. Agents that control or inhibit blebbing may be useful in therapy against *Neisseria* infections. However, the standard laboratory cultures of *Neisseria* typically show little or no blebbing after repeated laboratory passage and are therefore not useful in screening for agents that control or inhibit blebbing. Identification of the degree of blebbing of newly isolated strains from clinical samples may be an early indication of the virulence of the pathogen and whether immediate, aggressive therapy is necessary.

Factors affecting the blebbing process are poorly understood. Strains of meningococcal serogroup A, B and C release membrane blebs in the log phase of growth but not in the lag phase, suggesting some form of regulation. The identification of a putative regulatory system, the mutation of which produces clear differences in the blebbing process is potentially of great interest to improvement of therapeutic strategies for treatment of *Neisseria* infections in which blebbing is a serious complication.

During the course of investigation, mutation of the genes controlling blebbing was successful, and after insertion into a strain, the resultant high-blebbing bacterium is a “super blebber” that shows extensive membrane blebbing in all culture conditions. Unlike virulent, high-blebbing clinical isolates of *Neisseria*, these super blebber mutants, termed bmrRS, are stable and do not revert to non-blebbing strains on repeated culture. For this reason, they are particularly useful as controls to estimate the virulence of clinical samples, for the production of neisserial membranes for use as a vaccine and for screening of antibiotics targeted at virulent *Neisseria*.

Outer membrane proteins and LOS are known to be useful as vaccines. (See, e.g., U.S. Pat. No. 5,902,586 issued May 11, 1999 to Jennings et al.) Because the super blebbers produce two to four times as much outer membrane as do wild type strains, they are a particularly useful source of *neisseria* vaccine material. Thus, a novel two-component regulatory system has been identified in *N. gonorrhoeae*, *N. meningitidis* and *N. cinerea*. DNA sequence analysis suggested that the system may be non-functional in laboratory strains of *Neisseria*. However, experimental mutation of the system in each of the three species resulted in increased membrane production in the mutant compared to the wild type. This correlated with an increase in outer membrane blebbing. These data suggest that outer membrane blebbing, which is regarded as an important contributor to the pathogenesis of neisserial diseases, is regulated as part of a coordinated response to environmental cues.

The references identified in the specification are incorporated herein by reference to the extent that they supplement, explain, provide a background for or teach methodology, techniques, and/or compositions employed herein.

The following examples are included to demonstrate the preferred embodiments of the invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples that follow represent techniques discovered by the inventors to function well in the practice of the invention, and thus can be considered to constitute preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments

that are disclosed and still obtain a like or similar result without departing from the concept and scope of the invention. More specifically, it will be apparent that certain agents, such as DNA constructs, that are both chemically and biologically equivalent may be substituted for the agents described herein while the same or equivalent results would be achieved. It will also be apparent that the techniques are not limited by the order in which the steps are carried out. All such substitutions and modifications apparent to those skilled in the art are deemed to be within the spirit and scope of the invention as defined by the appended claims.

The present invention provides a target for control of the neisserial membrane synthesis. The present invention also provides a quantitative assay of neisserial membrane. The present invention also provides a neisserial bacterium with increased amounts of membrane, more specifically, blebs of outer membrane, which bacterium is useful as a production cell for isolation of vaccine related membrane proteins and glycolipids. The present inventions also provides a standard with which to compare clinical isolates of *Neisseria* spp. in order to estimate the pathogenicity of such isolates. The present invention also provides a screening assay for drugs that are specifically effective against virulent strains of *Neisseria*.

The invention is illustrated by the following examples.

EXAMPLE 1

Construction of the Super Blebber

A. Bacterial strains and plasmids

Wild type *Neisseria* strains used were *N. gonorrhoeae* 1291, *N. meningitidis* NMB and *N. cinerea* 601. *Neisseria* were grown on solid GC Base medium (Difco Laboratories, Detroit) supplemented with amino acids and vitamins. For growth of kanamycin resistant *Neisseria*, BHI agar was used (Difco) supplemented with 2.5% FCS. Liquid *Neisseria* cultures were grown in proteose peptone broth (proteose peptone No. 3, Difco) 15 grams per liter; soluble starch (Difco) 1 gram per liter, dibasic potassium phosphate 4 grams per liter; monobasic potassium phosphate 1 gram per liter; sodium chloride 15 grams per liter or in Morse's defined medium.

Escherichia coli strain XL1-Blue (Stratagene, LaJolla, Calif.) was used to maintain plasmids and was grown in liquid culture using LB broth or on solid medium using LB agar. Antibiotics were used at the following concentrations: ampicillin 100 $\mu\text{g/ml}$; kanamycin 50 $\mu\text{g/ml}$ (*E. coli*) or 25 $\mu\text{g/ml}$ (*Neisseria*), erythromycin 150 $\mu\text{g/ml}$ (*E. coli*) or 1 $\mu\text{g/ml}$ (*Neisseria*).

pBluescript (Stratagene), pUC18 (Pharmacia, Piscataway, N.J.) and pCRII.I (Invitrogen, Carlsbad, Calif.) were used as cloning vectors. PUC4K (Pharmacia) was used as a source of kanamycin cassette. Erythromycin resistance cassette was derived from pKermC' plasmids (Zhou, D. G. and M. A. Apicella (1996) Gene 171:133-134.)

Plasmid DNA was prepared using Qiagen plasmid DNA preparation (Qiagen Inc., Chatsworth, Calif.). Chromosomal DNA was prepared from plate grown bacteria which were scraped from the plate into 100 μl of phosphate buffered saline (PBS). Cells were then pelleted by spinning for one minute at top speed in a microfuge before resuspension in 250 μl of TNE (10 mM Tris-HCl pH 8, 100 mM NaCl, 1 mM EDTA.) 25 μl of 10% SDS were added followed by 25 μl of Proteinase K (25 mg/ml). The reaction was incubated overnight at 55° C. and then extracted twice with phenol and then twice with phenol/chloroform. DNA was precipitated by addition of 10 μl 3M sodium acetate pH 5.4 and 1 ml 100% ethanol, lifted from the tube on the end of a pipette tip and washed by repeated transfer to tubes containing 500 μl 70% ethanol. Washed DNA samples were air dried before resuspension in 100 μl TE RNase.

DNA sequencing was performed at the University of Iowa DNA sequencing facility using dye terminator sequencing chemistry with AmpliTaq DNA polymerase, FS enzyme (PE Applied Biosystems, Foster City, Calif.). The reactions were run on and analyzed using an Applied Biosystems Model 373A stretch fluorescent automated sequencer.

PCR was performed using Boehringer Mannheim (Philadelphia) reagents according to the protocols provided. PCR-DOP was performed using degenerate oligonucleotides as described by Wren et al (FEMS Microbial Letters (1992) 99:287-291). PCR products were directly cloned using the TA cloning system (Invitrogen) according to the manufacturer's instructions.

B. Results of PCR amplification and cloning in *N. gonorrhoeae*

PCR using the degenerate oligonucleotides described by Wren produced a product of the expected size (322 bp) from a *N. gonorrhoeae* genomic DNA template. The product was cloned and numerous clones were analyzed through DNA sequencing. This revealed that the PCR product contained many different DNA fragments that were clearly not from two-component regulator genes, which is in contrast to the report in which the same primer sequences amplified a region of two-component systems from numerous other bacteria without this apparent non-specific effect.

A single clone of 322 bp was obtained which through analysis of its deduced amino-acid sequence clearly contained the 5' region of the DNA-binding component of a two-component regulatory system. This clone was not identical to the pilAB system of *N. gonorrhoeae*.

Using the 322 bp PCR product as a probe, several genomic libraries were screened, but no positive signals were observed, although a Southern blot of *N. gonorrhoeae* genomic DNA identified several restriction fragments to which the probe stringently hybridized. Attempts to clone these fragments, all of which were 4 kbp or greater in size, failed. These fragments, which might have contained the entire two-component system sought, could not be cloned, suggesting that the system was toxic to *E. coli* when carried at high copy number. However, two Sau3AI fragments which were identified by the Southern analysis were successfully cloned and found to contain open reading frames (ORFs) that encoded for the majority of a two-component regulator, based on their deduced amino acid sequence. Successive cloning of other small genomic fragments, containing small parts of the two-component regulator, followed by reconstruction, produced a sequence contig of 3.97 kbp. that contained the two ORFs of the putative two-component regulator and partial ORFs upstream and downstream of the system. We have named this two-component system bmr (for bacterial membrane regulator, see below). The upstream ORF was homologous to a hypothetical membrane protein of *Bacillus subtilis* (Genbank submission number P42308) and the downstream ORF was homologous to the cytoplasmic axial filament protein CafA, of *E. coli* (FIG. 1.) *N. gonorrhoeae* bmrRS has been deposited with the American Type Culture Collection (P.O. Box 1549, Manassas, Va., 20108) under accession number PTA-801.

Comparison of the deduced amino acid sequence of the putative two-component regulator to the amino acid sequences of the Genbank database revealed homology to numerous other two-component regulators. The highest degree of homology was to the BasRS system of *E. coli* and *Salmonella typhimurium* and the putative Bas homologue in *Haemophilus influenzae*. The Bas system of *E. coli* and *S. typhimurium* is involved in regulation of lipopolysaccharide (LPS) substitution by phosphate and aminoarabinose, that in the case of *S. typhimurium* is implicated in LPS changes during the course of infection. This degree of homology is frequently found in bacteria. Biochemical capabilities of the bacteria are modified and extended by the duplication and

subsequent mutation of genes, resulting in gene products with considerable degrees of homology, but having different functions and characteristics, as is seen in the *bas* and *bmr* gene products.

An interesting feature of *N. gonorrhoeae* *bmr* was the presence of a stop-codon at the fourteenth codon position of the DNA-binding component that was predicted to terminate translation of this ORF at that Point (FIG. 2.)

C. Cloning of *bmr* from other *Neisseria*

N. meningitidis is closely related to *N. gonorrhoeae* and *N. cinerea* is a non-pathogenic member of the *Neisseria* genus. Southern blots of genomic DNA from these two bacteria, using the *N. gonorrhoeae* 322 bp PCR product as a probe, identified stringently hybridizing restriction fragments in each of these organisms. Using a strategy similar to that described above, the *bmr* locus was cloned from *N. meningitidis* and *N. cinerea*. The predicted amino acid sequences of the three loci are highly homologous, strongly suggesting that the equivalent locus from the three species had been cloned (FIG. 2). However, in *N. meningitidis*, a T to G substitution altered the premature TAA stop-codon of *N. gonorrhoeae* to a GAA glutamate codon. In *N. cinerea*, the codon was also for glutamate, but read GAG. However, the *N. meningitidis* locus contains a 591 bp deletion compared to both *N. gonorrhoeae* and *N. cinerea* that results in deletion of the 3' 486 bp of the putative DNA-binding component-encoding ORF, the intergenic region and the 5' 69 bp of the putative sensor-encoding ORF (FIG. 1.) In addition, other nucleotide differences in the *N. meningitidis* locus immediately preceding the deletion cause frame shifts of the *BmrR* ORF (FIG. 2.) Thus, *N. meningitidis* is not predicted to encode a *BmrR* protein. Although the 5' 23 codons of the sensor-encoding ORF are deleted, an alternative initiation codon may allow translation of a near-full length sensor protein in *N. meningitidis*. The *bmr* locus was amplified by PCR from numerous *N. meningitidis* strains, including several from serotypes A, C and W 135. The primers used (5' CCTGTCCGTGTGTTGCAATC 3' (SEQ ID NO:8) and 5' CCGTTCCCGTTATCCTCCAC 3' (SEQ ID NO:9)) amplify the region starting 166 bp upstream of the initiating codon of the deleted *BmrR* ORF and ending 200 bp upstream of the termination codon of the putative *BmrS* ORF. All of these PCR reactions produced a band of the same size as that from strain NMB (i.e., approximately 600 bp smaller than the fragment amplified from *N. gonorrhoeae*) (data not shown) suggesting that the deletion in *bmr* is common to many, if not all, *N. meningitidis* strains. *N. meningitidis* *bmrRS* has been deposited with the American Type Culture Collection (P.O. Box 1549, Manassas, Va., 20108) under accession number PTA-800.

An *N. gonorrhoeae* *bmrS* insertion mutant was constructed by insertion of an erythromycin resistance cassette at the site indicated in FIG. 1. An insertion mutant of *N. meningitidis* was constructed by insertion of a kanamycin resistance cassette within the sensor coding region (FIG. 1.) The locus was mutated in *N. cinerea* by deletion of an internal *BsrGI* fragment and insertion of a kanamycin resistance cassette in its place. This mutation deletes the 3' 418 bp of *bmrR* and the 5' 565 bp of *bmrS* (FIG. 1).

FIG. 2 shows the alignment of the amino acid sequences of the *BmrR* and *BmrS* ORFs from the three *Neisseria* spp. The GC *BmrR* ORF contains a stop codon (*) at the fourteenth codon. The NM *BmrR* sequence contains a small deletion compared to the GC and NC sequences after the 34th codon, resulting in a frameshift (denoted by the amino acid sequence written on the line above). The amino acid sequences in frame with the N-terminal NM *BmrR* ORF is closed by the presence of a stop codon (*). Following the frameshift, a small insertion results in the presence of the amino acids FVPLA in the NM sequence that is absent from both GC and NC. Following this is a large deletion that

removes the remainder of the *BmrR* ORF and the N-terminal 23 codons of the *BmrS* ORF. Outside of this region, the three sequences are highly homologous.

EXAMPLE 2

5 Characteristics of the Super Blebbers

A. Chemical analysis of the outer membrane of the super blebbers

The effect of the *bmr* mutation on several parameters was investigated by comparison of the *N. gonorrhoeae* wild-type and the mutant. Overnight cultures (10 ml) were killed by the addition of 0.5% phenol for two hours and outer membrane enriched samples were isolated as described by Zollinger et al., International Symposium, Hanasaari, Espoo, Finland (1991). Samples were analyzed by SDS-PAGE on a 6–15% gradient gel. LOS was isolated by phenol/chloroform/petroleum ether extraction as described by Galanos et al. Eur. J. Biochem. (1969) 9:45–249 and analyzed as described by Lee et al. J. Biol. Chem. (1995) 270:271151–9. The growth rate and outer membrane protein profile of the mutant were unaltered from the wild type.

20 Because of the homology of the *Neisseria* locus to the previously characterized *bas* locus of *E. coli* and *S. typhimurium* and the involvement of this system in LPS modulation in those bacteria, LOS was isolated from wild type and mutant and analyzed by mass spectrometry. No difference was found, showing that despite the homology, the system functioned differently in *Neisseria* and those other bacteria.

B. Morphology of the super blebbers

30 For scanning electron microscopy (SEM), samples of proteose peptone broth cultures were taken directly without centrifugation or washing. Processing included treatment with 1% osmium tetroxide prior to dehydration through a graded ethanol series, with a final clearance in hexamethyldisilane (HMDS from Polysciences, Inc. Warrington, Pa.). After coating with gold-palladium, the specimens were viewed on an S-4000 Hitachi scanning electron microscope at 5 kV accelerating voltage.

For transmission electron microscopy, the samples were dehydrated in a graded ethanol series prior to embedment in LR White resin (Ted Pella, Inc., Redding, Calif.) and sectioned to approximately 85 nm thickness using an ultramicrotome. Samples were counter-stained with 5% uranyl acetate for viewing with an H-7000 Hitachi transmission electron microscope at 75 kV accelerating voltage.

SEM analysis revealed that the wild type and mutant bacteria differed in two ways. Wild type *N. gonorrhoeae* were observed to be distinct diplococci with a smooth appearance (FIGS. 3A and 3B) However, the mutant produced large numbers of blebs. (FIGS. 3C and 3D) Shedding of outer-membrane particles (blebbing) is a characteristic of *Neisseria*. In these analyses the mutant bacteria were blebbing to a much higher degree than the wild type. In addition, many mutant diplococci were joined to others by membranous connections (FIGS. 3C and 3E). When viewed by TEM, the connections did not contain electron dense material of the cytoplasm, suggesting that they consist of hollow membrane tubes joining pairs of diplococci. (FIG. 3F). These connections were not observed with the wild type.

The morphological changes seen in the *N. meningitidis* *bmr* mutant are shown in FIG. 4A. FIGS. 4A and 4B show the parent strain of *N. meningitidis* NMB. FIG. 4C and FIG. 4D show the *N. meningitidis* NMB*bmr* mutant. As observed with the mutant *N. gonorrhoeae*, these mutants were also joined by membranous connections. The *bmr* mutant of *N. cinerea* showed similar changes in morphology.

65 C. Phospholipid Analysis

The electron micrographs suggested that the mutants had alterations in their membranes that resulted in an increase in

membrane material. To test this hypothesis, wild type and mutant *N. gonorrhoeae* were grown in the presence of ^{14}C -labeled acetate to label phospholipid and allow quantitation of membrane produced by the bacteria.

Bacteria were grown in 5 ml of Morse's defined medium supplemented with $4\ \mu\text{Ci/ml}$ ^{14}C -labeled acetate. *N. meningitidis* was grown for 5 hours and *N. gonorrhoeae* was grown for 20 hours at 37°C . in a shaking incubator to achieve the appropriate bacterial growth. For *N. gonorrhoeae*, the blebs were separated from the bacteria by underlying the cultures with 2.5 ml of 10% sucrose in balanced salt buffer in centrifuge tubes and centrifuging in a SW-40 Ti rotor (Beckman, Palo Alto, Calif.) at 10,000 rpm at 4°C . for 30 minutes. The supernatant above the sucrose cushion was transferred to fresh centrifuge tubes. The sucrose cushion was removed for scintillation counting. The supernatant was centrifuged in a SW-40 Ti rotor at 25,000 rpm ($100,000\times g$) for 75 minutes at 4°C . to pellet blebs. The supernatant from this spin was removed for scintillation counting. The centrifuge tube was partially dried using cotton swabs.

Labeled phospholipids were isolated by extraction with $100\ \mu\text{l}$ of chloroform and $200\ \mu\text{l}$ of methanol added to the sample and vortexed for 2 minutes. A further $100\ \mu\text{l}$ of chloroform was added followed by 30 seconds of vortexing. $100\ \mu\text{l}$ of deionized water was added followed by 30 seconds of vortexing. The mixture was centrifuged at 13,000 rpm in a microfuge for ten minutes at room temperature. The upper layer was removed for scintillation counting. The lower layer was removed from below the interface pellet and centrifuged before scintillation counting.

Blebs were separated from whole cells and the phospholipids in both samples extracted. The counts present in both bleb and cell-associated phospholipids are presented in Table I.

TABLE I

^{14}C -acetate Incorporation, Counts per Minute <i>N. gonorrhoeae</i> strain 1291, wild type and bmr mutant			
	Experiment 1	Experiment 2	Experiment 3
<u>Wild Type</u>			
Total CPM	7.4×10^7	5.7×10^7	7.1×10^7
CPM in cell-associated phospholipids	5.7×10^6	4×10^6	4.8×10^6
CPM in bleb-associated phospholipids	2.5×10^4	8.1×10^4	5×10^4
<u>Mutant</u>			
Total CPM	6.8×10^7	6.7×10^7	6.8×10^7
CPM in cell-associated phospholipids	7.3×10^6	4.8×10^6	5.6×10^6
CPM in bleb-associated phospholipids	5.2×10^4	3.1×10^5	2.1×10^6
Ratio of mutant to wild type cell-associated CPM	1.3	1.2	1.2
Ratio of mutant to wild type bleb-associated CPM	2.1	3.8	4.2

In each of these three experiments, *N. gonorrhoeae* 1291 bmr incorporated increased amounts of label compared to wild type. Both cell-associated and bleb-associated label was increased in the mutants. The increase in cell-associated counts was slight, which agrees with microscopy observations in which the mutant and wild type bacteria did not appear different except for the presence of connections between mutant diplococci and increased level of blebbing in the mutant. As would be expected from the morphology,

mutant bleb-associated counts were between two-and four-fold higher than wild type.

Labeling experiments were also conducted using *N. meningitidis*. For unknown reasons, it proved difficult to obtain consistent bleb-associated counts with these bacteria. The micrographs suggested that the membrane alteration observed with these mutants did not involve the type of blebbing seen in the gonococcus, but instead involved cell-associated membrane changes. However, total membrane was increased as was seen in the *N. gonorrhoeae* experiments. Table II shows a comparison of total outer membrane counts from both wild type and mutant bacteria. Increased incorporation of counts in the mutants was observed.

TABLE II

^{14}C -acetate Incorporation, Counts per Minute Wild type and bmr mutant of <i>N. meningitidis</i> NMB		
	Experiment 1	Experiment 2
<u>Wild-Type</u>		
Total CPM	4.0×10^8	6.0×10^7
Total phospholipids	1.0×10^7	5.4×10^6
<u>Mutant</u>		
Total CPM	4.0×10^8	6.0×10^7
Total phospholipids	1.9×10^7	9.7×10^6
Ratio of mutant to wild-type CPM	1.9	1.8

When larger quantities of membrane are desired, the culturing and purification may be easily scaled up.

EXAMPLE 4

Use of the Neisseria bmr Mutants to Estimate Virulence of Bacteria Recovered from Clinical Samples.

Swabs from suspected Neisseria infections are cultured for meningococcus or gonococcus by the methods disclosed in Example 1. A laboratory strain of Neisseria and a super blebber are grown in the same manner as controls. Overnight cultures are examined for blebbing in the SEM as described in Example 2B.

Phospholipids are isolated as described in Example 1. Phospholipids are quantitated as described by the incorporation of radionuclide or by HPLC analysis of total phospholipids. A level of phospholipid close to that of the super blebber indicates infection with a virulent neisserial pathogen, indicating that therapy should be immediate and aggressive.

As seen in Table III, a quantitative estimate on a scale of 1 (no blebbing) to 3 (many blebs and filamentous connections between cells) and the phospholipid content that correlates with blebbing are made to determine virulence of the clinical sample.

TABLE III

	Blebbing	Phospholipid CPM*	Diagnosis
<i>N. gonorrhoeae</i> strain 1291	1	6×10^4	
Super blebber	3	2×10^5	
Hypothetical Sample #1	1	6×10^4	Not virulent, therapy not needed
Hypothetical Sample #2	2	1×10^5	Moderate virulence, therapy indicated
Hypothetical Sample #3	3	$>1.8 \times 10^5$	Highly virulent, immediate and

TABLE III-continued

	Phospholipid Blebbing CPM*	Diagnosis
		aggressive therapy indicated

*Phospholipid content is expressed as counts per minute ¹⁴C-acetate incorporation into phospholipid fraction.

EXAMPLE 5

Use of super blebbers to screen biocidal or biostatic agents.

A ninety-six well culture is set up with *N. gonorrhoeae* Strain 1291 and the *N. gonorrhoeae* super blebber, and the *N. meningitidis* strain NMB and the *N. meningitidis* super blebber. Dilutions of the potentially selective antibiotics are added to respective wells. An antibiotic is considered selective if it inhibits the mutant organisms that show excessive blebbing as indicated by no growth in the super blebber chambers at lower concentrations than those that inhibit growth in the strains 1291 and/or NMB.

An compound may affect a specific property of the bacterium without outright killing of the organism. Such a compound may be particularly useful in that it does not disturb the normal microflora of a patient while inhibiting the specific property that renders a pathogen virulent. It would be particularly useful to identify a compound that

targets the Neisseria blebbing process. A ninety-six well culture is set up with a recently isolated, high-blebbing Neisseria that shows excessive outer membrane formation as indicated by blebbing. Dilutions of the potentially selective antibiotics are added to respective wells. An antibiotic is considered selective if it inhibits the blebbing at lower concentrations than those that inhibit growth. Such an antibiotic may be considered to have converted the high-blebbing, virulent bacterium to a non-virulent form.

All of the compositions and methods disclosed and claimed herein can be made and executed without undue experimentation in light of the previous disclosure. While the compositions and methods of this invention have been described in terms of preferred embodiments, it will be readily appreciated by those of skill in the art that many variations and modifications may be applied to the compositions and methods and used in the steps or in the sequence of steps of the methods described herein without departing from the concept and scope of the invention. More specifically, it will be apparent that certain agents, such as vectors, primers and bacterial strains, that are both chemically and physiologically equivalent may be substituted for the agents described herein while the same or similar results would be achieved. All such similar variations and modifications apparent to those of skill in the art are deemed to be within the scope and concept of the invention defined by the appended claims.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 9

<210> SEQ ID NO 1
<211> LENGTH: 73
<212> TYPE: PRT
<213> ORGANISM: Neisseria meningitidis

<400> SEQUENCE: 1

Met Arg Val Leu Leu Val Glu Asp Asp Ala Met Ile Ala Glu Ala Val
1 5 10 15

Ser Ala Ser Leu Lys Asp Gly Gly Tyr Ala Val Asp Trp Val Lys Asn
20 25 30

Gly Ala Arg Leu Pro Leu Pro Ser Leu Met Thr Ala Ala Ala Ala
35 40 45

Gln Pro Tyr Asp Leu Met Leu Leu Asp Leu Gly Leu Leu Gly Arg Asp
50 55 60

Gly Leu Asp Val Met Phe Val Pro Leu
65 70

<210> SEQ ID NO 2
<211> LENGTH: 13
<212> TYPE: PRT
<213> ORGANISM: Neisseria gonorrhoeae

<400> SEQUENCE: 2

Met Arg Val Leu Leu Val Glu Asp Asp Ala Met Ile Ala
1 5 10

<210> SEQ ID NO 3
<211> LENGTH: 210
<212> TYPE: PRT
<213> ORGANISM: Neisseria gonorrhoeae

-continued

<400> SEQUENCE: 3

Ala Val Ser Ala Ser Leu Lys Asp Gly Gly Tyr Ala Val Asp Trp Val
 1 5 10 15
 Lys Asn Gly Ala Gln Val Ala Ala Ala Ala Ala Gln Pro Tyr Asp
 20 25 30
 Leu Met Leu Leu Asp Leu Gly Leu Pro Gly Arg Asp Gly Leu Asp Val
 35 40 45
 Leu Ser Glu Ile Arg Ala Ala Gly Cys Thr Val Pro Val Leu Ile Val
 50 55 60
 Thr Ala Arg Asp Asp Leu Tyr Ser Arg Leu Asn Gly Leu Asp Gly Gly
 65 70 75 80
 Ala Asp Asp Tyr Ile Val Lys Pro Phe Asp Met Ala Glu Phe Lys Ala
 85 90 95
 Arg Met Arg Ala Val Leu Arg Arg Gly Ser Gly Gln Ala Gln Ala Cys
 100 105 110
 Leu Ser Asn Gly Ala Leu Ser Leu Asn Pro Ala Thr Tyr Gln Val Glu
 115 120 125
 Ile Ile Ala Glu Gly Arg Gln Val Ala Leu Ser Asn Gln Glu Phe Ser
 130 135 140
 Val Leu Gln Ala Leu Leu Ala Arg Pro Gly Val Ile Leu Ser Arg Ser
 145 150 155 160
 Asp Ser Glu Asp Lys Val Tyr Gly Trp Gly Gly Glu Val Glu Ser Asn
 165 170 175
 Ala Val Asp Phe Leu Ile His Gly Leu Cys Lys Lys Leu Gly Lys Glu
 180 185 190
 Ser Ile Gln Asn Val Arg Gly Val Gly Trp Leu Met Pro Arg Gln Asp
 195 200 205
 Ala Val
 210

<210> SEQ ID NO 4

<211> LENGTH: 222

<212> TYPE: PRT

<213> ORGANISM: *Neisseria cinerea*

<400> SEQUENCE: 4

Met Arg Val Leu Leu Val Glu Asp Asp Ala Met Ile Ala Glu Ala Val
 1 5 10 15
 Ser Ala Asn Leu Lys Asp Gly Gly Tyr Ala Val Asp Trp Val Lys Asn
 20 25 30
 Gly Ala Gln Val Ala Ala Ala Ala Ala Gln Pro Tyr Asp Leu Met
 35 40 45
 Leu Leu Asp Leu Gly Leu Pro Gly Arg Asp Gly Leu Asp Val Leu Ser
 50 55 60
 Glu Ile Arg Ala Ala Gly Cys Thr Val Pro Val Leu Ile Val Thr Ala
 65 70 75 80
 Arg Asp Asp Leu Tyr Ser Arg Leu Asn Gly Leu Asp Gly Gly Ala Asp
 85 90 95
 Asp Tyr Ile Val Lys Pro Phe Asp Met Ala Glu Leu Lys Ala Arg Met
 100 105 110
 Arg Ala Val Leu Arg Arg Gly Ser Gly Gln Ala Gln Ala Cys Leu Ser
 115 120 125
 Asn Gly Ala Leu Ser Leu Asn Pro Ala Thr Tyr Gln Val Glu Ile Ile
 130 135 140

-continued

Ala Glu Arg Arg Gln Val Ala Leu Ser Asn Lys Glu Phe Ala Val Leu
 145 150 155 160

Gln Ala Leu Leu Ala Arg Pro Gly Val Ile Leu Ser Arg Ser Asp Leu
 165 170 175

Glu Asp Lys Ile Tyr Gly Trp Gly Ala Glu Val Glu Ser Asn Ala Val
 180 185 190

Asp Phe Leu Ile His Ala Leu Arg Lys Lys Leu Gly Lys Glu Asn Ile
 195 200 205

Gln Asn Val Arg Gly Val Gly Trp Leu Val Pro Gly Ala Val
 210 215 220

<210> SEQ ID NO 5
 <211> LENGTH: 416
 <212> TYPE: PRT
 <213> ORGANISM: Neisseria meningitidis

<400> SEQUENCE: 5

Met Phe Val Pro Leu Ala Met Leu Ala Gly Met Phe Ser Tyr Tyr Glu
 1 5 10 15

Thr Phe His Glu Thr Glu Ala Leu Gln Asp Asp Leu Leu Arg Gln Ala
 20 25 30

Ala Leu Tyr Val Ala Pro Asp Ser Lys Pro Glu Thr Leu Pro Glu Gly
 35 40 45

Asp Gly Asp Thr Arg Ile Phe Val Gln Met Pro Gln Gln Glu Asp Pro
 50 55 60

Val Val Ser Leu Pro Ala His Leu Ala Asp Gly Leu His Thr Leu Gln
 65 70 75 80

Ala Asp Gly Asp Asp Tyr Tyr Arg Val Tyr Ile Arg Thr Thr Glu
 85 90 95

Gln Gly Arg Ile Ala Val Met Gln Glu Asn Glu Tyr Arg Glu Asp Leu
 100 105 110

Ala Ala Asp Ala Ala Arg Gln Ser Val Leu Pro Leu Leu Ala Ala Leu
 115 120 125

Pro Leu Met Ile Leu Leu Thr Val Trp Ile Thr His Lys Ala Met Arg
 130 135 140

Pro Val Arg Lys Leu Ser Gln Ser Leu Glu Gln Arg Arg Ile Asn Asp
 145 150 155 160

Leu Ser Ala Leu Ser Val Asp Asn Ile Pro Ser Glu Ile Arg Gly Phe
 165 170 175

Val Thr Ala Ile Asn Leu Leu Leu Lys Arg Ala Asp Glu Asp Ile Arg
 180 185 190

His Arg Gln Arg Phe Val Ala Asp Ala Ala His Glu Leu Arg Thr Pro
 195 200 205

Met Thr Ala Leu Ser Leu Gln Ala Glu Arg Leu Asn Asn Met Ser Leu
 210 215 220

Pro Pro Asp Ala Ala Arg Gln Pro Ala Val Leu Gln Gln Ser Ile Arg
 225 230 235 240

Arg Asn Lys His Leu Leu Glu Gln Leu Leu Ala Leu Ala Arg Ser Gln
 245 250 255

Ser Asp Glu Thr Pro Leu Thr Lys Thr Thr Phe Gly Leu Gln Ser Arg
 260 265 270

Phe Arg Gln Val Leu Gln Glu Leu Met Pro Leu Ala Leu Glu Lys Arg
 275 280 285

Gln Asp Ile Gly Val Ala Val Gly Gly Asp Val Glu Val Ser Ala Asp
 290 295 300

-continued

Glu Thr Glu Ile Tyr Thr Leu Ile Lys Thr Phe Ala Asp Asn Ala Val
 305 310 315 320
 Arg Tyr Thr Pro Pro Glu Gly Arg Ile Asp Leu Gly Phe Thr Asp Glu
 325 330 335
 Gly Lys Tyr Leu Ala Val Trp Val Glu Asp Asn Gly Asn Gly Ile Pro
 340 345 350
 Glu Ser Glu Arg Ala Arg Val Leu Asp Pro Phe Tyr Arg Ile Leu Gly
 355 360 365
 Thr Glu Gln Gln Gly Thr Gly Leu Gly Leu Ser Ile Ala Asp Thr Leu
 370 375 380
 Ala Lys Lys Tyr Gly Gly His Leu Glu Leu Thr Asp Ser Arg Arg Phe
 385 390 395 400
 Gly His Gly Leu Leu Ile Arg Ala Leu Leu Asp Lys Glu Thr Leu Lys
 405 410 415

<210> SEQ ID NO 6

<211> LENGTH: 439

<212> TYPE: PRT

<213> ORGANISM: Neisseria gonorrhoeae

<400> SEQUENCE: 6

Met Pro Asp Arg Phe Phe Lys Ile Leu Lys His Ser Leu Gln Val Arg
 1 5 10 15
 Ile Ser Leu Ala Leu Ile Trp Met Phe Val Pro Leu Ala Met Leu Ala
 20 25 30
 Gly Met Phe Ser Tyr Tyr Glu Thr Phe His Glu Thr Glu Ala Leu Gln
 35 40 45
 Asp Asp Leu Leu Arg Gln Ala Ala Leu Tyr Val Gly Pro Asp Ser Lys
 50 55 60
 Ser Glu Thr Leu Pro Glu Gly Asp Gly Asp Thr Arg Ile Leu Val Gln
 65 70 75 80
 Met Pro Gln Gln Glu Asp Pro Val Val Ser Leu Pro Ala His Leu Ala
 85 90 95
 Asp Gly Leu His Thr Leu Arg Ala Asp Gly Asp Asp Asp Tyr Tyr Arg
 100 105 110
 Ala Tyr Ile Arg Thr Thr Glu Gln Gly Arg Ile Ala Val Met Gln Glu
 115 120 125
 Asn Glu Tyr Arg Glu Asp Leu Ala Glu Asp Ala Ala Arg Gln Ser Val
 130 135 140
 Leu Pro Leu Leu Ala Ala Leu Pro Leu Met Ile Leu Leu Thr Val Trp
 145 150 155 160
 Ile Thr His Lys Ala Met Arg Pro Val Arg Lys Leu Ser Gln Ser Leu
 165 170 175
 Glu Gln Arg Arg Ile Asn Gly Leu Pro Ala Leu Ser Val Tyr Asn Ile
 180 185 190
 Pro Ser Glu Ile Arg Gly Phe Val Thr Ala Ile Asn Leu Leu Lys
 195 200 205
 Arg Val Asp Glu Asp Ile Arg Arg Arg Gln Arg Phe Val Ala Asp Ala
 210 215 220
 Ala His Glu Leu Arg Thr Pro Met Thr Ala Leu Ser Leu Gln Ala Glu
 225 230 235 240
 Arg Leu Asn Asn Met Pro Leu Pro Pro Asp Ala Gly Arg Gln Ser Ala
 245 250 255
 Val Leu Gln Gln Ser Ile Arg Arg Asn Lys His Leu Leu Glu Gln Leu

-continued

260					265					270					
Leu	Ala	Leu	Ala	Arg	Ser	Gln	Ser	Asp	Glu	Pro	Pro	Leu	Ala	Lys	Thr
	275						280					285			
Thr	Phe	Gly	Leu	Gln	Ser	Arg	Phe	Arg	Gln	Val	Leu	Gln	Glu	Leu	Met
	290					295					300				
Pro	Leu	Val	Leu	Glu	Lys	Arg	Gln	Asp	Ile	Gly	Val	Ala	Val	Gly	Gly
	305				310					315					320
Asp	Val	Glu	Val	Ser	Ala	Asp	Glu	Thr	Glu	Ile	Tyr	Thr	Leu	Val	Lys
				325					330					335	
Thr	Phe	Ala	Asp	Asn	Ala	Val	Arg	Tyr	Thr	Pro	Asn	Gly	Gly	Arg	Ile
			340					345					350		
Asp	Leu	Gly	Phe	Thr	Asp	Glu	Gly	Lys	Tyr	Leu	Ala	Val	Trp	Val	Glu
		355					360					365			
Asp	Asn	Gly	Asn	Gly	Ile	Pro	Glu	Ser	Glu	Cys	Ala	Arg	Val	Leu	Asp
	370					375					380				
Pro	Phe	Tyr	Arg	Ile	Leu	Gly	Thr	Glu	Gln	Gln	Gly	Ala	Gly	Leu	Gly
	385				390					395					400
Leu	Ser	Ile	Ala	Asp	Thr	Leu	Ala	Lys	Lys	Tyr	Gly	Gly	Tyr	Leu	Glu
				405					410					415	
Leu	Thr	Asp	Ser	Arg	Arg	Phe	Gly	His	Gly	Leu	Leu	Ile	Arg	Ala	Leu
			420					425					430		
Leu	Asp	Lys	Glu	Thr	Leu	Lys									
	435														
<210> SEQ ID NO 7															
<211> LENGTH: 378															
<212> TYPE: PRT															
<213> ORGANISM: Neisseria cinerea															
<400> SEQUENCE: 7															
Met	Leu	Asn	Arg	Leu	Ile	Arg	Met	Leu	Lys	Gln	Ser	Leu	Gln	Val	Arg
	1			5					10					15	
Ile	Cys	Ile	Ala	Leu	Ile	Leu	Met	Phe	Leu	Pro	Leu	Ala	Met	Val	Ala
			20					25					30		
Gly	Val	Phe	Ser	Tyr	Tyr	Asp	Thr	Phe	His	Glu	Ala	Glu	Glu	Leu	Gln
		35					40					45			
Asp	Asp	Leu	Leu	Arg	Gln	Thr	Ala	Leu	Tyr	Val	Gly	Pro	Asp	Tyr	His
	50					55					60				
Pro	Asp	Ala	Leu	Pro	Glu	Gly	Asp	Gly	Asp	Thr	Arg	Ile	Leu	Val	Gln
	65				70					75				80	
Met	Pro	Asp	Gln	Glu	Pro	Ile	Val	Ser	Leu	Pro	Met	His	Leu	Lys	Asp
			85						90					95	
Gly	Leu	His	Thr	Leu	Arg	Ala	Asp	Glu	Asp	Asp	Asp	Tyr	Tyr	Arg	Val
			100					105					110		
Tyr	Ile	Arg	Thr	Thr	Gly	Arg	Gly	Arg	Ile	Ala	Val	Met	Gln	Glu	Asn
			115				120					125			
Glu	Tyr	Arg	Glu	Asp	Leu	Ala	Ala	Asp	Ala	Ala	Met	Gln	Ser	Val	Leu
	130					135					140				
Pro	Leu	Leu	Ala	Ala	Leu	Pro	Leu	Ile	Ile	Leu	Leu	Thr	Val	Trp	Ile
	145				150					155					160
Thr	His	Gln	Ala	Met	Arg	Pro	Val	Arg	Ile	Leu	Ser	Gln	Asn	Leu	Glu
				165					170					175	
Gln	Arg	Arg	Leu	Asp	Asp	Leu	Ser	Ala	Leu	Asn	Thr	Asp	Asn	Ile	Pro
			180					185						190	

-continued

Ser Glu Ile Lys Gly Phe Val Thr Ala Ile Asn Leu Leu Leu Lys Arg
 195 200 205

Val Asp Glu Asp Ile Arg Arg Arg Gln Arg Phe Ile Ala Asp Ala Ala
 210 215 220

His Glu Leu Arg Thr Pro Met Thr Ala Leu Ser Leu Gln Ala Glu Arg
 225 230 235 240

Leu Asn Asn Met Pro Leu Pro Pro Asp Ala Ala Arg Gln Ser Ala Val
 245 250 255

Leu Gln Gln Ser Ile Arg Arg Asn Lys His Leu Leu Glu Gln Leu Leu
 260 265 270

Ala Leu Ala Arg Ser Gln Ser Asp Glu Thr Pro Leu Thr Lys Thr Thr
 275 280 285

Phe Gly Leu Gln Ser Arg Phe Arg Arg Val Leu Gln Glu Leu Met Pro
 290 295 300

Leu Ala Leu Glu Lys Arg Gln Asp Ile Gly Val Ala Val Glu Gly Asp
 305 310 315 320

Phe Glu Ile Phe Ala Asp Glu Thr Glu Ile Tyr Thr Leu Val Lys Thr
 325 330 335

Phe Val Asp Asn Ala Ile Arg Tyr Thr Pro Ser Glu Gly Arg Ile Asp
 340 345 350

Leu Gly Phe Thr Asp Glu Gly Lys Tyr Leu Ala Val Trp Val Glu Asp
 355 360 365

Asn Gly Lys Gly Ile Pro Glu Ser Glu Arg
 370 375

<210> SEQ ID NO 8
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Neisseria gonorrhoeae

<400> SEQUENCE: 8

cctgtccgtg tgttgcaatc 20

<210> SEQ ID NO 9
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Neisseria gonorrhoeae

<400> SEQUENCE: 9

ccgttccgt taccctccac 20

We claim:

1. A *Neisseria* bacterium comprising a mutant two-component regulatory (2CR) system that controls superblebbing, wherein said system comprises DNA encoding a BmrR open reading frame (ORF) and a BmrS ORF, wherein the BmrR ORF or the BmrS ORF contains an insertion or a deletion as compared to corresponding wild-type BmrR ORF or BmrS ORF.

2. The bacterium of claim 1 which is *Neisseria gonorrhoeae*.

50 3. The bacterium of claim 1 which is *Neisseria meningitidis*.

4. The bacterium of claim 2 which has been deposited with the American Type Culture Collection under accession number PTA-801.

55 5. The bacterium of claim 3 which has been deposited with the American Type Culture Collection under accession number PTA-800.

* * * * *