



US006803191B2

(12) **United States Patent**
Srikantha et al.

(10) **Patent No.:** **US 6,803,191 B2**
(45) **Date of Patent:** **Oct. 12, 2004**

(54) **CANDIDA ALBICANS TWO-COMPONENT HYBRID KINASE GENE, CANIK1, AND USE THEREOF**

FOREIGN PATENT DOCUMENTS

WO 96/40939 12/1996

(75) Inventors: **Thyagarajan Srikantha**, Coralville, IA (US); **David R. Soll**, Iowa City, IA (US)

OTHER PUBLICATIONS

(73) Assignee: **University of Iowa Research Foundation**, Iowa City, IA (US)

David R. Soll, "Gene regulation during high-frequency switching in *Candida albicans*" Microbiology, vol. 143, 1997, pp 279-288, XP002083237.

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

Database Swiss-Prot Accession No. p46588, Jun. 15, 1995 Ball T and Rosamond J: XP002083293 DNA Polymerase III gene (po13) from *Candida albicans*.

(21) Appl. No.: **09/424,951**

Nagahashi et al., Isolation of CaSLNI and CaNIK1, the genes for osmosensing histidine kinase homologues, from the pathogenic fungus *Candida albicans*: Microbiology, vol. 144, 1998, pp 425-432, XP002083238.

(22) PCT Filed: **Jun. 5, 1998**

Srikantha et al., The WH11 gene of *Candida albicans* is regulated in two distinct developmental programs through the same transcription activation sequences: Journal Of Bacteriology, vol. 179, No. 12, 1997, pp 3837-3844, XP002083239.

(86) PCT No.: **PCT/US98/11658**

§ 371 (c)(1),
(2), (4) Date: **Jan. 20, 2000**

Srikantha et al., "The sea pansy *Renilla reniformis* luciferas serves as a sensitive bioluminescent reporter, for differential gene expression in *Candida albicans*" Journal of Bacteriology, vol. 178, No. 1, 1996 pp 121-129, XP00208236.

(87) PCT Pub. No.: **WO98/55654**

PCT Pub. Date: **Dec. 10, 1998**

Timerlake, W.E., "Cellular Reporters for Antifungal Drug Discovery", PAP Conference Discovery Mode Action Antifungal Agent, 1995, pp 17-29 XP000603570.

(65) **Prior Publication Data**

US 2002/0137034 A1 Sep. 26, 2002

* cited by examiner

Related U.S. Application Data

(60) Provisional application No. 60/048,914, filed on Jun. 6, 1997.

Primary Examiner—James Ketter

Assistant Examiner—David Lambertson

(74) *Attorney, Agent, or Firm*—Foley & Lardner, LLP

(51) **Int. Cl.**⁷ **C12Q 1/68; C07H 21/04**

(57) **ABSTRACT**

(52) **U.S. Cl.** **435/6; 435/252.3; 435/254.22; 536/23.1; 536/24.1**

A *Candida albicans* gene, CaNik1, is involved in phenotypic switching which is significant because of a direct correlation between the switching and the level of virulence of the organism. A method of screening for anti-fungal pharmaceutical candidates entails bringing a test substance into contact with cells containing a CaNik1 gene or a variant thereof and then monitoring the effect, if any, on the level of expression of the gene.

(58) **Field of Search** **536/23.1, 24.1; 435/6, 252.3, 254.22**

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,939,306 A * 8/1999 Alex et al. 435/252.3

15 Claims, 10 Drawing Sheets

FIG. 1A

1531 Slb1 →
GAGATTAGAACACCCATTGAATGGGATTATTGGWATGACYCAGTTGTCRCTTGATACAGAG 1590
GluIleArgThrProLeuAsnGlyIleIleGlyMetThrGlnLeuSerLeuAspThrGlu 530
H1
TTGACRCAGTACCAACGAGAGATGTTGTCGATTGTGCATAACTTGGCAAATTCCTTGTTG 1650
LeuThrGlnTyrGlnArgGluMetLeuSerIleValHisAsnLeuAlaAsnSerLeuLeu 550

ACCATTATAGACGATATATTGGATATTTCTAAGATTGAGGCGAATAGAATGACGGTGGAA 1710
ThrIleIleAspAspIleLeuAspIleSerLysIleGluAlaAsnArgMetThrValGlu 570

CAGATTGATTTTTTCATTAAGAGGGACAGTGTGGTGCATTGAAAACGTTAGCCGTCAA 1770
GlnIleAspPheSerLeuArgGlyThrValPheGlyAlaLeuLysThrLeuAlaValLys 590

GCTATTGAAAAAACCTAGACTTGACCTATCAATGTGATTCATCGTTTCCAGATAATCTT 1830
AlaIleGluLysAsnLeuAspLeuThrTyrGlnCysAspSerSerPheProAspAsnLeu 610

ATTGGAGATAGTTTTAGATTACGACAAGTTATTCTTAACCTGGCTGGTAATGCTATTAAG 1890
IleGlyAspSerPheArgLeuArgGlnValIleLeuAsnLeuAlaGlyAsnAlaIleLys 630
N
TTTACTAAAGAGGGGAAAGTTAGTGTAGTGTGAAAAGTCTGATAAAATGGTGTTAGAT 1950
PheThrLysGluGlyLysValSerValSerValLysLysSerAspLysMetValLeuAsp 650

AGTAAGTTGTTGTTAGAGGTTTGTGTTAGCGACACGGGAATAGGTATAGAGAAAGACAAA 2010
SerLysLeuLeuLeuGluValCysValSerAspThrGlyIleGlyIleGluLysAspLys 670
G1
TTGGGATTGATTTTCGATACCTTCTGTCAAGCTGATGGTCTACTACAAGAAAGTTGGT 2070
LeuGlyLeuIlePheAspThrPheCysGlnAlaAspGlySerThrThrArgLysPheGly 690
← Slb2
GGTACAGGTTTAGGGTTGTCAATTTCCAAACAGTTGATACATTTAATGGGTGGAGAGATA 2130
GlyThrGlyLeuGlyLeuSerIleSerLysGlnLeuIleHisLeuMetGlyGlyGluIle 710
G2
TGGGTTACTTCGGAGTATGGATCCGGRTCAAACCTTTTATTTTACGGTGTGCGTGTCCGCA 2190
TrpValThrSerGluTyrGlySerGlySerAsnPheTyrPheThrValCysValSerPro 730

TCTAATATTAGATATACTCGACAAACCGAACAATTGTTACCATTTAGTTCCCATTATGTG 2250
SerAsnIleArgTyrThrArgGlnThrGluGlnLeuLeuProPheSerSerHisTyrVal 750

TTATTTGTATCGACTGAGCATACTCAAGAAGAACTTGATGTGTTGAGAGATGGAATTATA 2310
LeuPheValSerThrGluHisThrGlnGluGluLeuAspValLeuArtAspGlyIleIle 770

FIG. 1B

GAAGCTTGGATTGATACCTATAAATAGTGAGAAATATTGAAGATGCAACATTGACTGAGCCG 2370
 GluLeuGlyLeuIleProIleIleValArgAsnIleGluAspAlaThrLeuThrGluPro 790

GTGAAATATGATATAATTATGATTGATTTCGATAGAGATTGCCAAAAAGTTGAGGTTGTTA 2430
 ValLysTyrAspIleIleMetIleAspSerIleGluIleAlaLysLysLeuArgLeuLeu 810

TCGGAGGTTAAATATATTCCGTTGGTTTTGGTCCATCATTCTATTCCACAGTTGAATATG 2490
 SerGluValLysTyrIleProLeuValLeuValHisHisSerIleProGlnLeuAsnMet 830

AGAGTATGTATTGATTTGGGGATATCTTCCTATGCAAATACGCCATGTTTCGATCACGGAC 2550
 ArgValCysIleAspLeuGlyIleSerSerTyrAlaAsnThrProCysSerIleThrAsp 850

TTGGCCAGTGGCATTATAACCAGCGTTGGAGTCGAGATCTATATCACAGAACTCAGACGAG 2610
 LeuAlaSerAlaIleIleProAlaLeuGluSerArgSerIleSerGlnAsnSerAspGlu 870

TCGGTGAGGTACAAAATATTACTAGCAGAGGACAACCTCGTCAATCAGAACTTGCCAGTT 2670
 SerValArgTyrLysIleLeuLeuAlaGluAspAsnLeuValAsnGlnLysLeuAlaVal 890

AGGATATTAGAAAAGCAAGGCCATCTGGTGAAGTAGTTGAGAACGGACTCGAGGCCGTAC 2730
 ArgIleLeuGluLysGlnGlyHisIleValGluValValGluAsnGlyLeuGluAlaTyr 910

GAAGCGATTAAGAGGAATAAATATGATGTGGTGTGATGGATGTGCAAATGCCT 2784
 GluAlaIleLysArgAsnLysTyrAspValValLeuMetAspValGlnMetPro 928

← Slb3

D

FIG. 2A

ATGAACCCCACTAAAAACCTCGGTTATCACCAATGCAGCCCTCTGTTTTGAAATACTC 60
MetAsnProThrLysLysProArgLeuSerProMetGlnProSerValPheGluIleLeu 20

AACGACCCTGAGCTTTATAGTCAGCACTGTCATAGCCTTAGGGAAACACTTCTTGATCA 20
AsnAspProGluLeuTyrSerGlnHisCysHisSerLeuArgGluThrLeuLeuAspHis 40

TTCAACCATCAAGCTACACTTATCGACACTTATGAACATGAACTAGAAAAATCCAAAAAT 180
PheAsnHisGlnAlaThrLeuIleAspThrTyrGluHisGluLeuGluLysSerLysAsn 60

GCCAAACAAGCGTCCCAACAAGCACTTAGTGAAATAGGTACAGTTGTTATATCTGTTGCC 240
AlaAsnLysAlaSerGlnGlnAlaLeuSerGluIleGlyThrValValIleSerValAla 80

ATGGGAGACTTGTCGAAAAAGTTGAGATTCACACAGTAGAAAATGACCCTGAGATTTTA 300
MetGlyAspLeuSerLysLysValGluIleHisThrValGluAsnAspProGluIleLeu 100

AAAGTCAAATCACCATCAACACCATGATGGATCAATTACAGACATTTGCTAATGAGGTT 360
LysValLysIleThrIleAsnThrMetMetAspGlnLeuGlnThrPheAlaAsnGluVal 120

ACAAAAGTCGCCACCGAAGTCGCAAATGGTGAAC TAGGTGGACAAGCGAAAAATGATGGA 420
ThrLysValAlaThrGluValAlaAsnGlyGluLeuGlyGlyGlnAlaLysAsnAspGly 140

TCTGTTGGTATTTGGAGATCACTTACAGACAATGTTAATATTATGGCTCTTAATTTAACT 480
SerValGlyIleTrpArgSerLeuThrAspAsnValAsnIleMetAlaLeuAsnLeuThr 160

AACCAAGTGCAGAGAAATTGCTGATGTCACACGTGCTGTTGCCAAGGGGACTTGTCACGT 540
AsnGlnValArgGluIleAlaAspValThrArgAlaValAlaLysGlyAspLeuSerArg 180

AAAATTAATGTACACGCCCCAGGGTGAAATCCTTCAACTTCAACGTACAATAAACACCATG 600
LysIleAsnValHisAlaGlnGlyGluIleLeuGlnGluGlnArgThrIleAsnThrMet 200

GTGGATCAGTTACGAACGTTTGCATTCTGAAGTATCTAAAGTTGCTAGAGATGTTGGTGTG 660
ValAspGlnLeuArgThrPheAlaPheGluValSerLysValAlaArgAspValGlyVal 220

CTTGGTATATTAGGAGGACAAGCGTTGATTGAAAATGTTGAAGGTATTTGGGAAGAGTTG 720
LeuGlyIleLeuGlyGlyGlnAlaLeuIleGluAsnValGluGlyIleTrpGluGluLeu 240

ACTGATAATGTCAATGCCATGGCTCTTAATTTGACTACACAAGTGAGAAATATTGCCAAT 780
ThrAspAsnValAsnAlaMetAlaLeuAsnLeuThrThrGlnValArgAsnIleAlaAsn 260

FIG. 2B

GTCACCACTGCCGTTGCCAAGGGGGATTTGTCGAAAAAAGTCACTGCTGATTGTAAGGGA 840
ValThrThrAlaValAlaLysGlyAspLeuSerLysLysValThrAlaAspCysLycGly 280

GAAATYCTTGATTTGAAACTTACTATTAATCAAATGGTGGACCGATTACAGAATTTTGCT 900
GluIleLeuAspLeuLysLeuThrIleAsnGlnMetValAspArgLeuGlnAsnPheAla 300

CTTGCGGTGACGACATTGTCGAGAGAGGTTGGTACTTTGGGTATTTTGGGTGGACAAGCT 960
LeuAlaValThrThrLeuSerArgGluValGlyThrLeuGlyIleLeuGlyGlyGlnAla 320

AACGTACAGGATGTTGAAGGTGCTTGGAAACAGGTTACAGAAAATGTCAACCTAATGGCT 1020
AsnValGlnAspValGluGlyAlaTrpLysGlnValThrGluAsnValAsnLeuMetAla 340

ACTAATTTAACTAACCAAGTGAGATCTATTGCTACAGTTACTACTGCAGTTGCGCATGGT 1080
ThrAsnLeuThrAsnGlnValArgSerIleAlaThrValThrThrAlaValAlaHisGly 360

GATTTGTCGCAAAAGATTGATGGTCATCCCAAAGGAGAGATTTTACAATTGAAAAATACA 1140
AspLeuSerGlnLysIleAspGlyHisProLysGlyGluIleLeuGlnLeuLysAsnThr 380

ATCAACAAGATGGTGGACTCTTTGCAGTTGTTGCATCAGAAGTGTGCGAAAGTGGCACAA 1200
IleAsnLysMetValAspSerLeuGlnLeuPheAlaSerGluValSerLysValAlaGln 400

GATGTTGGTATTAATGGAAAATTAGGTATTCAAGCACAAGTTAGTGATGTTGATGGATTA 1260
AspValGlyIleAsnGlyLysLeuGlyIleGlnAlaGlnValSerAspValAspGlyLeu 420

TGGAAGGAGATTACGTCTAATGTAAATACCATGGCTTCAAATTTAACTTCGCAAGTGAGA 1320
TrpLysGluIleThrSerAsnValAsnThrMetAlaSerAsnLeuThrSerGlnValArg 440

GCTTTTGCACAGATTACTGCTGCTGCTACTGATGGGGATTTCACTAGATTTATTACTGTT 1380
AlaPheAlaGlnIleThrAlaAlaAlaThrAspGlyAspPheThrArgPheIleThrVal 460

GAAGCACTGGGAGAGATGGATGCGTTGAAAACAAAGATTAATCAAATGGTGTTTAACTTA 1440
GluAlaLeuGlyGluMetAspAlaLeuLysThrLysIleAsnGlnMetValPheAsnLeu 480

AGGGAATCGCTTCAAAGGAATACTGCGGCTAGAGAAGCTGCTGAGTTGGCCAATAGTGCG 1500
ArgGluSerLeuGlnArgAsnThrAlaAlaArgGluAlaAlaGluLeuAlaAsnSerAla 500

AAATCCGAGTTTTTTAGCAAACATGTCGCATGAGATTAGAACACCATTGAATGGGATTATT 1560
LysSerGluPheLeuAlaAsnMetSerHisGluIleArgThrProLeuAsnGlyIleIle 520

FIG. 2C

GGWATGACYCAGTTGTCRCTTGATACAGAGTTGACRCAGTACCAACGAGAGATGTTGTGCG 1620
 GlyMetThrGlnLeuSerLeuAspThrGluLeuThrGlnTyrGlnArgGluMetLeuSer 540

ATTGTGCATAACTTGGCAAATTCCTTGTTGACCATTATAGACGATATATTGGATATTTCT 1680
 IleValHisAsnLeuAlaAsnSerLeuLeuThrIleIleAspAspIleLeuAspIleSer 560

AAGATTGAGGCGAATAGAATGACGGTGAACAGATTGATTTTTCATTAAGAGGGACAGTG 1740
 LysIleGluAlaAsnArgMetThrValGluGlnIleAspPheSerLeuArgGlyThrVal 580

TTTGGTGCATTGAAAACGTTAGCCGTCAAAGCTATTGAAAAAACCTAGACTTGACCTAT 1800
 PheGlyAlaLeuLysThrLeuAlaValLysAlaIleGluLysAsnLeuAspLeuThrTyr 600

CAATGTGATTCATCGTTTCCAGATAATCTTATTGGAGATAGTTTTAGATTACGACAAGTT 1860
 GlnCysAspSerSerPheProAspAsnLeuIleGlyAspSerPheArgLeuArgGlnVal 620

ATTCTTAACTTGGCTGGTAATGCTATTAAGTTTACTAAAGAGGGGAAAGTTAGTGTTAGT 1920
 IleLeuAsnLeuAlaGlyAsnAlaIleLysPheThrLysGluGlyLysValSerValSer 640

N

GTGAAAAAGTCTGATAAAATGGTGTTAGATAGTAAGTTGTTGTTAGAGGTTTGTGTTAGC 1980
 ValLysLysSerAspLysMetValLeuAspSerLysLeuLeuLeuGluValCysValSer 660

GACACGGGAATAGGTATAGAGAAAGACAAATTGGGATTGATTTTCGATACCTTCTGTCAA 2040
AspThrGlyIleGlyIleGluLysAspLysLeuGlyLeuIlePheAspThrPheCysGln 680

G1

GCTGATGGTCTACTACAAGAAAGTTTGGTGGTACAGGTTTAGGGTTGTCAATTTCCAAA 2100
 AlaAspGlySerThrThrArgLysPheGlyGlyThrGlyLeuGlyLeuSerIleSerLys 700

G2

CAGTTGATACATTTAATGGGTGGAGAGATATGGGTTACTTCGGAGTATGGATCCGGRTCA 2160
 GlnLeuIleHisLeuMetGlyGlyGluIleTrpValThrSerGluTyrGlySerGlySer 720

AACTTTTATTTACGGTGTGCGTGTGCCATCTAATATTAGATATACTCGACAAACCGAA 2220
 AsnPheTyrPheThrValCysValSerproSerAsnIleArgTyrThrArgGlnThrGlu 740

CAATTGTTACCATTTAGTTCCCATATGTGTTATTTGTATCGACTGAGCATACTCAAGAA 2280
 GlnLeuLeuProPheSerSerHisTyrValLeuPheValSerThrGluHisThrGlnGlu 760

GAACTTGATGTGTTGAGAGATGGAATTATAGAACTTGGATTGATACCTATAATAGTGAGA 2340
 GluLeuAspValLeuArgAspGlyIleIleGluLeuGlyLeuIleProIleIleValArg 780

FIG. 2D

AATATTGAAGATGCAACATTGACTGAGCCGGTGAAATATGATATAATTATGATTGATTTCG 2400
 AsnIleGluAspAlaThrLeuThrGluProValLysTyrAspIleIleMetIleAspSer 800

ATAGAGATTGCCAAAAGTTGAGGTTGTTATCGGAGGTTAAATATATTCGGTTGGTTTTG 2460
 IleGluIleAlaLysLysLeuArgLeuLeuSerGluValLysTyrIleProLeuValLeu 820

GTCCATCATTCTATTCCACAGTTGAATATGAGAGTATGTATTGATTTGGGGATATCTTCC 2520
 ValHisHisSerIleProGlnLeuAsnMetArgValCysIleAspLeuGlyIleSerSer 840

TATGCAAATACGCCATGTTTCGATCACGGACTTGGCCAGTGGCATTATACCAGCGTTGGAG 2580
 TyrAlaAsnThrProCysSerIleThrAspLeuAlaSerAlaIleIleProAlaLeuGlu 860

TCGAGATCTATATCACAGAACTCAGACGAGTCGGTGAGGTACAAAATATTACTAGCAGAG 2640
 SerArgSerIleSerGlnAsnSerAspGluSerValArgTyrLysIleLeuLeuAlaGlu 880

GACAACCTCGTCAATCAGAACTTGCAGTTAGGATATTAGAAAAGCAAGGGCATCTGGTG 2700
 AspAsnLeuValAsnGlnLysLeuAlaValArgIleLeuGluLysGlnGlyHisLeuVal 900

GAAGTAGTTGAGAACGGACTCGAGGCGTACGAAGCGATTAAGAGGAATAAATATGATGTG 2760
 GluValValGluAsnGlyLeuGluAlaTyrGluAlaIleLysArgAsnLysTyrAspVal 920

GTGTTGATGGATGTGCAAATGCCTGTAATGGGTGGGTTTGAAGCTACGGAGAAGATTCCA 2820
 ValLeuMetAspValGlnMetProValMetGlyGlyPheGluAlaThrGluLysIleArg 940

D

CAATGGGAGAAAAAGTCTAACCCAATTGACTCGTTGACCTTTAGGACTCCAATTATTGCC 2880
 GlnTrpGluLysLysSerAsnProIleAspSerLeuThrPheArgThrProIleIleAla 960

CTCACTGCACACGCCATGTTAGGTGATAGAGAAAAGTCATTGGCCAAGGGGATGGACGAT 2940
 LeuThrAlaHisAlaMetLeuGlyAspArgGluLysSerLeuAlaLysGlyMetAspAsp 980

TATGTGAGTAAGCCATTGAAGCCGAAATTGTTAATGCAGACGATAAAGAAGTGTATTCAT 3000
 TyrValSerLysProLeuLysProLysLeuLeuMetGlnThrIleAsnLysCysIleHis 1000

H2

AATATTAACCAGTTGAAAGAATTGTCGAGAAATAGTAGGGGTAGCGATTTTGCAAAGAAG 3060
 AsnIleAsnGlnLeuLysGluLeuSerArgAsnSerArgGlySerAspPheAlaLysLys 1020

ATGACCCGAAACACACCCGGCCGCACGCCCGTCAGGGGAGTGATGAGGGGAGTGTAAG 3120
 MetThrArgAsnThrProGlySerThrThrArgGlnGlySerAspGluGlySerValLys 1040

FIG. 2E

GACATGATTGGGGACACTCCCCGTCAAGGGAGTGTGGAGGGAGGGGGTACAAGTAGTAGA 3180
AspMetIleGlyAspThrProArgGlnGlySerValGluGlyGlyGlyThrSerSerArg 1060

CCAGTACAGAGAAGGTCTGCCAGGGAGGGGTCGATCACTACAATTAGTGAACAAATCGAC 3240
ProValGlnArgArgSerAlaArgGluGlySerIleThrThrIleSerGluGlnIleAsp 1080

CGTTAG 3246
Arg*** 1082




FIG. 3




FIG. 4

**CANDIDA ALBICANS TWO-COMPONENT
HYBRID KINASE GENE, CANIK1, AND USE
THEREOF**

This application claims benefit of provisional application 60/048,914 filed Jun. 6, 1997.

This application describes microorganisms that have been deposited, in accordance with the Budapest Treaty, under ATCC Patent Deposit Designation: PTA-4456, with the following Deposit Identification Reference: Bacteriophage lambda EMBL3: Ca lambda 15.1.

BACKGROUND OF THE INVENTION

Candida is an opportunistic yeast that lives in the mouth, throat, intestines, and genitourinary tract of most humans. In a healthy human body, the population of *Candida* is kept in check by the immune system and by a competitive balance with other microorganisms. But when the body's immune system is compromised, as in AIDS patients and in patients undergoing immunosuppressive therapy, *Candida* will grow uncontrolled, leading to systemic infection called "*Candida mycosis*." If left untreated, such systemic infections frequently lead to the death of the patients.

Candida albicans is a species of particular interest to scientists and doctors because 90% of all cases of *Candida mycosis* are caused by this species.

At present, the therapy principally available for invasive infections is based on relatively few antimycotics, such as amphotericin B and flucytosine, or the azole derivatives fluconazole and itraconazole. These antimycotics cause serious side effects, such as renal insufficiency, hypocalcaemia and anaemia, as well as unpleasant constitutional symptoms such as fever, shivering and low blood pressure. Amphotericin B is toxic to the kidneys, for example, and yet the pharmaceutical is therapeutic only if administered at dose levels near to being toxic. A discussion of the pharmaceuticals used for treatment and their corresponding side effects can be found, for example, in Boyd, et al., *BASIC MEDICAL MICROBIOLOGY* (2d ed.), Little, Brown and Company, (1981).

Given the deficiencies of conventional therapies against *Candida*, a need exists for developing pharmaceuticals that are effective in this regard and also safe to use. One step in the development of such pharmaceuticals requires a method for screening compounds in order to identify pharmaceutical candidates.

SUMMARY OF THE INVENTION

It therefore is an object of the present invention to provide an isolated polynucleotide sequence coding for a protein that is linked to phenotypic switching in *Candida albicans*.

It is a further object of the invention to provide a method for screening compounds to identify pharmaceutical candidates for effectively inhibiting the pathogenicity of *C. albicans*.

In accomplishing these and other objects, there has been provided, according to one aspect of the present invention, an isolated polynucleotide that codes for such a protein and that hybridizes, under stringent conditions, to the polynucleotide sequence of SEQ ID NO:1, shown below in FIG. 1. In a preferred embodiment, the polynucleotide has the sequence of SEQ ID NO:3 (FIG. 2). In another preferred embodiment, the protein displays a kinase activity.

In accordance with another aspect of the present invention, a method is provided for screening compounds to identify pharmaceutical candidates. The inventive method comprises the steps of (A) providing a plurality of cells from yeast species that exhibit phenotypic switching, at least

some of which contain (i) a polynucleotide coding for a CaNIK1 protein and (ii) a promoter that is operably linked to the polynucleotide, such that the plurality of cells produces the protein; then (B) bringing the plurality into contact with a test substance; and (C) assessing what effect, if any, the test substance has on the expression of the DNA segment. Assessment step (C) can comprise, for example, of monitoring the level either of the protein or the corresponding mRNA transcript produced by the plurality of cells. In another embodiment, step (C) comprises monitoring the level of kinase activity, within the plurality, that typifies the protein.

In yet another embodiment of the present invention, a promoter is operably linked to a reporter gene. In this context, step (C) comprises monitoring the level of transcription of the reporter gene, after contact between the plurality of cells and the test substance.

Other objects, features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples, only indicate preferred embodiments of the invention.

BRIEF DESCRIPTION OF THE DRAWINGS

FIGS. 1A–B show the nucleotide sequence (SEQ ID NO:11) (top row) of the PCR product encoding the region spanning the H1 and D domains and the deduced amino acid sequence of the CaNIK1 protein (SEQ ID NO:2) (bottom row). The amino acid residues of functional domains are underlined. The three degenerate primers used to isolate the PCR products are shown as Slb1, Slb2, and Slb3.

FIGS. 2A–E show the nucleotide sequence (SEQ ID NO:3) (top row) of the gene CaNik1 and the deduced primary amino acid sequence of the CaNIK1 protein (SEQ ID NO:11) (bottom row). The beginning of each unique repeat is represented within the rectangle. The potential amino acid residues of different functional domains are underlined.

FIG. 3 is a schematic representation of the anatomy of two alleles in two strains of *C. albicans* according to the present invention. All the functional domains are shown as white bold letters inside each rectangle. A few of the unique restriction enzyme sites are shown at the top of the rectangle. The start of the protein coding region is shown as ATG. WO-1 and CAI8 are the two strains analyzed in this invention. H1 and H2 are two identical alleles of the strain WO-1. H1-L and H2-S represent large and small alleles respectively in strain CAI8. The five hatched rectangular units in each allele represent repeat units described in this invention. The gray rectangular area encompassing XhoI-PstI in H2-S represents the region containing a deletion of approximately one repeat unit length.

FIG. 4 illustrates the deletion strategy used to generate a homozygous deletion mutant, HH80, in strain CAI8. The region spanning AflII-XhoI was deleted and substituted by a hisG-Urablaster cassette in the plasmid pUNIK12.1 to create pCNH35 (FIG. 4c). Plasmid pUNIK12.1 (FIG. 4b) was derived by subcloning a PCR fragment using a pair of primers Slb8 and Slb7R and subcloning into pGEM-T easy plasmid vector. λ SA15.1 represent the lambda clone identified in a screen that contain the genomic fragment encompassing the entire CaNik1 gene and the flanking DNA sequence.

FIG. 5 shows the deletion strategy used to generate the homozygous deletion mutant in Red 3/6, an *ade2*⁻ derivative of strain WO-1. The deletion cassette pABX12 (FIG. 5b) was generated by deletion of all the functional domains except H2 and substitution with the ADE2 gene as an auxotrophic marker in pUNIK12.1 (FIG. 5c). FIG. 4 provides a description of λ SA15.1.

Table 1 summarizes the effects of the CaNik1 deletion in HH80 on growth in a variety of solution and conditions, high frequency phenotypic switching, and dimorphism.

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

Candida albicans is capable of differentiating in a reversible fashion between a bud and a hyphal growth form. Each strain of *C. albicans* can also undergo high frequency phenotypic switching between a limited number of general phenotypes that differ in a variety of traits including putative virulence factors. The frequencies of both of these developmental programs are influenced by environmental conditions. For example, pH and temperature influence the transition between bud and hypha while temperature, UV, white blood cell metabolites and colony aging affect the frequency of high frequency phenotypic switching. The morphological changes made by *C. albicans* in response to environmental cues indicates that the organism uses a sensory mechanism to register and assess environmental alterations.

Autophosphorylating histidine kinases, also known as "two-component response regulators," have been found, in lower eukaryotes such as fungi and slime molds, to play a pivotal role in relaying various environmental signals into the cell for inducing appropriate responses and in providing these organisms with the capacity to respond rapidly to an environmental perturbation. Two-component signal transducers all contain a sensory kinase, which autophosphorylates a histidine residue in response to an environmental cue, and a response regulator, which then is phosphorylated and, through a resultant conformational change, effects a signal that is transduced either directly to a molecular complex, as in the case of the bacterial CheY and the flagellar motor, or down a signal transduction pathway, as in the case of SLN1. These proteins have been shown to be involved in regulating morphogenesis and development in various prokaryotes and eukaryotes.

That two-component response regulators have been identified in other yeast species suggests that the two-component response regulators may also play a role in the developmental programs of *C. albicans*. The present invention relates to such a two-component response regulator, the hybrid kinase CaNIK1 from *Candida albicans*. A link between the gene encoding CaNik1 and the processes of phenotypic switching that includes the differential expression of pathogenic genes is evidenced by work with a CaNik1-deletion strain of *C. albicans*. See examples 3 and 5. Thus, CaNik1 is known to be involved in phenotypic switching.

Phenotypic switching is thought to be linked to the virulent characteristics of yeast. *Candida albicans* switches phenotypes with regard to its environment in order to maximize pathogenesis according to the demands of the particular environment. For example, in the WO-1 strain of *Candida albicans*, studies have shown that the yeast is more virulent in its opaque phenotype when located on the skin. When WO-1 is in the white phenotype, however, it is more pathogenic in systemic infections. A description of the relationship between the phenotypic switching and the pathogenic characteristics of *Candida albicans* can be found in Soll, "Switching and Gene Regulation in *Candida albicans*," in SOCIETY FOR GENERAL MICROBIOLOGY SYMPOSIUM 50 (1992). This relationship between phenotypic switching and pathogenicity can be exploited effectively, in a bioassay, for the purpose of discovering pharmaceutical candidates against *Candida albicans*.

1. Definitions

In this description, "isolated DNA" is a fragment of DNA that is not integrated in the genomic DNA of an organism. For example, the CaNik1 gene is a DNA fragment that has been isolated from the genomic DNA of *C. albicans*.

As used herein, "protein" refers to a polymer of amino acid residues. The terms apply to amino acid polymers in which one or more amino acid residue is an artificial chemical analogue of a corresponding naturally occurring amino acid, as well as to naturally occurring amino acid polymers. The essential nature of such analogues of naturally occurring amino acids is that, when incorporated into a protein, that protein is specifically reactive to antibodies elicited to the same protein but consisting entirely of naturally occurring amino acids. The terms "polypeptide", "peptide" and "protein" are also inclusive of modifications including, but not limited to, glycosylation, lipid attachment, sulfation, gamma-carboxylation of glutamic acid residues, hydroxylation and ADP-ribosylation. Exemplary modifications are described in most basic texts, such as PROTEINS—STRUCTURE AND MOLECULAR PROPERTIES (2d ed.), T. E. Creighton, W. H. Freeman and Company, New York (1993).

As used herein, "selectively hybridizes" includes reference to hybridization, under stringent hybridization conditions, of a nucleic acid sequence to a specified nucleic acid target sequence to a detectably greater degree (e.g., at least 2-fold over background) than its hybridization to non-target nucleic acid sequences and to the substantial exclusion of non-target nucleic acids. Selectively hybridizing sequences typically have about at least 80% sequence identity, preferably 90% sequence identity, and most preferably 100% sequence identity (i.e., complementary) with each other.

The terms "stringent conditions" or stringent hybridization conditions includes reference to conditions under which a probe will hybridize to its target sequence, to a detectably greater degree than other sequences (e.g., at least 2-fold over background). Stringent conditions are sequence-dependent and will be different in different circumstances. By controlling the stringency of the hybridization and/or washing conditions, target sequences can be identified which are 100% complementary to the probe (homologous probing). Alternatively, stringency conditions can be adjusted to allow some mismatching in sequences so that lower degrees of similarity are detected (heterologous probing). Generally, a probe is less than about 1000 nucleotides in length, preferably less than 500 nucleotides in length.

Specificity is typically the function of post-hybridization washes, the critical factors being the ionic strength and temperature of the final wash solution. For DNA-DNA hybrids, the T_m can be approximated from the equation of Meinkoth & Wahl, Anal. Biochem. 138: 267-84 (1984): $T_m = 81.5^\circ \text{C} + 16.6 (\log M) + 0.41 (\% \text{GC}) - 0.61 (\% \text{form}) - 500/L$; where M is the molarity of monovalent cations, % GC is the percentage of guanosine and cytosine nucleotides in the DNA, % form is the percentage of formamide in the hybridization solution, and L is the length of the hybrid in base pairs. The T_m is the temperature (under defined ionic strength and pH) at which 50% of a complementary target sequence hybridizes to a perfectly matched probe. T_m is reduced by about 1°C . for each 1% of mismatching; thus, T_m , hybridization and/or wash conditions can be adjusted to hybridize to sequences of the desired identity. For example, if sequences with $\geq 90\%$ identity are sought, the T_m can be decreased 10°C . Generally, stringent conditions are selected to be about 5°C . lower than the thermal melting point (T_m) for the specific sequence and its complement at a defined ionic strength and pH. But severely stringent conditions can utilize a hybridization and/or wash at 1, 2, 3, or 4°C . lower than the thermal melting point (T_m); moderately stringent conditions can utilize a hybridization and/or wash at 6, 7, 8, 9, or 10°C . lower than the thermal melting point (T_m); low stringency conditions can utilize a hybridization and/or wash at 11, 12, 13, 14, 15, or 20°C . lower than the thermal melting point (T_m). Using the equation, hybridization and

5

wash compositions, and desired T_m , those of ordinary skill will understand that variations in the stringency of hybridization and/or wash solutions are inherently described. If the desired degree of mismatching results in a T_m of less than 45° C. (aqueous solution) or 32° C. (formamide solution) it is preferred to increase the SSC concentration so that a higher temperature can be used. An extensive guide to the hybridization of nucleic acids is found in Tijssen, LABORATORY TECHNIQUES IN BIOCHEMISTRY AND MOLECULAR BIOLOGY—HYBRIDIZATION WITH NUCLEIC ACID PROBES, Part I, Chapter 2 “Overview of principles of hybridization and the strategy of nucleic acid probe assays,” Elsevier, New York (1993); and in Chapter 2 of CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, Greene Publishing and Wiley-Interscience, New York (1995) (hereafter “Ausubel et al.”).

Two nucleic acid molecules are considered to have a “substantial sequence similarity” if their nucleotide sequences share a similarity of at least 50%. Sequence similarity determinations can be performed, for example, using the FASTA program (Genetics Computer Group, Madison, Wis.). Alternatively, sequence similarity determinations can be performed using BLASTP (Basic Local Alignment Search Tool) of the Experimental GENINFO (R) BLAST Network Service. See Altschul et al., “Sequence Similarity Searches, Multiple Sequence Alignments, and Molecular Tree Building,” in METHODS IN PLANT MOLECULAR BIOLOGY AND BIOTECHNOLOGY, Glick et al. (eds.), pages 251–267 (CRC Press, 1993).

As used herein “promoter” includes reference to a region of DNA upstream from the start of transcription and involved in recognition and binding of RNA polymerase and other proteins to initiate transcription. Tissue-specific, tissue-preferred, cell type-specific, and inducible promoters constitute the class of “non-constitutive” promoters. A “constitutive” promoter is one that is active under most environmental conditions.

As used herein “operably linked” includes reference to a functional linkage between a promoter and a second sequence, wherein the promoter sequence initiates and mediates transcription of the DNA sequence corresponding to the second sequence. Generally, operably linked means that the nucleic acid sequences being linked are contiguous and, where necessary to join two protein coding regions, contiguous and in the same reading frame.

As used herein, “expression” refers to the biosynthesis of a gene product. For example, in the case of a structural gene, expression involves transcription of the structural gene into mRNA and the translation of mRNA into one or more polypeptides.

As used herein, “expression vector” is a polynucleotide molecule comprising a gene that is expressed in a host cell. Typically, gene expression is placed under the control of certain regulatory elements, including constitutive or inducible promoters, tissue-specific regulatory elements, and enhancers. Such a gene is said to be “operably linked” to the regulatory elements.

2. Isolating a *Candida albicans* Polynucleotide Segment Encoding CaNik1 Protein

An endogenous polynucleotide sequence from *Candida albicans* which encodes for the CaNIK1 protein was isolated using a polynucleotide probe derived from PCR amplification. See Example 1. Hybridization of the probe against a genomic library resulted in the determination of the full length polynucleotide sequence encoding the CaNIK1 protein. See Example 2. The full polynucleotide sequence encapsulating the CaNik1 gene is provided in FIG. 2.

3. Nucleic Acids

The present invention provides, inter alia, isolated nucleic acids of RNA, DNA, and analogs and/or chimeras thereof, comprising a polynucleotide encoding a CaNIK1 protein or

6

a polynucleotide probe which hybridizes to a polynucleotide encoding CaNIK1 protein. In this regard, the invention provides the nucleotide sequences of FIGS. 1 and 2. In addition, the present invention also provides other sequences as described below.

a. Polynucleotides Encoding A CaNIK1 Polypeptide or Conservatively Modified or Polymorphic Variants Thereof

As indicated above, the present invention provides isolated heterologous nucleic acids comprising a polynucleotide, wherein the polynucleotide encodes a CaNIK1 protein, disclosed herein in FIG. 2, or conservatively modified or polymorphic variants thereof. Those of skill in the art will recognize that the degeneracy of the genetic code allows for a plurality of polynucleotides to encode for the identical amino acid sequence. Such “silent variations” can be used, for example, to selectively hybridize and detect allelic variants of polynucleotides of the present invention. Accordingly, the present invention includes polynucleotides that are silent variations of the polynucleotides of FIG. 2. The present invention further provides isolated nucleic acids comprising polynucleotides encoding conservatively modified variants of CaNIK1 encoded by the sequences in FIG. 2. Conservatively modified variants can be used to generate or select antibodies immunoreactive to the non-variant polypeptide. Additionally, the present invention further provides isolated nucleic acids comprising polynucleotides encoding one or more polymorphic (allelic) variants of polypeptides/polynucleotides.

b. Polynucleotides That Selectively Hybridize

The present invention also provides isolated nucleic acids comprising polynucleotides, wherein the polynucleotides selectively hybridize, under selective hybridization conditions, to a polynucleotide as discussed above. In this regard, the present invention encompasses polynucleotides that selectively hybridize, under selective conditions, to a polynucleotide as discussed above, excluding the polynucleotide of FIG. 2. Thus, the polynucleotides of this embodiment can be used for isolating, detecting, and/or quantifying nucleic acids comprising the polynucleotides described above. For example, polynucleotides of the present invention can be used to identify, isolate, or amplify partial or full-length clones in a deposited library. In some embodiments, the polynucleotides are genomic or cDNA sequences isolated, or otherwise complementary to, a cDNA from a nucleic acid library. Preferably, the cDNA library comprises at least 80% full-length sequences, preferably at least 85% or 90% full-length sequences, and more preferably at least 95% full-length sequences. The cDNA libraries can be normalized to increase the representation of rare sequences. Low stringency hybridization conditions are typically, but not exclusively, employed with sequences having a reduced sequence identity relative to complementary sequences. Moderate and high stringency conditions can optionally be employed for sequences of greater identity. Low stringency conditions allow selective hybridization of sequences having about 70% sequence identity and can be employed to identify orthologous or paralogous sequences.

c. Polynucleotides Having at Least 60% Sequence Identity

The present invention further provides isolated nucleic acids comprising polynucleotides, wherein the polynucleotides have a specified identity at the nucleotide level to a polynucleotide as disclosed above. In this regard, the present invention encompasses polynucleotides that have a specified identity to the polynucleotides discussed above, but are not the same as the sequence of FIG. 2. The percentage of identity to a reference sequence is at least 60% and, rounded upwards to the nearest integer, can be expressed as an integer selected from the group of integers consisting of from 60 to 99. Thus, for example, the percentage of identity to a reference sequence can be at least 70%, 75%, 80%, 85%, 90%, or 95%.

4. Vectors

According to the present invention, the polynucleotide sequence encoding the CaNIK1 protein may be inserted into any suitable yeast vector with any method known to a person who has skill in the art. The vector will typically be comprised of a polynucleotide encoding the CaNIK1 protein operably linked to any suitable promoter which will direct the transcription of the polynucleotide in the intended host cell. Examples of suitable promoters include EF1 α 2 which is a constitutive promoter and is characterized in Sundstrom et al., *General Bacteriology*, 172: 2036–2045 (1990), and PCK1 which is an inducible promoter and is characterized in Leuker et al., *Gene* 192: 235–240 (1997). According to the present invention, the promoter is operably linked to the polynucleotide encoding for the CaNIK1 protein and inserted into a yeast transformation vector.

Yeast vectors are grouped into five general classes according to their mode of replication in the yeast: YIp, YRp, YCp, YE_p, YL_p. Comprehensive laboratory techniques regarding insertion of polynucleotides into yeast vectors can be found in Chapter 13 of Ausubel et al.

5. Bioassay

Another aspect of the invention is a bioassay useful for screening pharmaceutical candidates which can inhibit pathogenicity in *Candida albicans*. The bioassay is based on assessing a candidate's ability to inhibit expression or functionality of the CaNik1 gene or its gene product, which as explained above, is linked to the virulent characteristics of the yeast. A bioassay according to the present invention comprises the following steps: transformation of cells from yeast species that exhibit phenotypic switching with a polynucleotide encoding CaNIK1 protein, and a promoter linked to the polynucleotide segment which can drive protein expression; effecting contact between the yeast cells and a pharmaceutical candidate; and analyzing the effect of the pharmaceutical candidate on inhibition of the expression of the CaNik1 gene. In one embodiment, *C. albicans* cells harboring a CaNik1 deletion are transformed with a suitable construct containing a CaNIK1-encoding polynucleotide, and an operably linked promoter.

A. Transformation of Yeast Cells

The present invention contemplates the use of yeast cells with a phenotypic switching pathway similar to that of *Candida albicans*. Srikantha et al., *J. Bacteriol.* 179: 3837–3844 (1997). Transformation of the cells can be accomplished through any means known to a person with skill in the art. One example of a yeast transformation procedure is the lithium acetate procedure whereby yeast cells are briefly incubated in buffered lithium acetate and transforming DNA is introduced with carrier DNA. Addition of polyethylene glycol (PEG) and a heat shock trigger DNA uptake. An alternate method of transforming yeast cells is the electroporation procedure whereby concentrated cells are transformed using an exponential electric pulse. Comprehensive laboratory techniques regarding yeast transformation procedures can be found in Chapter 13 of Ausubel et al.

B. Contact of a Test Substance with Transformed Cells

According to the present invention, a test substance should make contact with at least some of a plurality of cells transformed with a polypeptide encoding CaNik1. Contact includes any exposure of the test substance to any surface of a transformed cell. A preferred method of contact would be incubation of the cells with the test substance.

The test substance includes any compound which may have characteristics inhibitory to the growth or the pathogenicity of *Candida albicans*. An example of a test substance is a pharmaceutical compound with antimycotic properties.

6. Assessing of the Effect of the Test Substance on CaNik1 Gene Expression

According to the present invention, the effect of the pharmaceutical compound on CaNik1 expression is analyzed after contact between the pharmaceutical compound and the plurality of transformed cells. CaNik1 expression can be measured through any means known by a person with skill in the art. Examples of methods which monitor the level of gene expression are: measuring levels of CaNIK1 protein and mRNA produced by the cells; or measuring the kinase activity within the cell; or monitoring the level of transcription of a reporter gene operably linked to a promoter.

An example of monitoring CaNik1 expression is the measurement of levels of CaNIK1 protein produced by the plurality of cells. This can be measured by performing two-dimensional gel electrophoresis using the techniques of isoelectric-focusing and SDS-polyacrylamide gel electrophoresis followed by autoradiography of the gel. Comprehensive laboratory techniques regarding two-dimensional gel electrophoresis and autoradiography can be found in Chapter 10 and Appendix 3 of Ausubel et al.

Another example of monitoring CaNik1 expression is to measure the level of mRNA encoded within the cell and produced by the plurality. mRNA levels within the cell can be measured with the following three techniques: Northern Blot, primer extension and ribonuclease protection. The Northern Blot procedure consists of fractionating mRNA with gel electrophoresis, transferring the mRNA fragments from the gel onto a filter and hybridizing the target mRNA molecules used a labeled DNA or RNA probe. The primer extension procedure includes hybridizing an oligonucleotide primer to the 5' end of the target mRNA and extending the primer using reverse transcriptase and unlabeled deoxynucleotides to form a single-stranded DNA complementary to the template RNA. The resultant DNA is analyzed on the sequencing gel. The yield of the primer extension product quantifies the amount of mRNA produced by the cell. The ribonuclease protection assay measures mRNA levels by hybridizing sequence specific RNA probes to sample RNAs. The probe anneals to homologous sequences in the sample RNA. The presence of target RNA is analyzed and quantified by gel electrophoresis. Comprehensive laboratory techniques regarding Northern Blot, primer extension and ribonuclease protection assays can be found in Chapter 4 of Ausubel et al.

A third example of monitoring CaNik1 expression is to monitor the level of kinase activity within the plurality of cells. Kinase activity within the cells can be monitored by labeling ATP with ³²P in vitro. The labeled ATP acts as the donor substrate, and the CaNIK1 protein acts as the acceptor substrate. Phosphotransfer is detected as the accumulation of ³²P-labeled protein within the cell. The accumulation of protein is measured with polyacrylamide gel electrophoresis and autoradiography. Target kinase activity can be distinguished from background kinase activity with autophosphorylation of the CaNIK1 protein on polyacrylamide gel. Comprehensive laboratory techniques regarding phosphorylation and measurement of kinase activity can be found in Chapter 18 of Ausubel et al.

In a further example, a reporter gene is operably linked to a promoter and the level of transcription of the reporter gene is monitored after contact between the plurality and the test substance. In accordance with the present invention, the promoter region of the CaNik1 gene is operably linked to the luciferase gene. Gene activity is thus linked to luciferase activity, which can then be measured quantitatively, with a luminometer, as a bioluminescent reaction.

The present invention is described further below by reference to the following examples, which are illustrative only.

EXAMPLE 1

PCR Amplification to Determine a CaNik1 Probe

The following, deoxyinosine-containing, degenerate primers were designed that encompassed the highly conserved regions of the two component response regulators LemA (Hrabak & Willis, *J Bacteriol* 174: 3011-3020 (1992)), BarA (Nagasawa et al., *Escherichia coli. Mol Microbiol*, 6: 799-807 (1992)) and SLN1 (Ota & Varshavsky, *Science* 263: 566-569 (1993)), respectively: 1) Slb1: (SEQ ID NO:5) 5-GAATTGAGAACGCCTITIAATGG-3, which corresponds to the histidine-autokinase domain; 2) Slb2: (SEQ ID NO:6) 5-AGTCCTAAGCCA GTACCACC-3, which corresponds to the ATP-binding domain; and 3) Slb3: (SEQ ID NO:7) 5-TTTAGGCATCTGGACITCCAT, which corresponds to the response regulator domain. Slb1 served as a 5'-end primer for PCR amplifications. The Slb1/Slb2 and Slb1/Slb3 pairs were used to amplify PCR products using the Hot-start wax gem (Perkin, Elmer) protocol. The Hot-start wax gem protocol which generates PCR products used the following reaction mixture: 10 mM Tris-HCl, pH 8.0, 50 mM KCl, 1.2 mM MgCl₂, 100 μM dNTP, 50 μM of each primer and 2.5 units of Taq polymerase, in a final volume of 100 μL. Conditions for PCR cycling included denaturation at 94° C. for 1 min, annealing at 40° C. for 1.5 min and extension at 72° C. for 2.5 min. For all amplifications, *S. cerevisiae* genomic DNA was used as a control for the amplification of the two component hybrid kinase gene SLN1, to monitor the quality of the PCR products. PCR products were gel purified and cloned into either PCR-Trap (Hunter Gen) or pGEM T-Easy (Promega Corp.). Three positive clones were chosen for each of the PCR products of the two sets of primer pairs. pCN.5/3, pCN.5/11 and pCN.5/21 were chosen from the products of Slb1/Slb2; and pCN1.3/5, pCN1.3/13 and pCN 1.3/16 were chosen from the products of Slb1/Slb3.

EXAMPLE 2

Isolation of CaNik1 Gene

To isolate a full-length gene, approximately 8×10⁴ plaques of a *C. albicans* genomic library were screened using a 1.2 kb DNA fragment isolated from pCN1.3/13, which spanned the histidine-autokinase (H1) and aspartyl receiver domain (D1). Lambda DNA from 20 positive clones was extracted and Southern blots probed with pCN1.3/13. Using combinations of primer pairs for the arms of the lambda DNA and either the degenerate primers for the histidine-autokinase domain (Slb1) or the response regulator domain (Slb3), lambda clones containing inserts larger than 4 kb were identified. The screen was performed with a high fidelity long PCR protocol (Boehringer Mannheim, Inc., Indianapolis, Ind.). Three lambda clones contained DNA fragments larger than 3 kb that flanked the upstream region of the histidine-autokinase domain and the downstream region of the aspartyl receiver domain. One of these clones, SA15.1, was chosen to determine the complete nucleotide sequence of the gene in both directions using the ABI automated sequencing system and fluorescent dideoxynucleotides as described earlier.

The DNA fragment generated by Slb1/Slb3 was used as a probe to screen a *C. albicans* EMBL3a lambda genomic library to identify clones containing the full-length gene. Of 10⁵ pfu's, twenty positive clones were identified. Clone λSA15.1, which contained a genomic fragment of approximately 4.8 kb with DNA flanking both the H1 and the D domains, was chosen for further characterization. The nucleotide sequence of the DNA insert was determined in both directions. The deduced amino acid sequence revealed an uninterrupted open reading frame of 1081 amino acids beginning with ATG as the initiation codon. The initiation codon was surrounded by an atypical Kozak consensus

sequence CTCCAATGA, with cytosine at the -3 position (Kozak, *Nucleic Acids Res*, 12: 857-871 (1984)). When total genomic DNA of *C. albicans* strain WO-1 was digested with a variety of restriction enzymes, and the resulting Southern blot hybridized under conditions of high stringency (65° C. in Church-Gilbert hybridization buffer) (Church & Gilbert, *Proc Natl Acad. Sci USA* 81: 1991-1995 (1984)) with the 1.2 kb probe spanning the 800 bp upstream of the gene, the banding pattern suggested that CaNik1 is encoded by a single copy gene. When total genomic DNA of strain WO-1 and strain 3153A was digested with BsaAI or NciI and hybridized with the 4.2 kb probe, the patterns were identical, but when TspI-digested DNA of the two strains were probed, the patterns differed, suggesting allelic differences exist between these strains. A comparison of the CaNik1 sequence published recently by Nagahashi et al., *Candida albicans. Microbiology*, 144: 425-432 (1998) for strain IF01060 and the sequence we obtained for strain WO-1 in the present invention differ at seven nucleotide positions in the open reading frame of 3243 bp.

EXAMPLE 3

Deletion of CaNik1 in *C. albicans* Strain CA18

In order to generate a CaNik1 deletion cassette, a DNA fragment of approximately 2.1 kb containing both the histidine-autokinase and aspartyl response regulator domains was amplified by PCR using as the template λSA15.1 (FIG. 4a), which contained the 545 bp sequence upstream of the histidine-autokinase domain. The PCR fragment was gel-purified and cloned into the PGEM-T easy vector (Promega). The DNA insert was again excised from the recombinant plasmid with EcoRI and subcloned into a PUC18 vector (Life Technologies) at the EcoRI site. The resultant recombinant plasmid was designated pUNIK12.1 (FIG. 4b). A deletion construct pCNH35 was generated that spanned the histidine-autokinase and ATP binding-domains. To construct pCNH35, pUNIK12.1 plasmid DNA (FIG. 4b) was digested with AflII and XhoI, and blunt-end repaired with the Klenow DNA polymerase I. The resultant plasmid DNA fragment was then gel purified and dephosphorylated with shrimp alkaline phosphatase (US Biochemical). A hisG-URA3-hisG cassette of 3.8 kb from pMB9 was then ligated to derive the disruption cassette (FIG. 4c). To isolate the CaNik1 disruption cassette from pCNH35, plasmid DNA was digested with PstI and the digested DNA extracted with phenol: chloroform. Approximately 25 μg of the digestion mixture was used to transform strain CA18, an ade2⁻ ura3⁻ derivative of wild type strain SC5314, by the lithium acetate protocol. Heterozygotes were selected for growth in minimal medium in the absence of uridine. Transformants were initially tested for the heterozygosity of one of the two CaNik1 alleles by Southern blot hybridization of genomic DNA digested with PstI. Positive heterozygotes were further confirmed by digesting genomic DNA with XhoI and by performing Southern blot hybridization. Because the genomic Southern revealed polymorphism between the two CaNik1 alleles, two distinct heterozygotes, NNL6 (L stands for large allele) and NNS7 (S stands for small allele) were selected. The heterozygote NNS7 was chosen to generate the knock-out for the second copy of the CaNik1 gene. Prior to the knock-out of the second copy, NNS7 was subjected to the 5-FOA selection protocol to convert it from uridine prototrophy to auxotrophy. Loss of the URA3 gene was again confirmed by digestion with XhoI and Southern blot analysis. In the final step, a single clone, NNS7.1.1, which was heterozygous for the L allele of the CaNik1 locus and URA3⁺, was subjected to a second round of transformation with pCNH35, and selected for growth on defined minimal medium lacking uridine. Transformants which had lost the second copy of CaNik1 were selected by Southern blot hybridization. One of the 125 transformants obtained with the pCNH35-based cassette, HH80, contained a homozygous deletion.

EXAMPLE 4

CaNik1 Transcription

To test whether transcription of CaNik1 was regulated by high frequency phenotypic switching, Northern blots of polyA⁺mRNA of white and opaque phase cell growth cultures of strain WO-1 were probed with the DNA fragment spanning the H1 and ATP binding domains of CaNik1. The CaNik1 transcript was detectable at very low levels in both white phase and opaque phase cells throughout the exponential phase of growth and in stationary phase. The level of CaNik1 transcript per cell remained constant throughout white phase cell growth, but increased steadily during opaque phase cell growth, reaching a level per cell roughly twice that of white phase cells at stationary phase (FIG. 5). Hypha-forming cells of both *C. albicans* strain WO-1 and *C. albicans* strain 3153A contained slightly higher levels of polyA⁺ CaNik1 transcript than budding cells. The hyphal-to-bud ratio of polyA⁺-containing CaNik1 transcript in strain WO-1 and strain 3153A was 1.2 and 1.3, respectively.

EXAMPLE 5

Functional Characterization of the CaNik1 Null Mutant of Strain CAI8

To test whether the CaNik1 deletion mutant HH80 underwent switching, we first had to characterize switching in this strain using a low dose ultraviolet irradiation protocol that increases switching frequencies. Cells were treated with ultraviolet irradiation for 0, 5, 10, 20 and 40 sec, and the percent kill as well as the frequency and type of switch variants were assessed on modified Lee's medium. The proportions of CAI8 and HH80 cells killed after 5, 10, 20, and 40 sec were similar. Identical variant phenotypes were stimulated by UV in both CAI8 and the homozygous deletion strain HH80. However, the frequency of variants induced by comparable levels of UV-irradiation was consistently lower in strain HH80, and this was true in a repeat experiment. For instance, 20 sec of UV irradiation resulted in 10.6% and 2.6% variants in CAI8 and HH80 cells, respectively. These results demonstrate that the CaNik1 gene product modulates phenotypic switching.

Since deletion of the *nik-1⁺* gene in *N. crassa* affects the morphology of hyphae, especially at high osmotic strength

(Alex et al., *Proc Natl Acad Sci USA*, 93: 3416–3421 (1996), the capability of the CaNik1-minus HH80 strain to form hyphae and the morphology of those hyphae were compared to that of the parent strain CAI8 and a URA3⁺ isogenic strain CAI8U5 at 0, 1.0 and 1.5 M NaCl. Under the regime of pH-regulated dimorphism, CAI8, CAI8U5, and HH80 cells formed buds at pH 4.5 and hyphae at pH 6.7. The kinetics of evagination for the three strains at low and high pH were similar at the three tested salt concentrations. At 1.5 M NaCl, the proportion of cells that formed evaginations at low and high pH was dramatically reduced in all three strains. The morphology of the hyphae that formed at pH 6.7 at 0, 1.0, and 1.5 M NaCl were comparable in the three strains. However, there was a significant and reproducible lag in hyphal growth at 1.5M NaCl in HH80 after 300 min. These results demonstrate that the CaNik1 gene product is not essential for hypha formation under the regime of pH regulated dimorphism, but its presence enhances hypha formation at high ionic strength.

Finally, growth of the CaNik1 deletion mutant HH80 was tested at 25° C. and 37° C. for differential sensitivity to osmotic strength and a variety of inhibitors. Patches of budding cells of CAI8, CAI8U5 and HH80 were plated on agar containing modified Lee's medium alone or with one of the following ingredients: 1.0 or 1.5M NaCl; 1M sorbitol; 0.8M KCl; 0.5M Mg₂SO₄; 20 or 40 μg per ml calcofluor; 1, 2 or 4 mg per ml caffeine; 10 or 20 mg per ml hygromycin; 0.002 or 0.004 μg per ml echinocandin; and 0.2 or 0.4M polymyxin B. In three independent experiments, no qualitative differences were observed between the growth of the control strains and the mutant strain HH80 for any of the tested conditions.

All publications and patent applications referred to in this specification are indicative of the level of skill of those in the art to which the invention pertains.

Other objects, features and advantages of the present invention will become apparent from the foregoing detailed description and examples. It should be understood, however, that the detailed description and the specific examples, while indicating preferred embodiments of the invention, are given only by way of illustration.

TABLE 1

Conditions used to test the effect of gene deletion	*Phenotypic effect in HH80
1. Growth kinetics in	
a) Lee's modified broth	Similar to SC5314, CAI8U5, and CAI8
b) YPD broth	Similar to SC5314, CAI8U5, and CAI8 ²⁰ .
2. Growth on agar plates with Lee's modified medium or YPD broth supplemented with:	
a) None	++++
b) 1 M NaCl	++
c) 1.5 M NaCl	+
d) 1 M KCl	++
e) 1.2 M Sorbitol	++++
f) 0.5 m MgSO ₄	++
g) Caffeine (1–4 mg/mL)	v
h) Calcofluor (20–40 μg/mL)	+++ ^v
i) Echinocandin (0.002–0.004 μg/mL)	±
j) 2% Trehalose	++++*
k) 2% Raffinose	++++*
l) 1 M Xylitol	++++
m) 10% Glycerol	++++*
3. Switching	
a) spontaneous frequency	No effect

TABLE 1-continued

Conditions used to test the effect of gene deletion	*Phenotypic effect in HH80
b) UV-stimulated frequency	Decreased
c) repertoire of switch phenotype	No effect
4. Hypha-induction under the regime of pH-regulated dimorphism. with no osmotic shock:	
a) time for 50% evagination	No effect
b) morphology of hypha	No effect
c) growth of hyphal filaments	
with osmotic shock using 1.5 M NaCl	
i) time for 50% evaginations	decreased in both wild type and the mutant
ii) morphology of hyphac	no difference between wild hyphae and the mutant
iii) growth of hyphal filaments	the growth of the hyphae after 300 min was reduced in the mutant as compared to that in wild type

In order to assess the effect of gene deletion on growth, exponentially grown cells of wild type (SC53 14), parental auxotrophic strain used to delete NIK1 gene (CAI8), URA3⁺ derivative of CAI8 (CAI8U5) and homozygous deletion mutant (HH80) were serially diluted and spot plated on agar plates with or without supplements in the medium. In all the growth medium used in this study, 2% glucose served as a carbon source except in the growth medium containing raffinose, trehalose and glycerol. The symbol "v" denote variable growth. Growth of the cultures were qualitatively assessed as very good (+++), good (++), fair (+), poor (+), poor to no growth (±). 0 indicates that colonies were very small (less than 1mm) as assessed by the colony size on agar plates spread with cultures to generate 50 to 100 individual colonies. The growth of the cultures were assessed after 2 or 3 days incubation both at 25° C. and 37° C.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 7

<210> SEQ ID NO 1

<211> LENGTH: 1254

<212> TYPE: DNA

<213> ORGANISM: Candida albicans

<220> FEATURE:

<221> NAME/KEY: CDS

<222> LOCATION: (1)..(1254)

<400> SEQUENCE: 1

```

gag att aga aca cca ttg aat ggg att att ggw atg acy cag ttg tcr      48
Glu Ile Arg Thr Pro Leu Asn Gly Ile Ile Gly Met Thr Gln Leu Ser
 1           5           10           15

cct gat aca gag ttg acr cag tac caa cga gag atg ttg tcg att gtg      96
Leu Asp Thr Glu Leu Thr Gln Tyr Gln Arg Glu Met Leu Ser Ile Val
          20           25           30

cat aac ttg gca aat tcc ttg ttg acc att ata gac gat ata ttg gat      144
His Asn Leu Ala Asn Ser Leu Leu Thr Ile Ile Asp Asp Ile Leu Asp
          35           40           45

att tct aag att gag gcg aat aga atg acg gtg gaa cag att gat ttt      192
Ile Ser Lys Ile Glu Ala Asn Arg Met Thr Val Glu Gln Ile Asp Phe
          50           55           60

tca tta aga ggg aca gtg ttt ggt gca ttg aaa acg tta gcc gtc aaa      240
Ser Leu Arg Gly Thr Val Phe Gly Ala Leu Lys Thr Leu Ala Val Lys
          65           70           75           80

gct att gaa aaa aac cta gac ttg acc tat caa tgt gat tca tcg ttt      288
Ala Ile Glu Lys Asn Leu Asp Leu Thr Tyr Gln Cys Asp Ser Ser Phe
          85           90           95

cca gat aat ctt att gga gat agt ttt aga tta cga caa gtt att ctt      336
Pro Asp Asn Leu Ile Gly Asp Ser Phe Arg Leu Arg Gln Val Ile Leu
          100          105          110

```

-continued

aac ttg gct ggt aat gct att aag ttt act aaa gag ggg aaa gtt agt Asn Leu Ala Gly Asn Ala Ile Lys Phe Thr Lys Glu Gly Lys Val Ser 115 120 125	384
gtt agt gtg aaa aag tct gat aaa atg gtg tta gat agt aag ttg ttg Val Ser Val Lys Lys Ser Asp Lys Met Val Leu Asp Ser Lys Leu Leu 130 135 140	432
tta gag gtt tgt gtt agc gac acg gga ata ggt ata gag aaa gac aaa Leu Glu Val Cys Val Ser Asp Thr Gly Ile Gly Ile Glu Lys Asp Lys 145 150 155 160	480
ttg gga ttg att ttc gat acc ttc tgt caa gct gat ggt tct act aca Leu Gly Leu Ile Phe Asp Thr Phe Cys Gln Ala Asp Gly Ser Thr Thr 165 170 175	528
aga aag ttt ggt ggt aca ggt tta ggg ttg tca att tcc aaa cag ttg Arg Lys Phe Gly Gly Thr Gly Leu Gly Leu Ser Ile Ser Lys Gln Leu 180 185 190	576
ata cat tta atg ggt gga gag ata tgg gtt act tcg gag tat gga tcc Ile His Leu Met Gly Gly Glu Ile Trp Val Thr Ser Glu Tyr Gly Ser 195 200 205	624
ggr tca aac ttt tat ttt acg gtg tgc gtg tcg cca tct aat att aga Gly Ser Asn Phe Tyr Phe Thr Val Cys Val Ser Pro Ser Asn Ile Arg 210 215 220	672
tat act cga caa acc gaa caa ttg tta cca ttt agt tcc cat tat gtg Tyr Thr Arg Gln Thr Glu Gln Leu Leu Pro Phe Ser Ser His Tyr Val 225 230 235 240	720
tta ttt gta tcg act gag cat act caa gaa gaa ctt gat gtg ttg aga Leu Phe Val Ser Thr Glu His Thr Gln Glu Glu Leu Asp Val Leu Arg 245 250 255	768
gat gga att ata gaa ctt gga ttg ata cct ata ata gtg aga aat att Asp Gly Ile Ile Glu Leu Gly Leu Ile Pro Ile Ile Val Arg Asn Ile 260 265 270	816
gaa gat gca aca ttg act gag ccg gtg aaa tat gat ata att atg att Glu Asp Ala Thr Leu Thr Glu Pro Val Lys Tyr Asp Ile Ile Met Ile 275 280 285	864
gat tcg ata gag att gcc aaa aag ttg agg ttg tta tcg gag gtt aaa Asp Ser Ile Glu Ile Ala Lys Lys Leu Arg Leu Leu Ser Glu Val Lys 290 295 300	912
tat att ccg ttg gtt ttg gtc cat cat tct att cca cag ttg aat atg Tyr Ile Pro Leu Val Leu Val His His Ser Ile Pro Gln Leu Asn Met 305 310 315 320	960
aga gta tgt att gat ttg ggg ata tct tcc tat gca aat acg cca tgt Arg Val Cys Ile Asp Leu Gly Ile Ser Ser Tyr Ala Asn Thr Pro Cys 325 330 335	1008
tcg atc acg gac ttg gcc agt gcg att ata cca gcg ttg gag tcg aga Ser Ile Thr Asp Leu Ala Ser Ala Ile Ile Pro Ala Leu Glu Ser Arg 340 345 350	1056
tct ata tca cag aac tca gac gag tcg gtg agg tac aaa ata tta cta Ser Ile Ser Gln Asn Ser Asp Glu Ser Val Arg Tyr Lys Ile Leu Leu 355 360 365	1104
gca gag gac aac ctc gtc aat cag aaa ctt gca gtt agg ata tta gaa Ala Glu Asp Asn Leu Val Asn Gln Lys Leu Ala Val Arg Ile Leu Glu 370 375 380	1152
aag caa ggg cat ctg gtg gaa gta gtt gag aac gga ctc gag gcg tac Lys Gln Gly His Leu Val Glu Val Val Glu Asn Gly Leu Glu Ala Tyr 385 390 395 400	1200
gaa gcg att aag agg aat aaa tat gat gtg gtg ttg atg gat gtg caa Glu Ala Ile Lys Arg Asn Lys Tyr Asp Val Val Leu Met Asp Val Gln 405 410 415	1248
atg cct Met Pro	1254

-continued

```

<210> SEQ ID NO 2
<211> LENGTH: 418
<212> TYPE: PRT
<213> ORGANISM: Candida albicans

<400> SEQUENCE: 2

Glu Ile Arg Thr Pro Leu Asn Gly Ile Ile Gly Met Thr Gln Leu Ser
 1           5           10
Leu Asp Thr Glu Leu Thr Gln Tyr Gln Arg Glu Met Leu Ser Ile Val
          20           25           30
His Asn Leu Ala Asn Ser Leu Leu Thr Ile Ile Asp Asp Ile Leu Asp
          35           40           45
Ile Ser Lys Ile Glu Ala Asn Arg Met Thr Val Glu Gln Ile Asp Phe
          50           55           60
Ser Leu Arg Gly Thr Val Phe Gly Ala Leu Lys Thr Leu Ala Val Lys
          65           70           75           80
Ala Ile Glu Lys Asn Leu Asp Leu Thr Tyr Gln Cys Asp Ser Ser Phe
          85           90           95
Pro Asp Asn Leu Ile Gly Asp Ser Phe Arg Leu Arg Gln Val Ile Leu
          100          105          110
Asn Leu Ala Gly Asn Ala Ile Lys Phe Thr Lys Glu Gly Lys Val Ser
          115          120          125
Val Ser Val Lys Lys Ser Asp Lys Met Val Leu Asp Ser Lys Leu Leu
          130          135          140
Leu Glu Val Cys Val Ser Asp Thr Gly Ile Gly Ile Glu Lys Asp Lys
          145          150          155          160
Leu Gly Leu Ile Phe Asp Thr Phe Cys Gln Ala Asp Gly Ser Thr Thr
          165          170          175
Arg Lys Phe Gly Gly Thr Gly Leu Gly Leu Ser Ile Ser Lys Gln Leu
          180          185          190
Ile His Leu Met Gly Gly Glu Ile Trp Val Thr Ser Glu Tyr Gly Ser
          195          200          205
Gly Ser Asn Phe Tyr Phe Thr Val Cys Val Ser Pro Ser Asn Ile Arg
          210          215          220
Tyr Thr Arg Gln Thr Glu Gln Leu Leu Pro Phe Ser Ser His Tyr Val
          225          230          235          240
Leu Phe Val Ser Thr Glu His Thr Gln Glu Glu Leu Asp Val Leu Arg
          245          250          255
Asp Gly Ile Ile Glu Leu Gly Leu Ile Pro Ile Ile Val Arg Asn Ile
          260          265          270
Glu Asp Ala Thr Leu Thr Glu Pro Val Lys Tyr Asp Ile Ile Met Ile
          275          280          285
Asp Ser Ile Glu Ile Ala Lys Lys Leu Arg Leu Leu Ser Glu Val Lys
          290          295          300
Tyr Ile Pro Leu Val Leu Val His His Ser Ile Pro Gln Leu Asn Met
          305          310          315          320
Arg Val Cys Ile Asp Leu Gly Ile Ser Ser Tyr Ala Asn Thr Pro Cys
          325          330          335
Ser Ile Thr Asp Leu Ala Ser Ala Ile Ile Pro Ala Leu Glu Ser Arg
          340          345          350
Ser Ile Ser Gln Asn Ser Asp Glu Ser Val Arg Tyr Lys Ile Leu Leu
          355          360          365
Ala Glu Asp Asn Leu Val Asn Gln Lys Leu Ala Val Arg Ile Leu Glu
          370          375          380

```

-continued

Lys Gln Gly His Leu Val Glu Val Val Glu Asn Gly Leu Glu Ala Tyr
385 390 395 400

Glu Ala Ile Lys Arg Asn Lys Tyr Asp Val Val Leu Met Asp Val Gln
405 410 415

Met Pro

<210> SEQ ID NO 3
<211> LENGTH: 3246
<212> TYPE: DNA
<213> ORGANISM: Candida albicans
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(3243)

<400> SEQUENCE: 3

```

atg aac ccc act aaa aaa cct cgg tta tca cca atg cag ccc tct gtt      48
Met Asn Pro Thr Lys Lys Pro Arg Leu Ser Pro Met Gln Pro Ser Val
  1          5          10          15

ttt gaa ata ctc aac gac cct gag ctt tat agt cag cac tgt cat agc      96
Phe Glu Ile Leu Asn Asp Pro Glu Leu Tyr Ser Gln His Cys His Ser
          20          25          30

ctt agg gaa aca ctt ctt gat cat ttc aac cat caa gct aca ctt atc     144
Leu Arg Glu Thr Leu Leu Asp His Phe Asn His Gln Ala Thr Leu Ile
          35          40          45

gac act tat gaa cat gaa cta gaa aaa tcc aaa aat gcc aac aaa gcg     192
Asp Thr Tyr Glu His Glu Leu Glu Lys Ser Lys Asn Ala Asn Lys Ala
          50          55          60

tcc caa caa gca ctt agt gaa ata ggt aca gtt gtt ata tct gtt gcc     240
Ser Gln Gln Ala Leu Ser Glu Ile Gly Thr Val Val Ile Ser Val Ala
          65          70          75          80

atg gga gac ttg tgg aaa aaa gtt gag att cac aca gta gaa aat gac     288
Met Gly Asp Leu Ser Lys Lys Val Glu Ile His Thr Val Glu Asn Asp
          85          90          95

cct gag att tta aaa gtc aaa atc acc atc aac acc atg atg gat caa     336
Pro Glu Ile Leu Lys Val Lys Ile Thr Ile Asn Thr Met Met Asp Gln
          100          105          110

tta cag aca ttt gct aat gag gtt aca aaa gtc gcc acc gaa gtc gca     384
Leu Gln Thr Phe Ala Asn Glu Val Thr Lys Val Ala Thr Glu Val Ala
          115          120          125

aat ggt gaa cta ggt gga caa gcg aaa aat gat gga tct gtt ggt att     432
Asn Gly Glu Leu Gly Gly Gln Ala Lys Asn Asp Gly Ser Val Gly Ile
          130          135          140

tgg aga tca ctt aca gac aat gtt aat att atg gct ctt aat tta act     480
Trp Arg Ser Leu Thr Asp Asn Val Asn Ile Met Ala Leu Asn Leu Thr
          145          150          155          160

aac caa gtg cga gaa att gct gat gtc aca cgt gct gtt gcc aag ggg     528
Asn Gln Val Arg Glu Ile Ala Asp Val Thr Arg Ala Val Ala Lys Gly
          165          170          175

gac ttg tca cgt aaa att aat gta cac gcc cag ggt gaa atc ctt caa     576
Asp Leu Ser Arg Lys Ile Asn Val His Ala Gln Gly Glu Ile Leu Gln
          180          185          190

ctt caa cgt aca ata aac acc atg gtg gat cag tta cga acg ttt gca     624
Leu Gln Arg Thr Ile Asn Thr Met Val Asp Gln Leu Arg Thr Phe Ala
          195          200          205

ttc gaa gta tct aaa gtt gct aga gat gtt ggt gtg ctt ggt ata tta     672
Phe Glu Val Ser Lys Val Ala Arg Asp Val Gly Val Leu Gly Ile Leu
          210          215          220

gga gga caa gcg ttg att gaa aat gtt gaa ggt att tgg gaa gag ttg     720
Gly Gly Gln Ala Leu Ile Glu Asn Val Glu Gly Ile Trp Glu Glu Leu
          225          230          235          240
    
```

-continued

act gat aat gtc aat gcc atg gct ctt aat ttg act aca caa gtg aga Thr Asp Asn Val Asn Ala Met Ala Leu Asn Leu Thr Thr Gln Val Arg 245 250 255	768
aat att gcc aat gtc acc act gcc gtt gcc aag ggg gat ttg tcg aaa Asn Ile Ala Asn Val Thr Thr Ala Val Ala Lys Gly Asp Leu Ser Lys 260 265 270	816
aaa gtc act gct gat tgt aag gga gaa aty ctt gat ttg aaa ctt act Lys Val Thr Ala Asp Cys Lys Gly Glu Ile Leu Asp Leu Lys Leu Thr 275 280 285	864
att aat caa atg gtg gac cga tta cag aat ttt gct ctt gcg gtg acg Ile Asn Gln Met Val Asp Arg Leu Gln Asn Phe Ala Leu Ala Val Thr 290 295 300	912
aca ttg tcg aga gag gtt ggt act ttg ggt att ttg ggt gga caa gct Thr Leu Ser Arg Glu Val Gly Thr Leu Gly Ile Leu Gly Gly Gln Ala 305 310 315 320	960
aac gta cag gat gtt gaa ggt gct tgg aaa cag gtt aca gaa aat gtc Asn Val Gln Asp Val Glu Gly Ala Trp Lys Gln Val Thr Glu Asn Val 325 330 335	1008
aac cta atg gct act aat tta act aac caa gtg aga tct att gct aca Asn Leu Met Ala Thr Asn Leu Thr Asn Gln Val Arg Ser Ile Ala Thr 340 345 350	1056
gtt act act gca gtt gcg cat ggt gat ttg tcg caa aag att gat ggt Val Thr Thr Ala Val Ala His Gly Asp Leu Ser Gln Lys Ile Asp Gly 355 360 365	1104
cat ccc aaa gga gag att tta caa ttg aaa aat aca atc aac aag atg His Pro Lys Gly Glu Ile Leu Gln Leu Lys Asn Thr Ile Asn Lys Met 370 375 380	1152
gtg gac tct ttg cag ttg ttt gca tca gaa gtg tcg aaa gtg gca caa Val Asp Ser Leu Gln Leu Phe Ala Ser Glu Val Ser Lys Val Ala Gln 385 390 395 400	1200
gat gtt ggt att aat gga aaa tta ggt att caa gca caa gtt agt gat Asp Val Gly Ile Asn Gly Lys Leu Gly Ile Gln Ala Gln Val Ser Asp 405 410 415	1248
gtt gat gga tta tgg aag gag att acg tct aat gta aat acc atg gct Val Asp Gly Leu Trp Lys Glu Ile Thr Ser Asn Val Asn Thr Met Ala 420 425 430	1296
tca aat tta act tcg caa gtg aga gct ttt gca cag att act gct gct Ser Asn Leu Thr Ser Gln Val Arg Ala Phe Ala Gln Ile Thr Ala Ala 435 440 445	1344
gct act gat ggg gat ttc act aga ttt att act gtt gaa gca ctg gga Ala Thr Asp Gly Asp Phe Thr Arg Phe Ile Thr Val Glu Ala Leu Gly 450 455 460	1392
gag atg gat gcg ttg aaa aca aag att aat caa atg gtg ttt aac tta Glu Met Asp Ala Leu Lys Thr Lys Ile Asn Gln Met Val Phe Asn Leu 465 470 475 480	1440
agg gaa tcg ctt caa agg aat act gcg gct aga gaa gct gct gag ttg Arg Glu Ser Leu Gln Arg Asn Thr Ala Ala Arg Glu Ala Ala Glu Leu 485 490 495	1488
gcc aat agt gcg aaa tcc gag ttt tta gca aac atg tcg cat gag att Ala Asn Ser Ala Lys Ser Glu Phe Leu Ala Asn Met Ser His Glu Ile 500 505 510	1536
aga aca cca ttg aat ggg att att gdw atg acy cag ttg tcr ctt gat Arg Thr Pro Leu Asn Gly Ile Ile Gly Met Thr Gln Leu Ser Leu Asp 515 520 525	1584
aca gag ttg acr cag tac caa cga gag atg ttg tcg att gtg cat aac Thr Glu Leu Thr Gln Tyr Gln Arg Glu Met Leu Ser Ile Val His Asn 530 535 540	1632
ttg gca aat tcc ttg ttg acc att ata gac gat ata ttg gat att tct Leu Ala Asn Ser Leu Leu Thr Ile Ile Asp Asp Ile Leu Asp Ile Ser	1680

-continued

545	550	555	560	
aag att gag gcg aat aga atg acg gtg gaa cag att gat ttt tca tta Lys Ile Glu Ala Asn Arg Met Thr Val Glu Gln Ile Asp Phe Ser Leu	565	570	575	1728
aga ggg aca gtg ttt ggt gca ttg aaa acg tta gcc gtc aaa gct att Arg Gly Thr Val Phe Gly Ala Leu Lys Thr Leu Ala Val Lys Ala Ile	580	585	590	1776
gaa aaa aac cta gac ttg acc tat caa tgt gat tca tcg ttt cca gat Glu Lys Asn Leu Asp Leu Thr Tyr Gln Cys Asp Ser Ser Phe Pro Asp	595	600	605	1824
aat ctt att gga gat agt ttt aga tta cga caa gtt att ctt aac ttg Asn Leu Ile Gly Asp Ser Phe Arg Leu Arg Gln Val Ile Leu Asn Leu	610	615	620	1872
gct ggt aat gct att aag ttt act aaa gag ggg aaa gtt agt gtt agt Ala Gly Asn Ala Ile Lys Phe Thr Lys Glu Gly Lys Val Ser Val Ser	625	630	635	1920
gtg aaa aag tct gat aaa atg gtg tta gat agt aag ttg ttg tta gag Val Lys Lys Ser Asp Lys Met Val Leu Asp Ser Lys Leu Leu Leu Glu	645	650	655	1968
gtt tgt gtt agc gac acg gga ata ggt ata gag aaa gac aaa ttg gga Val Cys Val Ser Asp Thr Gly Ile Gly Ile Glu Lys Asp Lys Leu Gly	660	665	670	2016
ttg att ttc gat acc ttc tgt caa gct gat ggt tct act aca aga aag Leu Ile Phe Asp Thr Phe Cys Gln Ala Asp Gly Ser Thr Thr Arg Lys	675	680	685	2064
ttt ggt ggt aca ggt tta ggg ttg tca att tcc aaa cag ttg ata cat Phe Gly Gly Thr Gly Leu Gly Leu Ser Ile Ser Lys Gln Leu Ile His	690	695	700	2112
tta atg ggt gga gag ata tgg gtt act tcg gag tat gga tcc ggr tca Leu Met Gly Gly Glu Ile Trp Val Thr Ser Glu Tyr Gly Ser Gly Ser	705	710	715	2160
aac ttt tat ttt acg gtg tgc gtg tcg cca tct aat att aga tat act Asn Phe Tyr Phe Thr Val Cys Val Ser Pro Ser Asn Ile Arg Tyr Thr	725	730	735	2208
cga caa acc gaa caa ttg tta cca ttt agt tcc cat tat gtg tta ttt Arg Gln Thr Glu Gln Leu Leu Pro Phe Ser Ser His Tyr Val Leu Phe	740	745	750	2256
gta tcg act gag cat act caa gaa gaa ctt gat gtg ttg aga gat gga Val Ser Thr Glu His Thr Gln Glu Glu Leu Asp Val Leu Arg Asp Gly	755	760	765	2304
att ata gaa ctt gga ttg ata cct ata ata gtg aga aat att gaa gat Ile Ile Glu Leu Gly Leu Ile Pro Ile Ile Val Arg Asn Ile Glu Asp	770	775	780	2352
gca aca ttg act gag ccg gtg aaa tat gat ata att atg att gat tcg Ala Thr Leu Thr Glu Pro Val Lys Tyr Asp Ile Ile Met Ile Asp Ser	785	790	795	2400
ata gag att gcc aaa aag ttg agg ttg tta tcg gag gtt aaa tat att Ile Glu Ile Ala Lys Lys Leu Arg Leu Leu Ser Glu Val Lys Tyr Ile	805	810	815	2448
ccg ttg gtt ttg gtc cat cat tct att cca cag ttg aat atg aga gta Pro Leu Val Leu Val His His Ser Ile Pro Gln Leu Asn Met Arg Val	820	825	830	2496
tgt att gat ttg ggg ata tct tcc tat gca aat acg cca tgt tcg atc Cys Ile Asp Leu Gly Ile Ser Ser Tyr Ala Asn Thr Pro Cys Ser Ile	835	840	845	2544
acg gac ttg gcc agt gcg att ata cca gcg ttg gag tcg aga tct ata Thr Asp Leu Ala Ser Ala Ile Ile Pro Ala Leu Glu Ser Arg Ser Ile	850	855	860	2592
tca cag aac tca gac gag tcg gtg agg tac aaa ata tta cta gca gag				2640

-continued

Ser	Gln	Asn	Ser	Asp	Glu	Ser	Val	Arg	Tyr	Lys	Ile	Leu	Leu	Ala	Glu		
865					870					875					880		
gac	aac	ctc	gtc	aat	cag	aaa	ctt	gca	ggt	agg	ata	tta	gaa	aag	caa	2688	
Asp	Asn	Leu	Val	Asn	Gln	Lys	Leu	Ala	Val	Arg	Ile	Leu	Glu	Lys	Gln		
				885					890					895			
ggg	cat	ctg	gtg	gaa	gta	ggt	gag	aac	gga	ctc	gag	gcg	tac	gaa	gcg	2736	
Gly	His	Leu	Val	Glu	Val	Val	Glu	Asn	Gly	Leu	Glu	Ala	Tyr	Glu	Ala		
			900					905					910				
att	aag	agg	aat	aaa	tat	gat	gtg	gtg	ttg	atg	gat	gtg	caa	atg	cct	2784	
Ile	Lys	Arg	Asn	Lys	Tyr	Asp	Val	Val	Leu	Met	Asp	Val	Gln	Met	Pro		
		915					920					925					
gta	atg	ggt	ggg	ttt	gaa	gct	acg	gag	aag	att	cga	caa	tgg	gag	aaa	2832	
Val	Met	Gly	Gly	Phe	Glu	Ala	Thr	Glu	Lys	Ile	Arg	Gln	Trp	Glu	Lys		
		930				935					940						
aag	tct	aac	cca	att	gac	tcg	ttg	acc	ttt	agg	act	cca	att	att	gcc	2880	
Lys	Ser	Asn	Pro	Ile	Asp	Ser	Leu	Thr	Phe	Arg	Thr	Pro	Ile	Ile	Ala		
					950					955					960		
ctc	act	gca	cac	gcc	atg	tta	ggt	gat	aga	gaa	aag	tca	ttg	gcc	aag	2928	
Leu	Thr	Ala	His	Ala	Met	Leu	Gly	Asp	Arg	Glu	Lys	Ser	Leu	Ala	Lys		
				965					970					975			
ggg	atg	gac	gat	tat	gtg	agt	aag	cca	ttg	aag	ccg	aaa	ttg	tta	atg	2976	
Gly	Met	Asp	Asp	Tyr	Val	Ser	Lys	Pro	Leu	Lys	Pro	Lys	Leu	Leu	Met		
			980					985					990				
cag	acg	ata	aag	aag	tgt	att	cat	aat	att	aac	cag	ttg	aaa	gaa	ttg	3024	
Gln	Thr	Ile	Lys	Lys	Cys	Ile	His	Asn	Ile	Asn	Gln	Leu	Lys	Glu	Leu		
		995				1000						1005					
tcg	aga	aat	agt	agg	ggt	agc	gat	ttt	gca	aag	aag	atg	acc	cga	aac	3072	
Ser	Arg	Asn	Ser	Arg	Gly	Ser	Asp	Phe	Ala	Lys	Lys	Met	Thr	Arg	Asn		
		1010			1015							1020					
aca	ccc	ggc	cgc	acg	acc	cgt	cag	ggg	agt	gat	gag	ggg	agt	gta	aag	3120	
Thr	Pro	Gly	Arg	Thr	Thr	Arg	Gln	Gly	Ser	Asp	Glu	Gly	Ser	Val	Lys		
				1025		1030				1035				1040			
gac	atg	att	ggg	gac	act	ccc	cgt	caa	ggg	agt	gtg	gag	gga	ggg	ggt	3168	
Asp	Met	Ile	Gly	Asp	Thr	Pro	Arg	Gln	Gly	Ser	Val	Glu	Gly	Gly	Gly		
			1045					1050					1055				
aca	agt	agt	aga	cca	gta	cag	aga	agg	tct	gcc	agg	gag	ggg	tcg	atc	3216	
Thr	Ser	Ser	Arg	Pro	Val	Gln	Arg	Arg	Ser	Ala	Arg	Glu	Gly	Ser	Ile		
			1060				1065						1070				
act	aca	att	agt	gaa	caa	atc	gac	cgt	tag							3246	
Thr	Thr	Ile	Ser	Glu	Gln	Ile	Asp	Arg									
		1075				1080											

<210> SEQ ID NO 4
 <211> LENGTH: 1081
 <212> TYPE: PRT
 <213> ORGANISM: Candida albicans

<400> SEQUENCE: 4

Met	Asn	Pro	Thr	Lys	Lys	Pro	Arg	Leu	Ser	Pro	Met	Gln	Pro	Ser	Val		
1				5					10					15			
Phe	Glu	Ile	Leu	Asn	Asp	Pro	Glu	Leu	Tyr	Ser	Gln	His	Cys	His	Ser		
			20					25					30				
Leu	Arg	Glu	Thr	Leu	Leu	Asp	His	Phe	Asn	His	Gln	Ala	Thr	Leu	Ile		
		35				40						45					
Asp	Thr	Tyr	Glu	His	Glu	Leu	Glu	Lys	Ser	Lys	Asn	Ala	Asn	Lys	Ala		
	50					55					60						
Ser	Gln	Gln	Ala	Leu	Ser	Glu	Ile	Gly	Thr	Val	Val	Ile	Ser	Val	Ala		
	65				70					75					80		
Met	Gly	Asp	Leu	Ser	Lys	Lys	Val	Glu	Ile	His	Thr	Val	Glu	Asn	Asp		

-continued

85					90					95					
Pro	Glu	Ile	Leu	Lys	Val	Lys	Ile	Thr	Ile	Asn	Thr	Met	Met	Asp	Gln
			100					105					110		
Leu	Gln	Thr	Phe	Ala	Asn	Glu	Val	Thr	Lys	Val	Ala	Thr	Glu	Val	Ala
		115						120				125			
Asn	Gly	Glu	Leu	Gly	Gly	Gln	Ala	Lys	Asn	Asp	Gly	Ser	Val	Gly	Ile
	130					135					140				
Trp	Arg	Ser	Leu	Thr	Asp	Asn	Val	Asn	Ile	Met	Ala	Leu	Asn	Leu	Thr
145					150					155					160
Asn	Gln	Val	Arg	Glu	Ile	Ala	Asp	Val	Thr	Arg	Ala	Val	Ala	Lys	Gly
				165					170					175	
Asp	Leu	Ser	Arg	Lys	Ile	Asn	Val	His	Ala	Gln	Gly	Glu	Ile	Leu	Gln
			180					185					190		
Leu	Gln	Arg	Thr	Ile	Asn	Thr	Met	Val	Asp	Gln	Leu	Arg	Thr	Phe	Ala
		195					200					205			
Phe	Glu	Val	Ser	Lys	Val	Ala	Arg	Asp	Val	Gly	Val	Leu	Gly	Ile	Leu
	210					215					220				
Gly	Gly	Gln	Ala	Leu	Ile	Glu	Asn	Val	Glu	Gly	Ile	Trp	Glu	Glu	Leu
225					230					235					240
Thr	Asp	Asn	Val	Asn	Ala	Met	Ala	Leu	Asn	Leu	Thr	Thr	Gln	Val	Arg
				245					250					255	
Asn	Ile	Ala	Asn	Val	Thr	Thr	Ala	Val	Ala	Lys	Gly	Asp	Leu	Ser	Lys
			260					265					270		
Lys	Val	Thr	Ala	Asp	Cys	Lys	Gly	Glu	Ile	Leu	Asp	Leu	Lys	Leu	Thr
		275					280					285			
Ile	Asn	Gln	Met	Val	Asp	Arg	Leu	Gln	Asn	Phe	Ala	Leu	Ala	Val	Thr
	290					295					300				
Thr	Leu	Ser	Arg	Glu	Val	Gly	Thr	Leu	Gly	Ile	Leu	Gly	Gly	Gln	Ala
305					310					315					320
Asn	Val	Gln	Asp	Val	Glu	Gly	Ala	Trp	Lys	Gln	Val	Thr	Glu	Asn	Val
				325					330					335	
Asn	Leu	Met	Ala	Thr	Asn	Leu	Thr	Asn	Gln	Val	Arg	Ser	Ile	Ala	Thr
			340					345					350		
Val	Thr	Thr	Ala	Val	Ala	His	Gly	Asp	Leu	Ser	Gln	Lys	Ile	Asp	Gly
		355					360					365			
His	Pro	Lys	Gly	Glu	Ile	Leu	Gln	Leu	Lys	Asn	Thr	Ile	Asn	Lys	Met
	370					375					380				
Val	Asp	Ser	Leu	Gln	Leu	Phe	Ala	Ser	Glu	Val	Ser	Lys	Val	Ala	Gln
385					390					395					400
Asp	Val	Gly	Ile	Asn	Gly	Lys	Leu	Gly	Ile	Gln	Ala	Gln	Val	Ser	Asp
				405					410					415	
Val	Asp	Gly	Leu	Trp	Lys	Glu	Ile	Thr	Ser	Asn	Val	Asn	Thr	Met	Ala
			420					425					430		
Ser	Asn	Leu	Thr	Ser	Gln	Val	Arg	Ala	Phe	Ala	Gln	Ile	Thr	Ala	Ala
		435					440					445			
Ala	Thr	Asp	Gly	Asp	Phe	Thr	Arg	Phe	Ile	Thr	Val	Glu	Ala	Leu	Gly
	450					455					460				
Glu	Met	Asp	Ala	Leu	Lys	Thr	Lys	Ile	Asn	Gln	Met	Val	Phe	Asn	Leu
465					470					475					480
Arg	Glu	Ser	Leu	Gln	Arg	Asn	Thr	Ala	Ala	Arg	Glu	Ala	Ala	Glu	Leu
				485					490					495	
Ala	Asn	Ser	Ala	Lys	Ser	Glu	Phe	Leu	Ala	Asn	Met	Ser	His	Glu	Ile
			500					505					510		

-continued

Arg Thr Pro Leu Asn Gly Ile Ile Gly Met Thr Gln Leu Ser Leu Asp
 515 520 525

Thr Glu Leu Thr Gln Tyr Gln Arg Glu Met Leu Ser Ile Val His Asn
 530 535 540

Leu Ala Asn Ser Leu Leu Thr Ile Ile Asp Asp Ile Leu Asp Ile Ser
 545 550 555 560

Lys Ile Glu Ala Asn Arg Met Thr Val Glu Gln Ile Asp Phe Ser Leu
 565 570 575

Arg Gly Thr Val Phe Gly Ala Leu Lys Thr Leu Ala Val Lys Ala Ile
 580 585 590

Glu Lys Asn Leu Asp Leu Thr Tyr Gln Cys Asp Ser Ser Phe Pro Asp
 595 600 605

Asn Leu Ile Gly Asp Ser Phe Arg Leu Arg Gln Val Ile Leu Asn Leu
 610 615 620

Ala Gly Asn Ala Ile Lys Phe Thr Lys Glu Gly Lys Val Ser Val Ser
 625 630 635 640

Val Lys Lys Ser Asp Lys Met Val Leu Asp Ser Lys Leu Leu Glu
 645 650 655

Val Cys Val Ser Asp Thr Gly Ile Gly Ile Glu Lys Asp Lys Leu Gly
 660 665 670

Leu Ile Phe Asp Thr Phe Cys Gln Ala Asp Gly Ser Thr Thr Arg Lys
 675 680 685

Phe Gly Gly Thr Gly Leu Gly Leu Ser Ile Ser Lys Gln Leu Ile His
 690 695 700

Leu Met Gly Gly Glu Ile Trp Val Thr Ser Glu Tyr Gly Ser Gly Ser
 705 710 715 720

Asn Phe Tyr Phe Thr Val Cys Val Ser Pro Ser Asn Ile Arg Tyr Thr
 725 730 735

Arg Gln Thr Glu Gln Leu Leu Pro Phe Ser Ser His Tyr Val Leu Phe
 740 745 750

Val Ser Thr Glu His Thr Gln Glu Leu Asp Val Leu Arg Asp Gly
 755 760 765

Ile Ile Glu Leu Gly Leu Ile Pro Ile Ile Val Arg Asn Ile Glu Asp
 770 775 780

Ala Thr Leu Thr Glu Pro Val Lys Tyr Asp Ile Ile Met Ile Asp Ser
 785 790 795 800

Ile Glu Ile Ala Lys Lys Leu Arg Leu Leu Ser Glu Val Lys Tyr Ile
 805 810 815

Pro Leu Val Leu Val His His Ser Ile Pro Gln Leu Asn Met Arg Val
 820 825 830

Cys Ile Asp Leu Gly Ile Ser Ser Tyr Ala Asn Thr Pro Cys Ser Ile
 835 840 845

Thr Asp Leu Ala Ser Ala Ile Ile Pro Ala Leu Glu Ser Arg Ser Ile
 850 855 860

Ser Gln Asn Ser Asp Glu Ser Val Arg Tyr Lys Ile Leu Leu Ala Glu
 865 870 875 880

Asp Asn Leu Val Asn Gln Lys Leu Ala Val Arg Ile Leu Glu Lys Gln
 885 890 895

Gly His Leu Val Glu Val Val Glu Asn Gly Leu Glu Ala Tyr Glu Ala
 900 905 910

Ile Lys Arg Asn Lys Tyr Asp Val Val Leu Met Asp Val Gln Met Pro
 915 920 925

-continued

Val Met Gly Gly Phe Glu Ala Thr Glu Lys Ile Arg Gln Trp Glu Lys
 930 935 940

Lys Ser Asn Pro Ile Asp Ser Leu Thr Phe Arg Thr Pro Ile Ile Ala
 945 950 955 960

Leu Thr Ala His Ala Met Leu Gly Asp Arg Glu Lys Ser Leu Ala Lys
 965 970 975

Gly Met Asp Asp Tyr Val Ser Lys Pro Leu Lys Pro Lys Leu Leu Met
 980 985 990

Gln Thr Ile Lys Lys Cys Ile His Asn Ile Asn Gln Leu Lys Glu Leu
 995 1000 1005

Ser Arg Asn Ser Arg Gly Ser Asp Phe Ala Lys Lys Met Thr Arg Asn
 1010 1015 1020

Thr Pro Gly Arg Thr Thr Arg Gln Gly Ser Asp Glu Gly Ser Val Lys
 1025 1030 1035 1040

Asp Met Ile Gly Asp Thr Pro Arg Gln Gly Ser Val Glu Gly Gly Gly
 1045 1050 1055

Thr Ser Ser Arg Pro Val Gln Arg Arg Ser Ala Arg Glu Gly Ser Ile
 1060 1065 1070

Thr Thr Ile Ser Glu Gln Ile Asp Arg
 1075 1080

<210> SEQ ID NO 5
 <211> LENGTH: 23
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Degenerate
 primer
 <221> NAME/KEY: modified_base
 <222> LOCATION: (16)
 <223> OTHER INFORMATION: i
 <221> NAME/KEY: modified_base
 <222> LOCATION: (18)
 <223> OTHER INFORMATION: i
 <400> SEQUENCE: 5

gaattgagaa cgctntnaa tgg 23

<210> SEQ ID NO 6
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Degenerate
 primer
 <221> NAME/KEY: modified_base
 <222> LOCATION: (3)
 <223> OTHER INFORMATION: i
 <400> SEQUENCE: 6

agncctaagc cagtaccacc 20

<210> SEQ ID NO 7
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Degenerate
 primer
 <221> NAME/KEY: modified_base
 <222> LOCATION: (16)

-continued

<223> OTHER INFORMATION: i

<400> SEQUENCE: 7

tttaggcattc tggacntcca t

21

What is claimed is:

1. An isolated polynucleotide that encodes a protein linked to phenotypic switching in *Candida albicans* that exhibits 70% or greater overall sequence identity to SEQ. ID No. 3, wherein said protein displays kinase activity.

2. The polynucleotide of claim 1 that exhibits 80% or greater identity to SEQ ID No 3.

3. The polynucleotide of claim 1 that exhibits 90% or greater identity to SEQ ID NO 3.

4. A polynucleotide according to claim 1, comprising the sequence of SEQ ID No. 3.

5. A method of screening for a compound with the ability to inhibit expression or functionality of the CaNIK1 protein comprising:

(A) contacting a yeast cell that exhibits phenotypic switching with a test substance, wherein said yeast cell comprises:

- (i) a polynucleotide according to claim 1 and
- (ii) a promoter operably linked to said polynucleotide, such that said yeast cell produces a protein encoded by said polynucleotide; then

(B) monitoring the ability of said test substance to inhibit expression or functionality of said protein encoded by said polynucleotide in said yeast cell.

6. The method according to claim 5, wherein step (B) comprises monitoring the level of said protein produced in said cell.

7. The method according to claim 6, wherein step (B) comprises effecting a two-dimensional gel electrophoresis.

10 8. The method according to claim 5, wherein step (B) comprises monitoring the level of m RNA encoded by said polynucleotide and produced by said cell.

15 9. The method according to claim 8, wherein step (B) comprises effecting a Northern blot, a primer extension, or a ribonuclease protection assay.

10. The method according to claim 5, wherein step (B) comprises monitoring the level of kinase activity within said yeast cell, wherein said kinase activity typifies said protein.

20 11. The method according to claim 10, wherein step (B) comprises:

(A) labeling ATP with ³²P in vitro;

(B) running cellular proteins on a polyacrylamide gel; and

(C) determining the amount of ³²P labeled protein using autoradiography.

25 12. The method according to claim 5, wherein a promoter is operably linked to a reporter gene and wherein step (B) comprises monitoring the level of transcription of said reporter gene within said yeast cell.

30 13. The method according to claim 12, wherein said reporter gene is a luciferase gene and luciferase activity is monitored using a luminometer.

14. An isolated polynucleotide encoding the amino acid sequence of SEQ ID. NO. 4.

35 15. A culture of a bacterial strain containing the lambda phage λSG15.1.

* * * * *