



US007498151B2

(12) **United States Patent**  
**Schwartz et al.**

(10) **Patent No.:** **US 7,498,151 B2**  
(45) **Date of Patent:** **Mar. 3, 2009**

(54) **VARIANT TLR4 NUCLEIC ACID AND USES THEREOF**

(75) Inventors: **David A. Schwartz**, Iowa City, IA (US);  
**Brian C. Schutte**, Iowa City, IA (US)

(73) Assignee: **University of Iowa Research Foundation**, Iowa City, IA (US)

(\* ) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **11/517,172**

(22) Filed: **Sep. 7, 2006**

(65) **Prior Publication Data**

US 2007/0009992 A1 Jan. 11, 2007

**Related U.S. Application Data**

(63) Continuation of application No. 10/010,066, filed on Dec. 7, 2001, now Pat. No. 7,141,367, which is a continuation of application No. PCT/US00/15723, filed on Jun. 8, 2000, which is a continuation-in-part of application No. 09/329,515, filed on Jun. 10, 1999, now Pat. No. 6,740,487.

(51) **Int. Cl.**

**C12N 15/12** (2006.01)  
**C12N 15/64** (2006.01)  
**C12N 5/10** (2006.01)  
**C12P 21/02** (2006.01)  
**C07K 14/47** (2006.01)

(52) **U.S. Cl.** ..... **435/69.1**; 435/70.1; 435/320.1;  
435/325; 435/471; 536/23.5; 530/351

(58) **Field of Classification Search** ..... None  
See application file for complete search history.

(56) **References Cited**

**FOREIGN PATENT DOCUMENTS**

WO WO-98/50547 11/1998

**OTHER PUBLICATIONS**

Arbour, N. C., et al., "Missense mutations of the TLR4 gene are associated with hyporesponsiveness to lipopolysaccharides (LPS) in humans", *American Journal of Human Genetics*, 65(4), (Oct. 1999),A97.

Arbour, Nancy C., et al., "TLR4 mutations are associated with endotoxin hyporesponsiveness in humans", *Nature Genetics*, 25(2), (Jun. 2000), 187-191.

Bonner, J. C., et al., "Induction of the Lung Myofibroblast PDGF Receptor System by Urban Ambient Particles From Mexico City", *American Journal of Respiratory Cell and Molecular Biology*, 19(4), (Oct. 1998),672-680.

Brigham, K. L., et al., "Endotoxin and Lung Injury", *American Review of Respiratory Disease*, (May 1986),913-927.

Chow, J. C., et al., "Toll-like Receptor-4 Mediates Lipopolysaccharide-induced Signal Transduction", *Journal of Biological Chemistry*, 274(16), (Apr. 1999),10689-10692.

Favorite, G. O., et al., "Effects Produced by the Intravenous Injection in Man of a Toxic Antigenic Material Derived From Eberthella Typhosa: Clinical Hematological, Chemical and Serological Studies", *Journal of Clinical Investigation*, (May 1942),589-599.

Hoshino, K., et al., "Cutting Edge: Toll-like Receptor 4 (TLR4)—Deficient Mice are Hyporesponsive to Lipopolysaccharide: Evidence for TLR4 as the Lps Gene Product", *Journal of Immunology*, 162(7), (1999),3749-3752.

Kuhns, D. B., et al., "Endotoxin and IL-1 Hyporesponsiveness in a Patient With Recurrent Bacterial Infections", *Journal of Immunology*, 158(8), (Apr. 1997),3959-3964.

Medzhitov, R., et al., "A Human Homologue of the Drosophila Toll Protein Signals Activation of Adaptive Immunity", *Nature*, 388(6640), (1997),394-397.

Medzhitov, R., et al., "Innate Immunity: The Virtues of a Nonclonal System of Recognition", *Cell*, 91(3), (1997),295-298.

Michel, O., et al., "Effects of Inhaled Endotoxin on Bronchial Reactivity in Asthmatic and Normal Subjects", *Journal of Applied Physiology*, 66(3), (Mar. 1989),1059-1064.

Michel, O., et al., "Severity of Asthma is Related to Endotoxin in House Dust", *American Journal of Respiratory and Critical Care Medicine*, 154(6), (Dec. 1996),1641-1646.

Morris, B., et al., "A Novel Strategy for Directional Cloning of Random Primed cDNA", *InNovations Newsletter*, No. 3, Novagen, Inc.,(Jun. 1995),1-4.

Poltorak, A., et al., "Defective LPS Signaling in C3H/HeJ and C57BL/10ScCr Mice: Mutations in Tlr4 Gene", *Science*, 282, (1998),2085-2088.

Qureshi, S. T., et al., "Endotoxin-tolerant Mice Have Mutations in Toll-like Receptor 4 (tlr4)", *Journal of Experimental Medicine*, 189(4), (1999),615-625.

Rock, Fernando L., et al., "A family of human receptors structurally related to Drosophila Toll", *Proc. Natl. Acad. Sci. USA*, 95, (Jan. 1998),588-593.

Santamaria, P., et al., "Involvement of Class II MHC Molecules in the LPS-Induction of IL-1/TNF Secretions by Human Monocytes", *Journal of Immunology*, 143(3), (Aug. 1989),913-922.

Schwartz, D. A., et al., "Determinants of Longitudinal Changes in Spirometric Function Among Swine Confinement Operators and Farmers", *American Journal of Respiratory and Critical Care Medicine*, 151(1), (Jan. 1995),47-53.

Schwartz, D. A., et al., "The Role of Endotoxin in Grain Dust-induced Lung Disease", *American Journal of Respiratory and Critical Care Medicine*, 152(2), (Aug. 1995),603-608.

Wang, S., et al., "Effects of Inhalation of Steroids on Lung Permeability in Patients With Asthma", *Clinical Nuclear Medicine*, 20(6), (Jun. 1995),494-496.

Wright, S. D., et al., "CD14, a Receptor for Complexes of Lipopolysaccharide (LPS) and LPS Binding Protein", *Science*, 249(4975), (Sep. 1990),1431-1433.

Wright, S. D., "Toll, A New Prize in the Puzzle of Innate Immunity", *Journal of Experimental Medicine*, 189(4), (Feb. 1999),605-609.

Yang, R., et al., "Toll-like Receptor-2 Mediates Lipopolysaccharide-induced Cellular Signalling", *Nature*, 395(6699), (Sep. 1998),284-288.

*Primary Examiner*—Bridget E Bunner


*Assistant Examiner*—Fozia M Hamud

(74) *Attorney, Agent, or Firm*—Schwegman, Lundberg & Woessner, P.A.

(57) **ABSTRACT**

The invention provides methods to identify polymorphisms at the human TLR4 locus, as well as methods to identify individuals at risk of indications that increase their morbidity and mortality.

**18 Claims, 23 Drawing Sheets**



Intron	Position	5' to 3' sequence at EXON/intron boundary
1	135-136	GCGTGGAGgtatgtggctggagtcagct ---- attttgttttgtttttaaagACTTGGCC
2	255-256	TCACGGAGgttagaatgctgagcacgta ---- gtcatgtgtaatcattgcagGTGGTTCC
3	422-423	TTATCCAGgtaatgaatccacttttaca ---- atgtctttttattcctgtagGTGTGAAA

FIG. 1




FIG. 2A




FIG. 2B




FIG. 2C




FIG. 3A




FIG. 3B

↓

Human (aa 290)	.	.	.	L	A	Y	L	D	Y	Y	L	D	D	I	I	D	L	F	N	C	L	T	N	V	.	.	.
Mouse (aa 289)	.	.	.	L	T	Y	T	N	D	F	S	D	D	I	V	K	-	F	H	C	L	A	N	V	.	.	.
Rat (aa 289)	.	.	.	L	T	Y	I	N	H	F	S	D	D	I	Y	N	-	L	N	C	L	A	N	I	.	.	.
Hamster (aa 289)	.	.	.	F	T	Y	A	N	E	F	S	E	D	I	T	D	-	F	D	C	L	A	N	V	.	.	.

FIG. 4




FIG. 5




FIG. 6A






FIG. 6B




FIG. 6C




FIG. 6D

<u>Amino Acid</u>	<u>Codon</u>
Phe	UUU, UUC
Ser	UCU, UCC, UCA, UCG, AGU, AGC
Tyr	UAU, UAC
Cys	UGU, UGC
Leu	UUA, UUG, CUU, CUC, CUA, CUG
Trp	UGG
Pro	CCU, CCC, CCA, CCG
His	CAU, CAC
Arg	CGU, CGC, CGA, CGG, AGA, AGG
Gln	CAA, CAG
Ile	AUU, AUC, AUA
Thr	ACU, ACC, ACA, ACG
Asn	AAU, AAC
Lys	AAA, AAG
Met	AUG
Val	GUU, GUC, GUA, GUG
Ala	GCU, GCC, GCA, GCG
Asp	GAU, GAC
Gly	GGU, GGC, GGA, GGG
Glu	GAA, GAG

**FIG. 7**

Original Residue	Exemplary Substitutions	Preferred Substitutions
Ala (A)	val; leu; ile	val
Arg (R)	lys; gln; asn	lys
Asn (N)	gln; his; lys; arg	gln
Asp (D)	glu	glu
Cys (C)	ser	ser
Gln (Q)	asn	asn
Glu (E)	asp	asp
Gly (G)	pro	pro
His (H)	asn; gln; lys; arg	arg
Ile (I)	leu; val; met; ala; phe norleucine	leu
Leu (L)	norleucine; ile; val; met; ala; phe	ile
Lys (K)	arg; gln; asn	arg
Met (M)	leu; phe; ile	leu
Phe (F)	leu; val; ile; ala	leu
Pro (P)	gly	gly
Ser (S)	thr	thr
Thr (T)	ser	ser
Trp (W)	tyr	tyr
Tyr (Y)	trp; phe; thr; ser	phe
Val (V)	ile; leu; met; phe; ala; norleucine	leu

FIG. 8

## HUMAN TLR4 GENOMIC SEQUENCE

AAAATACTCC CTTGCCTCAA AAACCTGCTCG GTCAAACGGT  
 GATAGCAAAC CACGCATTCA CAGGGCCACT GCTGCTCACA  
 AAACCAGTGA GGATGATGCC AGGATGATGT CTGCCTCGCG  
 CCTGGCTGGG ACTCTGATCC CAGCCATGGC CTTCTCTCC  
 TGCCTGAGAC CAGAAAGCTG GGAGCCCTGC GTGGAGGTAT  
 GTGGCTGGAG TCAGCTCCTC TGAAC TTCC CTCACTTCTG  
 CCCAGA ACTT CTCACTGTGT GCCCTGGTTT GTTTATTTTT  
 GCAAAAAAAAA AAAGAGTTAA ATTACCTTAA AGACTCAAGA  
 AGCCACAGAG ATCAAATAAT TCATTGTTAC AGGGCACTAG  
 AGGCAGCCAT TGGGGGTTG TTCCATTGG AAATTTGAG  
 TGCTAACAGG GGCATGAGAT AACATAGATC TGCTTAAGGT  
 CCCTGCTCTG CTACCTTGTG GCTCTGTGAA GAAATTATCA  
 AACCTGTCTG AGACTAGTTT TCGCATCTGT AAGAGAATTA  
 TAATACCTTC TTCACTAGAG AGTAAGCAGA CTGCTTCAGT  
 GTCATTTCTT CCCACTGGTG GTCTTTACAC TCAGCTTCAA  
 GCAGTCACCC TGCTCCTTTC AATCTCAGGA AAAAGATGGC  
 TTTGTGTGTG TGTCTCT : A : G : AGAAAGAA CTTTCTAAGT  
 TGGTGCAGA CTTCTGTATG CAGTAATATA GTTTAGTCCA  
 GAGGATGAAA AAAATAAGAG A : ATGAAAAA GGAAAAGAGA  
 GAGAGAGA : G AAGAAAAAAG CAAGAGGGAA AT : ATGTATA  
 ATGTCAGCTA ATGCAAC : AG TTTCTTTCTT AGTGAATAC  
 CAATCAGCTG : GTTG : GTAA TCTT : ATTCA TGATGGATCT  
 CTTTTGTTTT TCCCCTGCGC AGACTTC : AC AGTTGCTTTA  
 GAAACCCATA GTAGAGCCGA A : CAGCTAAG AAAATGATTT  
 ACAGTGAGGC AGGGTCAGAA ACTCAAGAGA GAAAAAGCCA  
 GCTGCAGTC : CTGAAGT : TG AGGATATAGG : AGAAAATCA  
 AGTAATATTT AGCAAAGACT AATTCATTAT CTTGAAGCCA  
 TCCCTTCCCT CAATTCCCTG CCCATAGTCC TCCTCCTTGT  
 CCTCTTCTCT GNA : TCCCTC TGCTGTTAGG TTA : ATGG : A  
 GATAGATTTT CTAATTANGC TCACTGCGAG ATAAAACCCA  
 GCCCATGTTT CTATTAGNCA ATATTGTCTT TGAGGCTCCA  
 TGGCTTGCAN CATTTAAGCA GACATACGAA TGAAGATCTG  
 CATGTTTGAA CTCTGACTTT GCGCATATTA CTTCAATTTCT  
 TTGAATTTCC ATTTTCCTCA TCTTTAAATG CTTATTTGAA  
 GATTAAGTGA AAGTATATAA CAAACAAGAA CTATGCAGGC  
 GTATGGTAAG GGATTAATGA TAGATGATAA TAATTAATGT  
 TGACATCTAT TGATCACTTA TACTGTAGCG GGCTTTTAAA  
 TAAACTCTTT AAACACCTTA TCTCATTTAA TCCTTCAAAC  
 ATTCTATTGG TTTCAAACAA CAGAAAATA CAATTAGCTG  
 GCTTCTGCAA GGAATTTTGT TGGAGGAAAT GAGAGCATTC  
 AGAAATTAGA TGGGAGCGTT AGAGAATTAG GCTTACAAAG  
 AATGTGGGAA AGTAGGCTAG AAAGCAGTGT AAAAACAAAG  
 ACAGCATAAA GCACTTGACC TTATTTACTA GGTTCCACCA  
 TGGGAATCCA TGCACCTCTAA AGATTTCCCC CTATTTCTAC  
 ATCACTTTGC TCAAGGGTCA ATGAGCCAAG GAAAAGAATG

FIG. 9A

CAGTTGTCAA AATCTGGGCC ATGACTAAGG AAGGTCTGGA  
CATCTTGACT GCCAGACAGT CTCCCCAATG ATATGGAGTA  
TTTAGAATGA TACTGGATAT TTTATTTATT TTTTGTATTT  
TCAACTTTTA AGTTCAGAGG CACATGTGCA GAGCATGCAG  
GTTTATTACA TAAGTAAATG TGTGCCATGG TGATTTGCTG  
CATAGATCAT GAAAATATGG AACGCATCAT GGATTTGTGT  
GTCATCCTTG TGCAGGGGCC ATGCTCATCT TCTCTGTATC  
CTTCCAATTT TAGTATATGT GCTACTGCAG CAAGCACGAT  
ATTGGATATT TTATTACCTA CATTTTACAT ATGATAAAAT  
GAGGCTCACT GAGGTTTTTC TTTTGTTCGT TTTATTTTGT  
TTTGTTTTTA AAGACTTGSC CCTAAACCAC ACAGAAGAGC  
TGGCATGAAA CCCAGAGCTT TCAGACTCCG GAGCCTCAGC  
CCTTCACCCC GATTCCATTG CTTCCTTGCTA AATGCTGCCG  
TTTTATCNCG GAGGTTAGAA TGCTGAGCAC GTAGTAGGTG  
CTCTTTACTT TCTAATCTAG AGTAAGACAA TTTATAAGCA  
TGAATTGAGT GAATGGATGG ATGGATATAT GGATGGAAGG  
ATGGACAGAT GGATGAAAGG TTGACTGAAT TTTGTGCTTG  
CACAAAAGA GGCCCCTCTC CACCATCTCT GGTCTAGGAG  
AGGGGAGTTG GGAGACCATG CAGTAAAGAT ACTTCATGTC  
ATGTGTAATC ATTGCAGGTG GTTCCTAATA TTACTIONTCA  
ATGCATGGAG CTGAATTTCT ACAAATCCC CGACAACCTC  
CCCTTCTCAA CCAAGAACCT GGACCTGAGC TTTAATCCCC  
TGAGGCATTT AGGCAGCTAT AGCTTCTTCA GTTTCCAGA  
ACTGCAGGTG CTGGATTTAT CCAGGTAATG AATCCACTTT  
TACATACTGC ACAAGGTGAG GTGTTCATTG TCCTATCATT  
TCATTATGG ACTGGAAAGC TTGGTTTGTG GAGTCTCATC  
TTCATTCACT TATTCATTCA TACAACAGAT GTCTTATTAA  
CTATATAACC TTGAGCAAGC TACCTCTATT CTCCAGGTCT  
CAGTTTTCTA ATCTGTGAAG TAGGCAGTTG GCTGAGACAG  
CTTCTAAGGG CAATTCTAAT TTTAGGTTTT CTTTTAAGAC  
AGGAGAGAAA ATTAGCTTAA ATTCTTTCAT AAGCAGCTAT  
TTATTGACTA CTTGCTATAT GTTGTACACT CTGCAAGAAG  
ACAGGCATAT ATTGATATAT AACACACAGC CCCTGTTGTT  
AAGGAGGCAT ATCTTCTTGA AAGAGTTAAT ACCTTAAAGT  
CCTGGGTATG GTCCTGGGTA CATAGTATAT AGTCAACACA  
TTTTAATTAT GATTTTTTGG ATCTGGAAAC TGATATAAAG  
ATAGCGACAT ATAACAGTAG GTGATAAATT ATGTTTAAAC  
TAAAGGTAAC TAATTGTATT TTTCAGAAGA GGGCCTTCT  
CTGTGGTGGG TAGTCAAGAA AGATTCATGA ACTGCATAAG  
ATTCAAACAA TGCTTAGAAT ATTAAAATA GTGGTGGCAG  
GTGAAATGTC ATCTTGATAT TTTAGGGGAA CCAAATCTA  
AAAGGGTTTT CATCATCGGG GCCTTATTTG CAAATCGAAC  
TAGATAATGG ATCATGTTCT CTGCAATGGT TTGTAAAACA  
TTTCAAACA TTTTACATAT TTTTATTAT AGAAATTATT  
GATAAAGACT AAGGTCACAG TATAAAAATC CTTTTTAGAG  
CAGACATTTT TGTAGAAGAG TGAACATATG ACCTATTATA  
CTCTAATTTG GATATAGATA GGATGTAACA AAGGAGTAAT

FIG. 9A (Continued)

GGGAACAATT CAAAGGCAGT GGTATAGTGC ATANAGTCCT  
GTTGGGGTCA GAAGACCTGA GCCCAAGTTT ACCCCCAACA  
TTTATAACCC ATGTAACCTT AGCATATTAC TTCATCTCCC  
TTAATCCTTA GTTTCATATC TGATCAATGG AAATGATGAA  
ACTTATTCTG CTGGATTAAA TGTGATAATA AATATTAATA  
TGCTGTATAT ATTTAAATTT TTATAAAATA TATTTTATAA  
GCATAAAGTA TTCCTACAGA ATTTCATTAG GTTTTTAAAA  
TAATTTCAAC TTTTATTTTT GATTCAGGGA TTTACATGGT  
TATATTGCGT AATGCTGAGG TGTAGGTAC AATCGATACC  
ATCACTCAGG TAGTGAGCAT AGTACCCAAT AGTTAGTTTT  
TCAACCCTTG CTGCTTTCTC TCTATCCCCT CTCTAGTAAT  
CCCCAGGGTC TATTTTTGTC ATCTTTATGT CCATGTGTAC  
TCCATGTTTTG GATCCTACTT ATAAAGTGAG AACTCATGGT  
ATTTGGCTTT CTGTNCCTTT GTTNGCTAAT TTGCTTAGGA  
TAATGGCTAC TAGCTGCATC TATGCCATTA TGTTCTAAAT  
TTCANTTNC TGCATGAAAA TTTTGTCAAG TACTCTATTA  
AGGTAGACCA CCTCTCCCTT TTTTTTCAA ACAAGAAGTA  
GNTTTTCCCA AACAATGCCC TTATGGAATT NATCTTCAAT  
CCNNGGATAC CCAATAACTT GCCCAAANC CTTAATCTGN  
CTTACAGAGA GGCCACCTC CTTCTGTAAC CCATAGGAGA  
TTTGGATTGG TAAGAATGCT TTGTGATAGC CCAGCAGCCT  
TCTTTCCCT ATAGAAATAT ATATATANTC TTTTATAGG  
TGAGGAACTG AAGCTTGAAT AATTTAAATG ACTTATATAC  
ATNATCATTG CTTGTTAGCC ACAGACCAGA GATTTAAGTT  
CNCATCTCCA GAATCCAACT TAAATGTTTT CTTTGTCTTA  
ATACTCTACT TCTCTAAAGT GATTATCACC AATGTAATGA  
TATAGAGNCA CAGCAAGACC CTTTCCTTCT CACCTAATGT  
ATAGAGCAAT GCAGAGATAG AATGATGGGC TATAACAATC  
ATATAATTGA AAGAAAGAAC TTCAAAAATA ATCAAGTTCA  
GCTGTTTGAT TTATAAATGT GATAACTAAA ACCTAGAGAG  
GAAAAGAGGT ACTCAAGATC ACACAGTAGG AGAGGACTGC  
AGAAACACCA AACCCAAGCT CTTTTGTCCA CTCTTCCAGC  
GTTCTTTCTA CTATACTGCC TATCCTTTAT CTAGTTACCA  
ATAAATAACA AAAGCTTGGA CCACAATGCT TTTATTGTCT  
AGGAAACTCC TGAAGAAGCT AAATAAAATG GGTGGGGAAT  
ATTGTAAATG TAATTCAGGC TGGATTAAGA AAGAACTTAT  
TTGACATTGT AACTGACAAG CACCTGCAAT GCTGAAAGGA  
ATTTTTCAT TGGCNTGCTGT TTGCTGGGCT GCATCAAAGC  
CCTGTCTCTA GGACATGTCT CTGAACATTG TGTGTAGCAT  
GGCTTTCATT TCTTTTAGGA TAAAATTCAA AACCTTTTAT  
CTGGTTGGTA AACCTCTGCC TAATTGGGAA CCTTCTTTCT  
CCACAACCTC ATATTGTACA CTCCAATTC ATCTCTGTTC  
TCCAACCATG GAAGCTATTT GTCATGATTC CTCCTTGTGT  
CATTTTTTTT CTGTCAACCT TGGGGCTTTT GTGTTTGCTG  
TTCACTTCAC CTCCTTTTAT TGTTAACTTC TACTCATCTT  
TCAATTTTCA ACTTAAGTGT TCTCAGAGAA ACCTACTTTG  
ATTTTCTTGG TCCANAACGG TTCTCTGGAT GTGAACTCTT

FIG. 9A (Continued)



ATAGCACATA ATTTTCACTT TTTTCCACAA AACTCGCTCC  
 TATCACCTGT TACAAGCATT TACCTCTGAT AACAAGAACT  
 TTCAAATATC TAGCTGTCAT GTAAGCACTT TTCATAAACA  
 TTAAGAGTAT CTGTGACACT TATGTGTAAT GTTTCGTATC  
 TCTGAAATTG ATATTTACCA GTCATTTATC TTGGCTACCA  
 ACTAACAACF ATCCATATTA TCTGTACCAA TCAGATGTAT  
 AATCACAATT TTGTGTGACA GAAAATGGCT AACTTTGATC  
 CAAGGCTATT ACATGCTTT : ATCAACTGCA CAATCTTTAT  
 ATATGTCAAT TATTGATCTT TAACTGATTT CCTTCTTATG  
 : GATTTTCTC CTCTGCTTAT CATGTATGCC TAACAT : GAC  
 AAAAAAG : AG CCTA : TCATT GCAGCCAGTA TGATAAATACT  
 CA : GTCTGTG GGGCTTCTTA TTTGCTTAT : TCCATCATCA  
 TCTGTCCCTGC TTGATGTCTT TGCCTATGCA CAATCATATG  
 : ACCCATCAC ATCTGTATGA AGAGC : TGGA TGACTAGGAT  
 TAATATTCT : AT : : : TTTAG GTTCTTATT : CAGCAGAAAT  
 ATTAGATAA : TCAATGTCTT TTTATTCTTG TAGGTGTGAA  
 ATCCAGACAA TTGAAGATGG GGCATATCAG AGCCT : AAGC  
 CACCTCTCTA CCTTAATATT GACAGGAAAC CCCATCCAGA  
 GTTTAGCCCT GGGAGCCTTT TCTGGACTAT CAAGTTTACA  
 GAAGCTGGTG GCTGTGGAGA CAAATCTAGC ATCTCTAGAG  
 AACTTCCCCA TTGGACATCT CAAAACCTTG AAAGAACTTA  
 ATGTGGCTCA CAATCTTATC CAATCTTTCA AATTACCTGA  
 GTATTTTCT AATCTGACCA ATCTAGAGCA CTTGGACCTT  
 TCCAGCAACA AGATTCAAAG TATTTATTGC ACAGACTTGC  
 GGGTTCTACA TCAAATGCC CTACTCAATC TCTCTTTAGA  
 CCTGTCCCTG AACCCATATGA ACTTTATCCA ACCAGGTGCA  
 TTTAAAGAAA TTAGGCTTCA TAAGCTGACT TTAAGAAATA  
 ATTTTGATAG TTTAAATGTA ATGAAAACCT GTATTCAAGG  
 TCTGGCTGGT TTAGAAGTCC ATCGTTTGGT TCTGGGAGAA  
 TTTAGAAATG AAGGAACTT GGAAAAGTTT GACAAATCTG  
 CTCTAGAGGG CCTGTGCAAT TTGACCATTG AAGAATTCCC  
 GATTAGCATA CTTAGACTAC TACCTCGATG ATATTATTGA  
 CTTATTTAAT TGGTTGACAA ATGGTTCTTC ATTTTCCCTG  
 GTGAGTGTGA CTATTGAAAG GGTAAAAGAC TTTTCTTATA  
 ATTTCCGGATG GCAACATTTA GAATTAGTTA ACTGTAAATT  
 TGGACAGTTT CCCACATTGA AACTCAAATC TCTCAAAGG  
 CTTACTTTCA CTTCCAACAA AGGTGGGAAT GCTTTTTTCAG  
 AAGTTGATCT ACCAAGCCTT GAGTTTCTAG ATCTCAGTAG  
 AAATGGCTTG AGTTTCAAAG GTTGCTGTTC TCAAAGTGAT  
 TTTGGGACAA CCA : GCCT : A AAGTATTTAG ATCTGAGCTT  
 CAATGGTGTT A : TTACCATG AGTTCAAACF TCTTGGGCTT  
 AGAACA : ACT AGAACATCTG GATTTCCAGC ATTCCAATTT  
 GAAACA : AAT GAGTGAGTTT TCAGTATTCC TA : TCACTCA  
 GAAA : CCT : C ATTTACCTTG ACATTTCTCA TACTCACACC  
 AGAGTTGCTT TCAATGGCAT CTTCAATGGC TTGTCCAGTC  
 TCGAAGTCTT GAAAATGGCT GGCAATTCTT TCCAGGAAAA  
 CTTCCCTTCCA GATATCTTCA CAGAGCTGAG AACTTGACC

FIG. 9A (Continued)

TTCCTGGACC TCTCTCAGTG TCAACTGGAG CAGTTGTCTC  
CAACAGCATT TAACTCACTC TCCAGTCTTC AGGTACTAAA  
TATGAGCCAC AACAACTTCT TTTCATTGGA TACGTTTTCT  
TATAAGTGTC TGAACTCCCT CCAGGTCTTT GATTACAGTC  
TCAATCACAT AATGACTTCC AAAAAACAGG AACTACAGCA  
TTTTCCAAGT AGTCTAGCTT TCTTAAATCT TACTCAGAAT  
GACTTTGCTT GTACTTGTGA ACACCAGAGT TTCCTGCAAT  
GGATCAAGGA CCAGAGGCAG CTCTTGGTGG AAGTTGAACG  
AATGGAATGT GCAACACCTT CAGATAAGCA GGGCATGCCT  
GTGCTGAGTT TGAATATCAC CTGTCAGATG AATAAGACCA  
TCATTGGTGT GTCGGTCCTC AGTGTGCTTG TAGTATCTGT  
TGTAGCAGTT CTGGTCTATA AGTTCTATTT TCACCTGATG  
CTTCTTGCTG GCTGCATAAA GTATGGTAGA GGTGAAAACA  
TCTATGATGC CTTTGTTATC TACTCAAGCC AGGATGAGGA  
CTGGGTAAGG AATGAGCTAG TAAAGAATTT AGAAGAAGGG  
GTGCCTCCAT TTCAGCTCTG CCTTCACTAC AGAGACTTTA  
TTCCCGGTGT GGCCATTGCT GCCAACATCA TCCATGAAGG  
TTCCATAAA AGCCGAAAGG TGATTGTTGT GGTGTCCCAG  
CACTTCATCC AGAGCCGCTG GTGTATCTTT GAATATGAGA  
TTGCTCAGAC CTGGCAGTTT CTGAGCAGTC GTGCTGGTAT  
CATCTTCATT GTCCTGCAGA AGGTGGAGAA GACCCTGCTC  
AGGCAGCAGG TGGAGCTGFA CCGCCTTCTC AGCAGGAACA  
CTTACCTGGA GTGGGAGGAC AGTGTCCCTGG GGCGGCACAT  
CTTCTGGAGA CGACTCAGAA AAGCCCTGCT GGATGGTAAA  
TCATGGAATC CAGAAGGAAC AGTGGGTACA GGATGCAATT  
GGCAGGAAGC AACATCTATC TGAAGAGGAA AAATAAAAAC  
CTCCTGAGGC ATTTCTTGCC CAGCTGGGTC CAACACTTGT  
TCAGTTAATA AGTATTAAT GCTGCCACAT GTCAGGCCTT  
ATGCTAAGGG TGAGTAATTC CATGGTGCAC TAGATATGCA  
GGGCTGCTAA TCTCAAGGAG CTTCCAGTGC AGAGGGAATA  
AATGCTAGAC TAAAATACAG AGTCTTCCAG GTGGGCATTT  
CAACCAACTC AGTCAAGGAA CCCATGACAA AGAAAGTCAT  
TTCAACTCTT ACCTCATCAA GTTGAATAAA GACAGAGAAA  
ACAGAAAGAG ACATTGTTCT TTTCCTGAGT CTTTTGAATG  
GAAATTGTAT TATGTTATAG CCATCATAAA ACCATTTTGG  
TAGTTTTGAC TGAACTGGGT GTTCACTTTT TCCTTTTTGA  
TTGAATACAA TTTAAATCTT ACTTGATGAC TGCAGTCGTC  
AAGGGGCTCC TGATGCAAGA TGCCCCTTCC ATTTTAAGTC  
TGTCTCCTTA CAGAGGTAA AGTCTAGTGG CTAATTCCTA  
AGGAAACCTG ATTAACACAT GCTCACAACC ATCCTGGTCA  
TTCTCGAGCA TGTTCTATTT TTTAACTAAT CACCCCTGAT  
ATATTTTTAT TTTTATATAT CCAGTTTTCA TTTTTTTACG  
TCTTGCCTAT AAGCTAATAT CATAAATAAG GTTGTTTTAAG  
ACGTGCTTCA AATATCCATA TTAACCACTA TTTTCAAGG  
AAGTATGGAA AAGTACACTC TGTCACCTTG TCACTCGATG  
TCATTCAAA GTTATTGCCT ACTAAGTAAT GACTGTCATG  
AAAGCAGCAT TGAATAAATT TGTTTAAAGG GGCACCTCTT

FIG. 9A (Continued)

TTAAACGGGA	AGAAAATTTC	CGCTTCCTGG	TCTTATCATG
GACAATTTGG	GCTATAGGCA	TGAAGGAAGT	GGGATTACCT
CAGGAAGTCA	CCTTTTCTTG	ATTCCAGAAA	CATATGGGCT
GATAAACCCG	GGGTGACCTC	ATGAAATGAG	TTGCAGCAGA
TGTTTATTTT	TTTCAGAACA	AGTGATGTTT	GATGGACCTA
TGAATCTATT	TAGGGAGACA	CAGATGGCTG	GGATCCCTCC
CCTGTACCCT	TCTCACTGCC	AGGAGAACTA	CGTGTGAAGG
TATTCAAGGC	AGGGAGTATA	CATTGCTGTT	TCCTGTTGGG
CAATGCTCCT	TGACCACATT	TTGGGAAGAG	TGGATGTTAT
CATTGAGAAA	ACAATGTGTC	TGGAATTAAT	GGGGTTCTTA
TAAAGAAGGT	TCCCAGAAAA	GAATGTTTCA	TCCAGCTTCT
TCAGGAAACA	GGAACATTCA	AGGAAAAGGA	CAATCAGGAT
GTCATCAGGG	AAATGAAAAT	AAAAACCACA	ATGAGATATC
ACCTTATACC	AGGTAGATGG	CTACTATAAA	AAAATGAAGT
GTCATCAAGG	ATATAGAGAA	ATTGGAACCC	TTCTTCACTG
CTGGAGGGAA	TGGAAAATGG	TGTAGCCGTT	ATGAAAACA
GTACGGAGGT	TTCTCAAAAA	TTAAAAATAG	AACTGCTATA
TGATCCAGCA	ATCTCACTTC	TGTATATATA	CCCAAATAA
TTGAAATCAG	AATTTCAAGA	AAATATTTAC	ACTCCCATGT
TCATTGTGGC	ACTCTTCACA	ATCACTGTTT	CCAAAGTTAT
GGAAACAACC	CAAATTTCCA	TTGGAAAATA	AATGGACAAA
GGAAATGTGC	ATATAACGTA	CAATGGGGAT	ATTATTCAGC
CTAAAAAAG	GGGGGATCCT	GTTATTTATG	ACAACATGAA
TAAACCCGGA	GGCCATTATG	CTATGTAAAA	TGAGCAAGTA
ACAGAAAGAC	AAATACTGCC	TGATTTTATT	TATATGAGGT
TCTAAAATAG	TCAAACATCAT	AGAAGCAGAG	AATAGAACAG
TGGTTCCTAG	GGAAAAGGAG	GAAGGGAGAA	ATGAGGAAAT
AGGGAGTTGT	CTAATTGGTA	TAAAATTATA	GTATGCAAGA
TGAATTAGCT	CTAAAGATCA	GCTGTATAGC	AGAGTTCGTA
TAATGAACAA	TACTGTATTA	TGCACTTAAC	ATTTTGTTAA
GAGGGTACCCT	CTCATGTTAA	GTGTTCTTAC	CATATACATA
TACACAAGGA	AGCTTTTGGA	GGTGATGGAT	ATATTTATTA
CCTTGATTGT	GGTGATGGTT	TGACAGGTAT	GTGACTATGT
CTAAACTCAT	CAAATTGTAT	ACATTAATA	TATGCAGTTT
TATAATATCA	AAAAAAAAAA	AAAAAAAAAA	

FIG. 9A (Continued)

1 TTCCACTTCT AAGAGCTGCC TAGAGTAGTC AAGATTATAG AGACAAAAGT  
51 AGTGCATAGA TTCAAGGGCC TAGGGAAAGG GGAAATGGGG AGTTATTTAT  
101 TAATGAATAG TGGTGATGAT TGTACAAAAA TATGAACATA ATTAATGCCA  
151 CTAAATTGTN CACATACAAA TGGTCAAGAT AATAAATTTT ATGTTATGTC  
201 ATGTTATGTT ATGTGATTTT ACCATAATAC AGAAAATGAA AAAAGAAAAG  
251 AAAGAAAGTA AAGCTTAGCG GTTNCATGA CTTGNCCAAT GCCTCAAAGC  
301 CATGAGTCGA CCCAGCTGAG ATCTGANCTT CAGTATATTC CATTCTGAAA  
351 TCCCAGACTT TTCCAATCT TCTTGTA CTT TCAAACTGT GTTTCAGTTG  
401 AGGTTTATTT TCAGTTTGT ATGTGAGPTT CTTGCAAGA AGGGCGGGCC  
451 AAATTGTGTC CTGCAAAAAC CTACATATCG AAGTCCTAAC CCCTCTACCT  
501 CAGACTATGA CTGTATATGG AGAGAGAGCC TTGAAAGAGG TATGTAAGGT  
551 AGAATGAGGT CATTATGGTG GGCCCTAATC CAACATAACT GGTGTCCTTA  
601 TAAGAAGGGG AGATTAGAAT TCAGACACAC TTGCTGACAC CTTGAGTTCA  
651 GACTGGAAGC CTCTAGAATT GTGAGAAAAT GAATGTCTGT TGTTTAAGCC  
701 ACCCAGTCTG TGGTATTTCC TTATGGCAGC CCCAGCAAAC TAATACAAAT  
751 AGTGTTTCCA CAGCTGAAAC AAAATGGAA AATCACCGTC ATCCTAGAGA  
801 GTTACAAGGG CTATTTTAAT AGAACCTGAT TGTTTTCTTA AATTCACCAA  
851 GCCCAGGCAG AGGTCAGATG ACTAATTGGG ATAAAAGCCA ACTAGCTTCC  
901 TCTTGCTGTT TCTTTAGCCA CTGGTCTGCA GGCGTTTTCT TCTTCTAACT  
951 TCCTCTCCTG TGACAAAAGA GATAACTATT AGAGAAACAA AAGTCCAGAA  
1001 TGCTAAGGTT GCCGCTTCA CTTCTCTCA CCCTTTAGCC CAGAAGTCT  
1051 TTGAATACAC CAATTGCTGT GGGGCGGCTC GAGGAAGAGA AGACACCAGT  
1101 GCCTCAGAAA CTGCTCGGTC AGACGGTGAT AGCGAGCCAC GCATTCACAG  
1151 GGCCACTGCT GCTCACAGAA GCAGTGAGGA TGATGCCAGG ATGATGTCTG  
1201 CCTCGCGCCT GGCTGGGACT CTGATCCCAG CCATGGCCTT CCTCTCCTGC  
1251 GTGAGACCAG AAAGCTGGGA GCCCTGCGTG GAGGTATGTG GCTGGAGTCA  
1301 GCTCCTCTGA ACTTCCCTC ACTTCTGCCC AGAACTTCTC ACTGTGTGCC  
1351 CTGGTTTGTT

FIG. 9B

1 CGCATCATGG ATTTGTGTGT CATCCTTGTG CAGGGGCCAT GCTCATCTTC  
51 TCTGTATCCT TCCAATTTTA GTATATGTGC TACTGCAGCA AGCACGATAT  
101 TGGATATTTT ATTACCTACA TTTTACATAT GATAAAATGA GGCTCACTGA  
151 GGTTTTTCTT TTGTTTCGTTT TATTTTGTTT TGTTTTTAAA GACTTGGCCC  
201 TAAACCACAC AGAAGAGCTG GCATGAAACC CAGAGCTTTC AGACTCCGGA  
251 GCCTCAGCCC TTCACCCCGA TTCCATTGCT TCTTGCTAAA TGCTGCCGTT  
301 TTATCACGGA GGTTAGAATG CTGAGCACGT AGTAGGTGCT CTTTACTTTC  
351 TAATCTAGAG TAAGACAATT TATAAGCATG AATTGAGTGA ATGGATGGAT  
401 GGATATATGG ATGGAAGGAT GGACAGATGG ATGAAAGGTT GACTGAATTT  
451 TGTGCTTGCA CAAAAGAGG CCCCTCTCCA CCATCTCTGG TCTAGGAGAG  
501 GGGAGTTGGG AGACCATGCA GTAAAGATAC TTCATGTCAT GTGTAATCAT  
551 TGCAGGTGGT TCCTAATATT ACTTATCAAT GCATGGAGCT GAATTTCTAC  
601 AAAATCCCCG ACAACCTCCC CTTCTCAACC AAGAACCTGG ACCTGAGCTT  
651 TAATCCCCTG AGGCATTTAG GCAGCTATAG CTTCTTCAGT TTCCCAGAAC  
701 TGCAGGTGCT GGATTTATCC AGGTAATGAA TCCACTTTTA CATACTGCAC  
751 AAGGTGAGGT GTTCATTGTC CTATCATTTT ATTATTGGAC TGGAAAGCTT  
801 GGTTTGTGGA GTCTCATCTT CATTCACTTA TTCATTCATA CAACAGATGT  
851 CTTATTA ACT ATATAACCTT GAGCAAGCTA CCTCTATTCT CCAGGTCTCA  
901 GTTTTCTAAT CTGTGAAGTA GGCAGTTGGC TGAGACAGCT TCTAAGGGCA  
951 ATTCTAATTT TAGGTTTTCT TTTAAGACAG GAGAGAAAAT TAGCTTAAAT  
1001 TCTTTCATAA GCAGCTATTT ATTGACTACT TGCTATATGT TGTACTCTCT  
1051 GCAAGAAGAC AGGCATATAT TGATATATAA CACACAGCCC CTGTTGTTAA  
1101 GGAGGCATAT CTTCTTGAAA GAGTTAATAC CTTAAAGTCC TGGGTATGGT  
1151 CCTGGGTACA TAGTATATAG TCAACACATT TTAATTATGA TTTTTTGGAT  
1201 CTGGAAACTG ATATAAAGAT AGCGACATAT AACAGTAGGT GATAAATTAT  
1251 GTTTAAACTA AAGGTA ACTA ATTGTATTTT TCAGAAGAGG GGCCTTCTCT  
1301 GTGGTGGGTA GTCAAGAAAG ATTCATGAAC TGC

FIG. 9C

```

1   GGTAAGAATG CTTTGTGATA GCCCAGCAGC CTTCTTTCCC CTATAGAAAT
51  ATATATATAN TCTTTTTATA GGTGAGGAAC TGAAGCTTGA ATAATTTAAA
101 TGACTTATAT ACATNATCAT TGCTTGTTAG CCACAGACCA GAGATTTAAG
151 TTCNCATCTC CAGAATCCAA CTAAATGTT TTCTTTGTCT TAATACTCTA
201 CTTCTCTAAA GTGATTATCA CCAATGTAAT GATATAGAGN CACAGCAAGA
251 CCCTTTCCTT CTCACCTAAT GTATAGAGCA ATGCAGAGAT AGAATGATGG
301 GCTATAACAA TCATATAAAT GAAAGAAAGA ACTTCAAAAA TAATCAAGTT
351 CAGCTGTTTG ATTTATAAAT GTGATAACTA AAACCTAGAG AGGAAAAGAG
401 GTECTAAGA TCACACAGTA GGAGAGGACT GCAGAAACAC CAAACCCAAG
451 CTCTTTTGTC CACTCTTCCA GCGTCTTTC TACTATACTG CCTATCCTTT
501 ATCTAGTTAC CAATAAATAA CAAAAGCTTG GACCACAATG CTTTTATTGT
551 CTAGGAAACT CCTGAAGAAG CTAAATAAAA TGGGTGGGGA ATATTGTAAA
601 TGTAATTCAG GCTGGATTAA GAAAGAACTT ATTTGACATT GTAACGACA
651 AGCACCTGCA ATGCTGAAAG GAATTTTTCA TTGGCNTGCT GTTTGCTGGG
701 CTGCATCAA GCCCTGTCTC TAGGACATGT CTCTGAACAT TGTGTGTAGC
751 ATGGCTTTC TTTCTTTTAG GATAAAATTC AAAACCTTT ATCTGGTTGG
801 TAAACCTCTG CCTAATTGGG AACCTTCTTT CTCCACAAC CCATATTGTA
851 CACTCCAATT TCATCTCTGT TCTCCAACCA TGGAAAGCTAT TTGTCATGAT
901 TCCTCCTTGT GTCATTTTTT TTCTGTCAAC CTTGGGGCTT TTGTGTTTGC
951 TGTFCACTTC ACCTCCTTTT ATTGTTAACT TCTACTCATC TTTCAATTTT
1001 CAACTTAAGT GTTCTCAGAG AAACCTACTT TGATTTTCTT GGTCCANAAC
1051 GGTTCTCTGG ATGTGAACTC TTATAGCACA TAATTTTCAC TTTTTTCCAC
1101 AAAACTCGCT CCTATCACCT GTTACAAGCA TTIACCTCTG ATAACAAGAA
1151 CTTTCAAATA TCTAGCTGTC ATGTAAGCAC TTTTCATAAA CATTAAAGAGT
1201 ATCTGTGACA CTTATGTGTA ATGTTTCGTA TCTCTGAAAT TGATATTTAC
1251 CAGTCATTTA TCTTGGCTAC CAACTAACAA CTATCCATAT TATCTGTACC
1301 AATCAGATGT ATAATCACAA TTTTGTGTGA CAGAAAATGG CTAAACTTGA
1351 TCCAAGGCTA TTACATGCTT TATCAACTGC ACAATCTTTA TATATGTCAA
1401 TTATTGATCT TTANCTGATT TCCTTCTTAT GGATTTTCTC CTCTGCTTAT
1451 CATGTATGCC TAACATGACA AAAAAGAGCC TATCATTGCA GCCAGTATGA
1501 TAATACTCAG TCTGTGGGGC TTCTTATTTG CTTATTCCAT CATCATCTGT
1551 CCTGCTTGAT GTCTTTGCCT ATGCACAATC ATATGACCCA TCACATCTGT
1601 ATGAAGAGCT GGATGACTAG GATTAATATT CTATTTTAGG TTCTTATTCA
1651 GCAGAAATAT TAGATAATCA ATGTCTTTTT ATTCCTGTAG GTGTGAAATC
1701 CAGACAATTG AAGATGGGGC ATATCAGAGC CTAAGCCACC TCTCTACCTT
1751 AATATTGACA GGAAACCCCA TCCAGAGTTT AGCCCTGGGA GCCTTTTCTG
1801 GACTATCAAG TTTACAGAAG CTGGTGGCTG TGGAGACAAA TCTAGCATCT
1851 CTAGAGAACT TCCCATTGG ACATCTCAA ACTTTGAAAG AACTTAATGT
1901 GGCTCACAAT CTTATCCAAT CTTTCAAAT ACCTGAGTAT TTTTCTAATC
1951 TGACCAATCT AGAGCACTTG GACTTTCCA GCAACAAGAT TCAAAGTATT
2001 TATTGCACAG ACTTGCGGGT TCTACATCAA ATGCCCTAC TCAATCTCTC
2051 TTTAGACCTG TCCCTGAACC CTATGAACTT TATCCAACCA GGTGCATTTA
2101 AAGAAATTAG GCTTCATAAG CTGACTTTAA GAAATAATTT TGATAGTTTA
2151 AATGTAATGA AAACCTGTAT TCAAGGTCTG GCTGGTTTAG AAGTCCATCG
2201 TTTGGTTCTG GGAGAATTTA GAAATGAAGG AAACCTGGAA AAGTTTGACA

```

FIG. 9D

2251 AATCTGCTCT AGAGGGCCTG TGCAATTTGA CCATTGAAGA ATTCCGATTA  
2301 GCATACTTAG ACTACTACCT CGATGATATT ATTGACTTAT TTAATTGTTT  
2351 GACAAATGTT TCTTCATTTT CCCTGGTGAG TGTGACTATT GAAAGGGTAA  
2401 AAGACTTTTC TTATAATTTT GGATGGCAAC ATTTAGAATT AGTTAACTGT  
2451 AAATTTGGAC AGTTTCCCAC ATTGAAACTC AAATCTCTCA AAAGGCTTAC  
2501 TTTCACTTCC AACAAAGGTG GGAATGCTTT TTCAGAAGTT GATCTACCAA  
2551 GCCTTGAGTT TCTAGATCTC AGTAGAAATG GCTTGAGTTT CAAAGGTTGC  
2601 TGTTCTCAA GTGATTTTGG GACAACCAGC CTAAAGTATT TAGATCTGAG  
2651 CTTCAATGGT GTTATTACCA TGAGTTCAA CTTCTTGGGC TTAGAACAAC  
2701 TAGAACATCT GGATTTCCAG CATTCCAATT TGAAACAAAT GAGTGAGTTT  
2751 TCAGTATTCC TATCACTCAG AAACCTCATT TACCTTGACA TTTCTCATA  
2801 TCACACCAGA GTTGCTTTCA ATGGCATCTT CAATGGCTTG TCCAGTCTCG  
2851 AAGTCTTGAA AATGGCTGGC AATTCTTTCC AGGAAAACCT CCTTCCAGAT  
2901 ATCTTCACAG AGCTGAGAAA CTGACCTTC CTGGACCTCT CTCAGTGTCA  
2951 ACTGGAGCAG TTGTCTCCAA CAGCATTTAA CTCACTCTCC AGTCTTCAGG  
3001 TACTAAATAT GAGCCACAAC AACTTCTTTT CATTGGATAC GTTTCCTTAT  
3051 AAGTGTCTGA ACTCCCTCCA GGTTCCTGAT TACAGTCTCA ATCACATAAT  
3101 GACTTCCAAA AACAGGAAC TACAGCATT TCCAAGTAGT CTAGCTTTCT  
3151 TAAATCTTAC TCAGAATGAC TTTGCTTGTA CTTGTGAACA CCAGAGTTTC  
3201 CTGCAATGGA TCAAGGACCA GAGGCAGCTC TTGGTGGAAG TTGAACGAAT  
3251 GGAATGTGCA ACACCTTCAG ATAAGCAGGG CATGCCTGTG CTGAGTTTGA  
3301 ATATCACCTG TCAGATGAAT AAGACCATCA TTGGTGTGTC GGTCCTCAGT  
3351 GTGCTTGTAG TATCTGTTGT AGCAGTTCCTG GTCTATAAGT TCTATTTTCA  
3401 CCTGATGCTT CTGCTGGCT GCATAAAGTA TGGTAGAGGT GAAAACATCT  
3451 ATGATGCCTT TGTATCTAC TCAAGCCAGG ATGAGGACTG GGTAAGGAAT  
3501 GAGCTAGTAA AGAATTTAGA AGAAGGGGTG CCTCCATTC AGCTCTGCCT  
3551 TCACTACAGA GACTTTATTC CCGGTGTGGC CATTGCTGCC AACATCATCC  
3601 ATGAAGGTTT CCATAAAAGC CGAAAGGTGA TTGTTGTGGT GTCCCAGCAC  
3651 TTCATCCAGA GCCGCTGGT TATCTTTGAA TATGAGATTG CTCAGACCTG  
3701 GCAGTTTCTG AGCAGTCTGT CTGGTATCAT CTTTATTGTC CTGCAGAAGG  
3751 TGGAGAAGAC CCTGCTCAGG CAGCAGGTGG AGCTGTACCG CCTTCTCAGC  
3801 AGGAACACTT ACCTGGAGTG GGAGGACAGT GTCCTGGGGC GGCACATCTT  
3851 CTGGAGACGA CTCAGAAAAG CCCTGCTGGA TGGTAAATCA TGGAAATCCAG  
3901 AAGGAACAGT GGGTACAGGA TGCAATTTGGC AGGAAGCAAC ATCTATCTGA  
3951 AGAGGAAAAA TAAAAACCTC CTGAGGCATT TCTTGCCCAG CTGGGTCCAA  
4001 CACTTGTTC A GTTAATAAGT ATTAATGCT GCCACATGTC AGGCCTTATG  
4051 CTAAGGGTGA GTAATTCCAT GGTGCACTAG ATATGCAGGG CTGCTAATCT  
4101 CAAGGAGCTT CCAGTGCAGA GGAATAAAT GCTAGACTAA AATACAGAGT  
4151 CTTCCAGGTG GGCATTTCAA CCAACTCAGT CAAGGAACCC ATGACAAAGA  
4201 AAGTCATTC AACTCTTACC TCATCAAGTT GAATAAAGAC AGAGAAAACA  
4251 GAAAGAGACA TTGTTCTTTT CCTGAGTCTT TTGAATGGAA ATTGTATTAT  
4301 GTTATAGCCA TCATAAAACC ATTTTGGTAG TTTTGACTGA ACTGGGTGTT  
4351 CACTTTTCC TTTTGTATTG AATACAATTT AAATTCTACT TGATGACTGC  
4401 AGTCGTCAAG GGGCTCCTGA TGCAAGATGC CCCTTCCATT TTAAGTCTGT  
4451 CTCCTTACAG AGGTTAAAGT CTAGTGGCTA ATTCCTAAGG AAACCTGATT  
4501 AACACATGCT CACAACCATC CTGGTCATTC TCGAGCATGT TCTATTTTTT  
4551 AACTAATCAC CCCTGATATA TTTTTATTTT TATATATCCA GTTTTCATTT

FIG. 9D (Continued)

4601 TTTTACGTCT TGCCTATAAG CTAATATCAT AAATAAGGTT GTTTAAGACG  
 4651 TGCTTCAAAT ATCCATATTA ACCACTATTT TTCAAGGAAG TATGGAAAAG  
 4701 TACACTCTGT CACTTTGTCA CTCGATGTCA TTCCAAAGTT AFTGCCTACT  
 4751 AAGTAATGAC TGTTCATGAAA GCAGCATTGA AATAATTTGT TTAAAGGGGG  
 4801 CACTCTTTTA AACGGGAAGA AAATTTCCGC TTCCTGGTCT TATCATGGAC  
 4851 AATTTGGGCT AGAGGCAGGA AGGAAGTGGG ATGACCTCAG GAGGTCACCT  
 4901 TTTCTTGATT CCAGAAACAT ATGGGCTGAT AAACCCGGGG TGACCTCATG  
 4951 AAATGAGTTG CAGCAGAAGT TTATTTTTTT CAGAACAAGT GATGTTTGAT  
 5001 GGACCTCTGA ATCTCTTTAG GGAGACACAG ATGGCTGGGA TCCCTCCCT  
 5051 GTACCCTTCT CACTGCCAGG AGAACTACGT GTGAAGGTAT TCAAGGCAGG  
 5101 GAGTATACAT TGCTGTTTCC TGTTGGGCAA TGCTCCTTGA CCACATTTTG  
 5151 GGAAGAGTGG ATGTTATCAT TGAGAAAACA ATGTGTCTGG AATTAATGGG  
 5201 GTTCTTATAA AGAAGGTTCC CAGAAAAGAA TGTTCATCCA GCCTCCTCAG  
 5251 AAACAGAACA TTCAAGAAAA GGACAATCAG GATGTCATCA GGGAAATGAA  
 5301 AATAAAAACC ACAATGAGAT ATCACCTTAT ACCAGGTAGA ATGGCTACTA  
 5351 TAAAAAATG AAGTGTTCATC AAGGATATAG AGAAATTGGA ACCCTTCTTC  
 5401 ACTGCTGGAG GGAATGGAAA ATGGTGTAGC CGTTATGAAA AACAGTACGG  
 5451 AGGTTTCTCA AAAATTAAAA ATAGAACTGC TATATGATCC AGCAATCTCA  
 5501 CTTCTGTATA TATACCCAAA ATAATTGAAA TCAGAATTTT AAGAAAATAT  
 5551 TTACACTCCC ATGTTTCATTG TGGCACTCTT CACAATCACT GTTCCAAAG  
 5601 TTATGGAAAC AACCCAAATT TCCATTGAAA AATAAATGGA CAAAGAAAAT  
 5651 GTGCATATAC GTACAATGGG ATATTATTCA GCCTAAAAAA AGGGGGNATC  
 5701 CTGTTATTTA TGACAACATG AATAAACCCG GAGCCATTAT GCTATGTAAA  
 5751 ATGAGCAAGT AACAGAAAGA CAAATACTGC CTGATTTTAT TTATATGAGG  
 5801 TTCTAAAATA GTCAAACTCA TAGAAGCAGA GAATAGAACA GTGGTTCCTA  
 5851 GGGAAAAGGA GGAAGGGAGA AATGAGGAAA TAGGGAGTTG TCTAATTGGT  
 5901 ATAAAATTAT AGTATGCAAG ATGAATTAGC TCTAAAGATC AGCTGTATAG  
 5951 CAGAGTTCGT ATAATGAACA ATACTGTATT ATGCACTTAA CATTTTGTTA  
 6001 AGAGGGTACC TCTCATGTTA AGTGTTCCTA CCATATACAT ATACACAAGG  
 6051 AAGCTTTTGG AGGTGATGGA TATATTTATT ACCTTGATTG TGGTGTGGT  
 6101 TTGACAGGTA TGTGACTATG TCTAAACTCA TCAAATTGTA TACATTAAT  
 6151 ATATGCAGTT TTATAATATC AATTATGTCT GAATGAAGCT ATAAAAAAGA  
 6201 AAAGACAACA AAATTCAGTT GTCAAAACTG GAAATATGAC CACAGTCAGA  
 6251 AGTGTTTGTT ACTGAGTGT TCCAGAGTGT TTTGGTTTGA GCAGGTCTAG  
 6301 GGTGATTGAA CATCCCTGGG TGTGTTTCCA TGTCTCATGT ACTAGTGAAA  
 6351 GTAGATGTGT GCATTTGTGC ACATATCCCT ATGTATCCCT ATCAGGGCTG  
 6401 TGTGTATTTG AAAGTGTGTG TGTCCGCATG ATCATATCTG TATAGAAGAG  
 6451 AGTGTGATTA TATTTCTTGA AGAATACATC CATTTGAAAT GGATGTCTAT  
 6501 GGCTGTTTGA GATGAGTTCT CTA CTCTTGT GCTTGACAG TAGTCTCCCC  
 6551 TTATCCCTTA TGCTTGGTGG ATACGTTCTT AGACCCCAAG TGGATCTCTG  
 6601 AGACCGCAGA TGGTACCAA CCTCATATAT GCAATATTTT TTCCTATACA  
 6651 TAAATACCTA AGATAAAGTT CATCTTCTGA ATTAGGCACA GTAAGAGATT  
 6701 AACAAATACT AACAAATAAA TTGAATAGTT ATAATAATAT ATTGTAATAA  
 6751 AAGTTATGTG AATGTGATCT CTTTCTTTTC TCTCTC

FIG. 9D (Continued)



## VARIANT TLR4 NUCLEIC ACID AND USES THEREOF

### RELATED APPLICATIONS

This application is a continuation under 37 C.F.R. 1.53(b) of U.S. application Ser. No. 10/010,066 filed Dec. 7, 2001 now U.S. Pat. No. 7,141,367, which is a continuation under 35 USC 111(a) of International Application Serial No. PCT/US00/15723 filed on Jun. 8, 2000 and published in English on Dec. 21, 2000 as WO 00/77204 A1, which is a continuation-in-part of U.S. application Ser. No. 09/329,515, filed Jun. 10, 1999 (U.S. Pat. No. 6,740,487); which applications and publication are incorporated herein by reference.

### STATEMENT OF GOVERNMENT RIGHTS

This invention was made at least in part with a grant from the Government of the United States of America (grants ES06537, ES07498 and ES05605 from the National Institute of Environmental Sciences, grant HL62628 from the National Heart and Lung Institute, and grant RR00059 from the General Clinical Research Centers Program). The Government may have certain rights in the invention.

### BACKGROUND OF THE INVENTION

Endotoxin or lipopolysaccharide (LPS), released from the cell wall of bacteria, plays a central role in a broad spectrum of human disease. The pathogenic importance of LPS in gram-negative sepsis is well established. Intravenous LPS induces all of the clinical features of gram-negative sepsis, including fever, shock, leukopenia followed by leukocytosis, and disseminated intravascular coagulation (Favorite et al., 1942). Higher concentrations of circulating levels of endotoxin have been associated with manifestations of systemic inflammatory response syndrome (Wang et al., 1995) and the development of acute respiratory distress syndrome following sepsis (Brigham et al., 1986). Inhaled endotoxin can induce airflow obstruction in naive or previously unexposed subjects (Michel et al., 1992) and is the most important occupational exposure associated with the development Schwartz et al., 1995a), and progression (Schwartz et al., 1995b), of airway disease among exposed workers. The concentration of endotoxin in the domestic setting appears to be associated with the clinical severity of asthma (Michel et al., 1996). Moreover, recent studies have shown that endotoxin is a contaminant of particulate matter in air pollution and may play a role in the pathophysiologic consequences of air pollution (Bonner et al., 1998). Thus, endotoxin is an important cause of morbidity and mortality.

The ability of the host to respond to endotoxin may play an important role in determining the severity of the physiologic and biologic response to this frequently encountered toxin. In mice, genetic differences in susceptibility to LPS have been established. LPS hyporesponsiveness arose spontaneously and was first identified in the C3H/HeJ strain. This strain had an LD<sub>50</sub> for LPS at least 20 times that observed in A/HeJ mice (Sultzter et al., 1968). In addition to C3H/HeJ, two other mouse strains, C57BL/10ScCR (Coutinho et al., 1978) and its progenitor strain, C57BL/10ScN, (Vogel et al., 1979), are hyporesponsive to LPS.

Moreover, several reports suggest that humans may also respond differently to LPS. A patient with recurrent bacterial infections has been reported to be refractory to the in vivo and in vitro effects of LPS (Kuhns et al., 1997). Following challenge with intravenous LPS, this patient had no systemic

increase in IL-6 or G-CSF and had a minimal rise in the concentration of leukocytes, TNF- $\alpha$ , and IL-8. Inter-individual differences have also been reported in the release and synthesis of cytokines by monocytes stimulated with LPS in vitro (Santamaria et al., 1989).

LPS is thought to cause much of its morbidity and mortality by activating kinases (DeFranco et al., 1998) that control the function of transcription factors (nuclear factor- $\kappa$ B and AP-1) and ultimately lead to production of pro-inflammatory cytokines and co-stimulatory molecules (Wright, 1999). Several lines of evidence suggest that the toll receptor (TLR) family, and specifically TLR4 and TLR2 regulate the interaction between LPS and intracellular kinases and may serve as a proximal target to interrupt LPS signaling (Wright, 1998; Medzhitov et al., 1997). Both TLR4 and TLR2 activate signaling through NF- $\kappa$ B and AP-1 in transfected human cell lines (Medzhitov et al., 1997; Yang et al., 1998), and TLR4 mediates LPS induced signal transduction (Chow et al., 1999). CD14, a glycosylphosphatidyl inositol-linked receptor that binds LPS (Poltorak et al., 1998a) enhances LPS induced TLR2 (Yang et al., 1998) and TLR4 (Chow et al., 1998) signaling, suggesting that the toll receptors interact with CD14 to initiate the cellular response to LPS. Studies in mice indicate that 1) the TLR4 gene maps to the critical region in LPS hyporesponsive mice (Poltorak et al., 1998), 2) mutations in the TLR4 gene (Poltorak et al., 1998; Qureshi et al., 1999) are found in mouse strains (C3H/HeJ and C57BL10/ScCr) that are defective in their response to LPS, and 3) disruption of the TLR4 gene results in a LPS hyporesponsive phenotype (Hoshino et al., 1999).

Thus, there is need to determine whether the human TLR4 gene is polymorphic, and whether any particular polymorphism is associated with disease, e.g., LPS hyporesponsiveness.

### SUMMARY OF THE INVENTION

The invention provides a method to identify a mammal, e.g., a human, at risk of, or having, an indication associated with altered innate immunity, e.g., to bacterial infection. The method comprises contacting an amount of DNA obtained from a human physiological sample with an amount of at least one TLR4-specific oligonucleotide under conditions effective to amplify the DNA so as to yield amplified DNA. Then it is determined whether the amplified DNA comprises a nucleotide substitution, e.g., one that results in an amino acid substitution, i.e., the TLR4 DNA of the human encodes a variant TLR4. Thus, the invention is useful to detect polymorphisms in the TLR4 gene.

Normal healthy, non-asthmatic subjects demonstrate a reproducible airway response to an incremental LPS inhalation challenge test, with some subjects developing airflow obstruction when challenged with low concentrations of LPS and others virtually unaffected by high concentrations of inhaled LPS. These findings suggest that the spectrum of LPS responsiveness in humans is quite variable from one individual to the next (but reproducible within an individual), and that a substantial portion of the population may be hyporesponsive to inhaled LPS. As described hereinbelow, an incremental LPS inhalation challenge test was employed to reliably phenotype individuals as either responsive (at least a 20% decline in the forced expiratory volume in one second (FEV<sub>1</sub>) after inhaling up to 41.5  $\mu$ g LPS) or hyporesponsive (FEV<sub>1</sub> >80% of their baseline after inhaling 41.5  $\mu$ g of LPS) to inhaled LPS. Fifty-two (63%) of these individuals were responsive to inhaled LPS and 31 (37%) were hyporesponsive to inhaled LPS.

These results were employed to determine the relationship between polymorphisms in the TLR4 gene and the airway response to inhaled LPS in the 83 normal healthy, non-asthmatic subjects. Using single stranded conformational variant (SSCV) analysis and direct sequencing, a missense mutation (A896G) was identified in the fourth exon of the TLR4 gene that results in replacement of a conserved aspartic acid residue with glycine at position 299 in the extracellular domain of the TLR4 receptor. The Asp299Gly sequence variant occurred in 3 LPS responsive (5.8%) and 7 LPS hyporesponsive (22.6%) study subjects ( $p=0.03$ ). Among the subjects with the common TLR4 allele ( $N=73$ ), the dose-response slope (percent decline  $FEV_1$ /cumulative dose of inhaled LPS) averaged a 1.86% decline in  $FEV_1/\mu\text{g}$  inhaled LPS (range 0.01%-19.78%), while the dose-response slope for the subjects with the Asp299Gly allele ( $N=10$ ) was significantly less ( $p=0.007$ ), averaging 0.59% decline in  $FEV_1/\mu\text{g}$  inhaled LPS (range 0.00%-1.59%). Thus, a sequence polymorphism in the TLR4 gene, i.e., a missense mutation (Asp299Gly) in the fourth exon of the TLR4 gene, occurs in a substantial portion of the population, and is associated with an airway hyporesponsive in humans challenged with inhaled LPS. The allelic frequency of the A896G substitution was 6.6% in the study population, 7.9% in a normal control population from Iowa (Lidral et al., 1998), and 3.3% in the parental chromosomes of the CEPH population (NIH-CEPH, 1992). As also described herein, the presence of a TLR4 mutation was associated with gram negative sepsis, severity of sepsis, pre-term delivery, and respiratory distress syndrome in pre-term infants.

The invention also provides an isolated and purified nucleic acid molecule comprising a nucleic acid segment, e.g., genomic DNA or cDNA, encoding TLR4, such as a variant TLR4. Also provided are primers, oligonucleotides and probes comprising the isolated nucleic acid sequences of the invention. The nucleic acid molecules of the invention may be single stranded or double stranded.

Transfection of CHO cells with either the wild-type or the mutant (Asp299Gly) allele of the TLR4 gene demonstrated that this mutation interrupts TLR4-mediated LPS signaling. Moreover, the wild-type allele of TLR4 rescues the LPS hyporesponsive phenotype in either airway epithelial cells or alveolar macrophages obtained from individuals with the TLR4 mutation. Thus, these results provide the first genetic evidence that a common mutation causes differences in LPS responsiveness that may contribute to several disease states in humans.

Therefore, the invention further provides an expression cassette comprising a nucleic acid molecule of the invention, a host cell transformed with the expression cassette, and TLR4 polypeptides isolated therefrom. The transformed host cells, or isolated TLR4 polypeptides, may be useful in identifying agents that modulate, i.e., enhance or inhibit, TLR4 activity. For example, an expression cassette comprising a nucleic acid molecule of the invention which encodes a variant TLR4 polypeptide is introduced to murine cells, e.g., oocytes via microinjection (see Sigmund et al., 1993). The resulting pups are screened for the presence of the nucleic acid molecule. Hence, the invention also provides a transgenic mouse, the genome of the cells of which is augmented with variant human TLR4 DNA. Human TLR4 transgenic mice of the invention have altered innate immunity, e.g., they are more susceptible to gram negative sepsis than their non-transgenic counterparts.

The invention also provides a method to treat an individual at risk of, or having, an indication associated with altered innate immunity, in which an agent that alters TLR4 activity is administered to the individual.

## BRIEF DESCRIPTION OF THE FIGURES

FIG. 1. Genomic structure for the human TLR4 gene (exons are represented by boxes and introns are represented by lines). The coding sequence is in black and the nucleotides encoding the transmembrane domain (TM) are shaded. The positions of the introns in the published TLR4 cDNA sequence (Rock et al., 1998; Genbank Accession No. U88880) are listed. The exon (caps) and intron (lower case) sequences at each of the splice junctions is shown (sequences at 5' and 3' splice junctions for intron 1, SEQ ID NO:1 and SEQ ID NO:63, respectively; for intron 2, SEQ ID NO:2 and SEQ ID NO:64, respectively; and for intron 3, SEQ ID NO:3 and SEQ ID NO:65, respectively).

FIG. 2. The human TLR4 gene is alternatively spliced. Photographs are of agarose gels containing the products amplified from a multi-tissue cDNA panel from human adult (panel a; Clontech #K1420-1, Palo Alto, Calif.), and human fetal (panel b; Clontech #K1425-1, Palo Alto, Calif.) cDNA. The tissues included brain (B), heart (H), kidney (K), liver (Li), lung (Lu), pancreas (Pa), placenta (PI), skeletal muscle (Sk), spleen (Sp), and thymus (T). The first lane of each gel included a 100 bp molecular weight standard (MW). The forward and reverse PCR primers were derived from exons 1 and 4, respectively (see panel c) and amplified three products of 453, 333, and 167 bp. The DNA sequence for these bands showed that both exons 2 and 3 were present in the 453 bp band, that exon 2 was absent in the 333 bp product, and that both exons 2 and 3 were absent in the 167 bp band. The 453 bp and 333 bp sequences are identical to previously published sequences for the human TLR cDNA (Medzhitov et al., 1997; Rock et al., 1998). The open, closed, and shaded boxes indicate the untranslated, translated, and transmembrane domain portions of the TLR4 exons, respectively. The ends of the cDNAs were arbitrarily terminated at the stop codon (STP).

FIG. 3. Mutations in the human TLR4 gene. The SSCV and sequence analysis were performed blinded to the LPS response phenotype of the study subjects. The SSCV gel in panel a contains the products amplified from two samples that are homozygous for the 1060A allele (lanes 1 and 3), a heterozygous sample (lane 2) with both the 896A and 896G alleles, and a homozygous sample (lane 4) with only the 896G allele. The SSCV gel in panel b contains the products amplified from a sample that is homozygous for the T allele at position -11 and the T allele at position 315 (lane 1), and from a sample that is heterozygous with a T and a deletion at position -11 and a T and a C allele at position 315. The nucleotide numbers are based on the previously published TLR4 cDNA sequence (Rock et al., 1998).

FIG. 4. The aspartic acid residue at position 299 is conserved. A portion of the predicted amino acid sequence is aligned for the TLR4 genes from human (Rock et al., 1998; SEQ ID NO:4), mouse (Poitorak et al., 1998; SEQ ID NO:5), rat (Genbank Accession No. AF057025; SEQ ID NO:6), and hamster (D. Golenbock, SEQ ID NO:7). The position of the first amino acid in each sequence is given. The aspartic acid at position 299 is indicated with an arrow. Conserved amino acids are shaded.

FIG. 5. Frequency distribution histogram of the dose-response slope (% decline in  $FEV_1/\mu\text{g}$  LPS). The % decline in  $FEV_1/\mu\text{g}$  LPS was calculated following administration of the cumulative LPS dose that either resulted in at least a 20% decline in  $FEV_1$  or the decline in  $FEV_1$  following a cumulative inhaled dose of 41.5  $\mu\text{g}$  LPS. Subjects above the x-axis (solid bars) are homozygous for the 896A allele (WT/WT); subjects below the x-axis (open bars) are either heterozygous or homozygous (\*) for the missense Asp299Gly allele. The

data is replotted in the inset after the values on the x-axis were log normalized. P values are presented for the comparison of the % decline in FEV<sub>1</sub>/μg LPS between subjects with the WT/WT genotype (N=73) and those with at least one Asp299Gly allele (N=10) using absolute values (P=0.037) and log normalized values (P=0.026). Since the distribution of the dose-response slope (% decline FEV<sub>1</sub>/cumulative dose of inhaled LPS) was highly skewed, the two-sample Monte-Carlo permutation test based on 10,000 permutations was used to calculate P values (Fisher et al., 1993) the missense mutation (N=10) using absolute values (p=0.04) and the log normalized values (p=0.04) and the log normalized values (p=0.025).

FIG. 6. A) CHO/CD-14 cells that express the CD-14 receptor (generous gift of Doug Golenbock) were transfected with a wild-type or mutant TLR4 allele, and then exposed to LPS. Cells were co-transfected with a luciferase reporter construct. B) IL-1α levels from airway epithelial cells of individuals which had been genotyped for TLR4. IL-1α levels before and after LPS exposure are shown. C) IL-1α levels from airway epithelial cells from a TLR4 heterozygote (WT/Asp299Gly). The cells were transfected with a recombinant adenovirus which expresses TLR4 (Rock et al., 1998), or a recombinant adenovirus which expresses green fluorescent protein (GFP), and then exposed to LPS. D) TNF-α levels from alveolar macrophage from a TLR4 homozygote (Asp299Gly/Asp299Gly). The cells were transfected with a recombinant adenovirus which expresses TLR4 (Rock et al., 1998), and then exposed to LPS.

FIG. 7. Codons.

FIG. 8. Exemplary amino acid substitutions.

FIG. 9. A) A partial nucleotide sequence of genomic human TLR4 DNA (SEQ ID NO:62). B) The 5' UTR, exon 1 (nt 1100-1283), and a partial sequence of intron 1 (Genbank Accession No. AF172169; SEQ ID NO:70). C) Exon 2 (nt 192-311), exon 3 (nt 556-722), and intron 2 (244 bp) (Genbank Accession No. AF172170; SEQ ID NO:71). D) Exon 4 (nt 1691-6172) and the 3' UTR (Genbank Accession No. AF172171; SEQ ID NO:72).

#### DETAILED DESCRIPTION OF THE INVENTION

As used herein, the terms "isolated and/or purified" refer to in vitro preparation, isolation and/or purification of a nucleic acid molecule or polypeptide, so that it is not associated with in vivo substances. Thus, with respect to an "isolated nucleic acid molecule", which includes a polynucleotide of genomic, cDNA, or synthetic origin or some combination thereof, the "isolated nucleic acid molecule" (1) is not associated with all or a portion of a polynucleotide in which the "isolated nucleic acid molecule" is found in nature, (2) is operably linked to a polynucleotide which it is not linked to in nature, or (3) does not occur in nature as part of a larger sequence. An isolated nucleic acid molecule means a polymeric form of nucleotides of at least 10 bases in length, either ribonucleotides or deoxy-nucleotides or a modified form of either type of nucleotide. The term includes single and double stranded forms of DNA. The term "oligonucleotide" referred to herein includes naturally occurring, and modified nucleotides linked together by naturally occurring, and non-naturally occurring oligonucleotide linkages. Oligonucleotides are a polynucleotide subset with 200 bases or fewer in length. Preferably, oligonucleotides are 10 to 60 bases in length and most preferably 12, 13, 14, 15, 16, 17, 18, 19, or 20 to 40 bases in length. Oligonucleotides are usually single stranded, e.g., for primers or probes; although oligonucleotides may be double stranded, e.g., for use in the construction of a variant. Oligonucleotides of the

invention can be either sense or antisense oligonucleotides. The term "naturally occurring nucleotides" referred to herein includes deoxyribonucleotides and ribonucleotides. The term "modified nucleotides" referred to herein includes nucleotides with modified or substituted sugar groups and the like. The term "oligonucleotide linkages" referred to herein includes oligonucleotides linkages such as phosphorothioate, phosphorodithioate, phosphoroselenoate, phosphorodiselenoate, phosphoroanilothioate, phosphoraniladate, phosphoroamidate, and the like. An oligonucleotide can include a label for detection, if desired.

The term "isolated polypeptide" means a polypeptide encoded by genomic DNA, cDNA or recombinant RNA, or is synthetic origin, or some combination thereof, which isolated polypeptide (1) is not associated with proteins found in nature, (2) is free of other proteins from the same source, e.g., free of human proteins, (3) is expressed by a cell from a different species, or (4) does not occur in nature.

The term "sequence homology" means the proportion of base matches between two nucleic acid sequences or the proportion amino acid matches between two amino acid sequences. When sequence homology is expressed as a percentage, e.g., 50%, the percentage denotes the proportion of matches over the length of sequence from one TLR4 allele that is compared to another TLR4 allele. Gaps (in either of the two sequences) are permitted to maximize matching; gap lengths of 15 bases or less are usually used, 6 bases or less are preferred with 2 bases or less more preferred. When using oligonucleotides as probes or treatments, the sequence homology between the target nucleic acid and the oligonucleotide sequence is generally not less than 17 target base matches out of 20 possible oligonucleotide base pair matches (85%); preferably not less than 9 matches out of 10 possible base pair matches (90%), and more preferably not less than 19 matches out of 20 possible base pair matches (95%).

The term "selectively hybridize" means to detectably and specifically bind. Polynucleotides, oligonucleotides and fragments of the invention selectively hybridize to nucleic acid strands under hybridization and wash conditions that minimize appreciable amounts of detectable binding to non-specific nucleic acids. High stringency conditions can be used to achieve selective hybridization conditions as known in the art and discussed herein. Generally, the nucleic acid sequence homology between the polynucleotides, oligonucleotides, and fragments of the invention and a nucleic acid sequence of interest is at least 65%, and more typically with preferably increasing homologies of at least about 70%, about 90%, about 95%, about 98%, and 100%.

Two amino acid sequences are homologous if there is a partial or complete identity between their sequences. For example, 85% homology means that 85% of the amino acids are identical when the two sequences are aligned for maximum matching. Gaps (in either of the two sequences being matched) are allowed in maximizing matching; gap lengths of 5 or less are preferred with 2 or less being more preferred. Alternatively and preferably, two protein sequences (or polypeptide sequences derived from them of at least 30 amino acids in length) are homologous, as this term is used herein, if they have an alignment score of at more than 5 (in standard deviation units) using the program ALIGN with the mutation data matrix and a gap penalty of 6 or greater. See Dayhoff, M. O., in Atlas of Protein Sequence and Structure, 1972, volume 5, National Biomedical Research Foundation, pp. 101-110, and Supplement 2 to this volume, pp. 1-10. The two sequences or parts thereof are more preferably homologous if their amino acids are greater than or equal to 50% identical when optimally aligned using the ALIGN program.

The term “corresponds to” is used herein to mean that a polynucleotide sequence is homologous (i.e., is identical, not strictly evolutionarily related) to all or a portion of a reference polynucleotide sequence, or that a polypeptide sequence is identical to a reference polypeptide sequence. In contradistinction, the term “complementary to” is used herein to mean that the complementary sequence is homologous to all or a portion of a reference polynucleotide sequence. For illustration, the nucleotide sequence “TATAC” corresponds to a reference sequence “TATAC” and is complementary to a reference sequence “GTATA”.

The following terms are used to describe the sequence relationships between two or more polynucleotides: “reference sequence”, “comparison window”, “sequence identity”, “percentage of sequence identity”, and “substantial identity”. A “reference sequence” is a defined sequence used as a basis for a sequence comparison; a reference sequence may be a subset of a larger sequence, for example, as a segment of a full-length cDNA or gene sequence given in a sequence listing, or may comprise a complete cDNA or gene sequence. Generally, a reference sequence is at least 20 nucleotides in length, frequently at least 25 nucleotides in length, and often at least 50 nucleotides in length. Since two polynucleotides may each (1) comprise a sequence (i.e., a portion of the complete polynucleotide sequence) that is similar between the two polynucleotides, and (2) may further comprise a sequence that is divergent between the two polynucleotides, sequence comparisons between two (or more) polynucleotides are typically performed by comparing sequences of the two polynucleotides over a “comparison window” to identify and compare local regions of sequence similarity.

A “comparison window”, as used herein, refers to a conceptual segment of at least 20 contiguous nucleotides and wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) of 20 percent or less as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. Optimal alignment of sequences for aligning a comparison window may be conducted by the local homology algorithm of Smith and Waterman (1981) *Adv. Appl. Math.* 2: 482, by the homology alignment algorithm of Needleman and Wunsch (1970) *J. Mol. Biol.* 48: 443, by the search for similarity method of Pearson and Lipman (1988) *Proc. Natl. Acad. Sci. (U.S.A.)* 85: 2444, by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package Release 7.0, Genetics Computer Group, 575 Science Dr., Madison, Wis.), or by inspection, and the best alignment (i.e., resulting in the highest percentage of homology over the comparison window) generated by the various methods is selected.

The term “sequence identity” means that two polynucleotide sequences are identical (i.e., on a nucleotide-by-nucleotide basis) over the window of comparison. The term “percentage of sequence identity” means that two polynucleotide sequences are identical (i.e., on a nucleotide-by-nucleotide basis) over the window of comparison. The term “percentage of sequence identity” is calculated by comparing two optimally aligned sequences over the window of comparison, determining the number of positions at which the identical nucleic acid base (e.g., A, T, C, G, U, or I) occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison (i.e., the window size), and multiplying the result by 100 to yield the percentage of sequence identity. The terms “substantial identity” as used herein denote a characteristic of a polynucleotide

sequence, wherein the polynucleotide comprises a sequence that has at least 85 percent sequence identity, preferably at least 90 to 95 percent sequence identity, more usually at least 99 percent sequence identity as compared to a reference sequence over a comparison window of at least 20 nucleotide positions, frequently over a window of at least 20-50 nucleotides, wherein the percentage of sequence identity is calculated by comparing the reference sequence to the polynucleotide sequence which may include deletions or additions which total 20 percent or less of the reference sequence over the window of comparison.

As applied to polypeptides, the term “substantial identity” means that two peptide sequences, when optimally aligned, such as by the programs GAP or BESTFIT using default gap weights, share at least about 80 percent sequence identity, preferably at least about 90 percent sequence identity, more preferably at least about 95 percent sequence identity, and most preferably at least about 99 percent sequence identity.

As used herein, “substantially pure” means an object species is the predominant species present (i.e., on a molar basis it is more abundant than any other individual species in the composition), and preferably a substantially purified fraction is a composition wherein the object species comprises at least about 50 percent (on a molar basis) of all macromolecular species present. Generally, a substantially pure composition will comprise more than about 80 percent of all macromolecular species present in the composition, more preferably more than about 85%, about 90%, about 95%, and about 99%. Most preferably, the object species is purified to essential homogeneity (contaminant species cannot be detected in the composition by conventional detection methods) wherein the composition consists essentially of a single macromolecular species.

An isolated “variant” TLR4 polypeptide has at least 50%, preferably at least about 80%, and more preferably at least about 90%, but less than 100%, contiguous amino acid sequence homology or identity to the amino acid sequence of a reference (wild-type) TLR4 polypeptide. Preferably, the TLR4 polypeptides of the invention are biologically active. Biologically active polypeptides include those that induce an immune response when administered to an organism, are bound by antibodies specific for TLR4, activate signaling through NF-KB and AP-1, interact with CD14, or induce cytokine release following LPS stimulation. While it is preferred that a variant TLR4 has at least about 0.1%, preferably at least about 1%, and more preferably at least about 10%, of the activity of wild-type TLR4, the invention includes variant TLR4 polypeptides having no detectable biological activity. Likewise, a “variant” TLR4 nucleic acid molecule has at least about 80%, preferably at least about 90% and more preferably at least about 95%, but less than 100% contiguous nucleic acid sequence homology or identity to the nucleic acid sequence of a wild-type TLR4 gene.

As used herein, an “indication or condition associated with aberrant, modified or altered innate immunity” includes, but is not limited to, hyporesponsiveness to LPS, susceptibility to infection with gram-negative bacteria, susceptibility to sepsis by gram-negative bacteria, susceptibility to chronic airway disease, susceptibility to asthma, susceptibility to arthritis, susceptibility to pyelonephritis, susceptibility to gall bladder disease, susceptibility to pneumonia, susceptibility to bronchitis, susceptibility to chronic obstructive pulmonary disease, severity of cystic fibrosis, and susceptibility to local and systemic inflammatory conditions, e.g., systemic inflammatory response syndrome (SIRS), local gram negative bacterial infection, or acute respiratory distress syndrome (ARDS).

## A. Nucleic Acid Molecules of the Invention

## 1. Sources of the Nucleic Acid Molecules of the Invention

Sources of nucleotide sequences from which the present nucleic acid molecules encoding TLR4, a portion (fragment) thereof, a variant thereof or the nucleic acid complement thereof, include total or polyA<sup>+</sup> RNA from any mammalian, preferably human, cellular source from which cDNAs can be derived by methods known in the art. Other sources of the DNA molecules of the invention include genomic libraries derived from any mammalian cellular source. Moreover, the present DNA molecules may be prepared in vitro, e.g., by synthesizing an oligonucleotide of about 100, preferably about 75, more preferably about 50, and even more preferably about 40, nucleotides in length, or by subcloning a portion of a DNA segment that encodes a particular TLR4.

## 2. Isolation of a Gene Encoding TLR4

A nucleic acid molecule encoding TLR4 can be identified and isolated using standard methods, as described by Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor, NY (1989). For example, reverse-transcriptase PCR (RT-PCR) can be employed to isolate and clone TLR4 cDNAs. Oligo-dT can be employed as a primer in a reverse transcriptase reaction to prepare first-strand cDNAs from isolated RNA which contains RNA sequences of interest, e.g., total RNA isolated from human tissue. RNA can be isolated by methods known to the art, e.g., using TRIZOL™ reagent (GIBCO-BRL/Life Technologies, Gaithersburg, Md.). Resultant first-strand cDNAs are then amplified in PCR reactions.

“Polymerase chain reaction” or “PCR” refers to a procedure or technique in which amounts of a preselected fragment of nucleic acid, RNA and/or DNA, are amplified as described in U.S. Pat. No. 4,683,195. Generally, sequence information from the ends of the region of interest or beyond is employed to design oligonucleotide primers comprising at least 7-8 nucleotides. These primers will be identical or similar in sequence to opposite strands of the template to be amplified. PCR can be used to amplify specific RNA sequences, specific DNA sequences from total genomic DNA, and cDNA transcribed from total cellular RNA, bacteriophage or plasmid sequences, and the like. See generally Mullis et al., *Cold Spring Harbor Symp. Quant. Biol.*, 51, 263 (1987); Erlich, ed., *PCR Technology*, (Stockton Press, NY, 1989). Thus, PCR-based cloning approaches rely upon conserved sequences deduced from alignments of related gene or polypeptide sequences.

Primers are made to correspond to highly conserved regions of polypeptides or nucleotide sequences which were identified and compared to generate the primers, e.g., by a sequence comparison of other mammalian TLR4. One primer is prepared which is predicted to anneal to the antisense strand, and another primer prepared which is predicted to anneal to the sense strand, of a DNA molecule which encodes TLR4.

The products of each PCR reaction are separated via an agarose gel and all consistently amplified products are gel-purified and cloned directly into a suitable vector, such as a known plasmid vector. The resultant plasmids are subjected to restriction endonuclease and dideoxy sequencing of double-stranded plasmid DNAs.

Another approach to identify, isolate and clone cDNAs which encode TLR4 is to screen a cDNA or genomic library. Screening for DNA fragments that encode all or a portion of a DNA encoding TLR4 can be accomplished by probing the library with a probe which has sequences that are highly conserved between genes believed to be related to TLR4, e.g.,

the homolog of a particular TLR4 from a different species, or by screening of plaques for binding to antibodies that specifically recognize TLR4. DNA fragments that bind to a probe having sequences which are related to TLR4, or which are immunoreactive with antibodies to TLR4, can be subcloned into a suitable vector and sequenced and/or used as probes to identify other cDNAs encoding all or a portion of TLR4.

Thus, “isolated and/or purified TLR4” nucleic acid refers to in vitro isolation of a nucleic acid molecule from its natural cellular environment, and from association with other components of the cell, such as nucleic acid or polypeptide, so that it can be sequenced, replicated, and/or expressed. For example, “isolated TLR4 nucleic acid” is RNA or DNA containing greater than 9, preferably 36, and more preferably 45 or more, sequential nucleotide bases that encode at least a portion of TLR4, a variant thereof, RNA or DNA complementary thereto, or which hybridizes to, RNA or DNA comprising TLR4 sequences, and remains stably bound under stringent conditions, as defined by methods well known in the art, e.g., in Sambrook et al., *supra*. Thus, the RNA or DNA is “isolated” in that it is free from at least one contaminating nucleic acid with which it is normally associated in the natural source of the RNA or DNA and is preferably substantially free of any other mammalian RNA or DNA. The phrase “free from at least one contaminating source nucleic acid with which it is normally associated” includes the case where the nucleic acid is reintroduced into the source or natural cell but is in a different chromosomal location or is otherwise flanked by nucleic acid sequences not normally found in the source cell.

As used herein, the term “recombinant nucleic acid”, e.g., “recombinant DNA sequence or segment”, refers to a nucleic acid, e.g., to DNA, that has been derived or isolated from any appropriate tissue source, that may be subsequently chemically altered in vitro, so that its sequence is not naturally occurring, or corresponds to naturally occurring sequences that are not positioned as they would be positioned in a genome which has not been transformed with exogenous DNA. An example of DNA “derived” from a source, would be a DNA sequence that is identified as a useful fragment within a given organism, and which is then chemically synthesized in essentially pure form. An example of such DNA “isolated” from a source would be a useful DNA sequence that is excised or removed from said source by chemical means, e.g., by the use of restriction endonucleases, so that it can be further manipulated, e.g., amplified, for use in the invention, by the methodology of genetic engineering.

Thus, recovery or isolation of a given fragment of DNA from a restriction digest can employ separation of the digest on polyacrylamide or agarose gel by electrophoresis, identification of the fragment of interest by comparison of its mobility versus that of marker DNA fragments of known molecular weight, removal of the gel section containing the desired fragment, and separation of the gel from DNA. See Lawn et al., *Nucleic Acids Res.*, 9, 6103 (1981), and Goeddel et al., *Nucleic Acids Res.*, 8, 4057 (1980). Therefore, “isolated DNA” includes completely synthetic DNA sequences, semi-synthetic DNA sequences, DNA sequences isolated from biological sources, and DNA sequences derived from RNA, as well as mixtures thereof.

As used herein, the term “derived” with respect to a RNA molecule means that the RNA molecule has complementary sequence identity to a particular DNA molecule.

## 3. Variants of the Nucleic Acid Molecules of the Invention

Nucleic acid molecules encoding amino acid sequence variants of TLR4 are prepared by a variety of methods known

in the art. These methods include, but are not limited to, isolation from a natural source (in the case of naturally occurring amino acid sequence variants) or preparation by oligonucleotide-mediated (or site-directed) mutagenesis, PCR mutagenesis, and cassette mutagenesis of an earlier prepared variant or a non-variant version of TLR4 nucleic acid.

Oligonucleotide-mediated mutagenesis is a preferred method for preparing amino acid substitution variants of TLR4. This technique is well known in the art as described by Adelman et al., *DNA*, 2, 183 (1983). Briefly, TLR4 DNA is altered by hybridizing an oligonucleotide encoding the desired mutation to a DNA template, where the template is the single-stranded form of a plasmid or bacteriophage containing the unaltered or native DNA sequence of TLR4. After hybridization, a DNA polymerase is used to synthesize an entire second complementary strand of the template that will thus incorporate the oligonucleotide primer, and will code for the selected alteration in the TLR4 DNA.

Generally, oligonucleotides of at least 25 nucleotides in length are used. An optimal oligonucleotide will have 12 to 15 nucleotides that are completely complementary to the template on either side of the nucleotide(s) coding for the mutation. This ensures that the oligonucleotide will hybridize properly to the single-stranded DNA template molecule. The oligonucleotides are readily synthesized using techniques known in the art such as that described by Crea et al., *Proc. Natl. Acad. Sci. U.S.A.*, 75, 5765 (1978).

The DNA template can be generated by those vectors that are either derived from bacteriophage M13 vectors (the commercially available M13mp18 and M13mp19 vectors are suitable), or those vectors that contain a single-stranded phage origin of replication as described by Viera et al., *Meth. Enzymol.*, 153, 3 (1987). Thus, the DNA that is to be mutated may be inserted into one of these vectors to generate single-stranded template. Production of the single-stranded template is described in Sections 4.214.41 of Sambrook et al., *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Laboratory Press, N.Y. 1989).

Alternatively, single-stranded DNA template may be generated by denaturing double-stranded plasmid (or other) DNA using standard techniques.

For alteration of the native DNA sequence (to generate amino acid sequence variants, for example), the oligonucleotide is hybridized to the single-stranded template under suitable hybridization conditions. A DNA polymerizing enzyme, usually the Klenow fragment of DNA polymerase I, is then added to synthesize the complementary strand of the template using the oligonucleotide as a primer for synthesis. A heteroduplex molecule is thus formed such that one strand of DNA encodes the mutated form of TLR4, and the other strand (the original template) encodes the native, unaltered sequence of TLR4. This heteroduplex molecule is then transformed into a suitable host cell, usually a prokaryote such as *E. coli* JM101. After the cells are grown, they are plated onto agarose plates and screened using the oligonucleotide primer radiolabeled with 32-phosphate to identify the bacterial colonies that contain the mutated DNA. The mutated region is then removed and placed in an appropriate vector for peptide or polypeptide production, generally an expression vector of the type typically employed for transformation of an appropriate host.

The method described immediately above may be modified such that a homoduplex molecule is created wherein both strands of the plasmid contain the mutations(s). The modifications are as follows: The single-stranded oligonucleotide is annealed to the single-stranded template as described above. A mixture of three deoxyribonucleotides, deoxyriboadenosine (dATP), deoxyribothymine (dTTP), and deoxyribo-

midine (dTTP), is combined with a modified thiodeoxyribo-cytosine called dCTP-( $\alpha$ S) (which can be obtained from the Amersham Corporation). This mixture is added to the template-oligonucleotide complex. Upon addition of DNA polymerase to this mixture, a strand of DNA identical to the template except for the mutated bases is generated. In addition, this new strand of DNA will contain dCTP-( $\alpha$ S) instead of dCTP, which serves to protect it from restriction endonuclease digestion.

After the template strand of the double-stranded heteroduplex is nicked with an appropriate restriction enzyme, the template strand can be digested with ExoIII nuclease or another appropriate nuclease past the region that contains the site(s) to be mutagenized. The reaction is then stopped to leave a molecule that is only partially single-stranded. A complete double-stranded DNA homoduplex is then formed using DNA polymerase in the presence of all four deoxyribonucleotide triphosphates, ATP, and DNA ligase. This homoduplex molecule can then be transformed into a suitable host cell such as *E. coli* JM101.

For example, a preferred embodiment of the invention is an isolated and purified DNA molecule comprising a human DNA segment encoding a variant TLR4 having a nucleotide substitution at position 896 (A896G) which encodes an amino acid substitution (Asp299Gly). Other nucleotide substitutions which result in silent mutations, missense mutations, or a nonsense mutations, can be ascertained by reference to FIG. 7, FIG. 8 and page D1 in Appendix D in Sambrook et al., *Molecular Cloning: A Laboratory Manual* (1989).

#### B. Chimeric Expression Cassettes

To prepare expression cassettes for transformation herein, the recombinant DNA sequence or segment may be circular or linear, double-stranded or single-stranded. A DNA sequence which encodes an RNA sequence that is substantially complementary to a mRNA sequence encoding TLR4 is typically a "sense" DNA sequence cloned into a cassette in the opposite orientation (i.e., 3' to 5' rather than 5' to 3'). Generally, the DNA sequence or segment is in the form of chimeric DNA, such as plasmid DNA, that can also contain coding regions flanked by control sequences which promote the expression of the DNA present in the resultant cell line.

As used herein, "chimeric" means that a vector comprises DNA from at least two different species, or comprises DNA from the same species, which is linked or associated in a manner which does not occur in the "native" or wild type of the species.

Aside from DNA sequences that serve as transcription units for TLR4, or portions thereof, a portion of the recombinant DNA may be untranscribed, serving a regulatory or a structural function. For example, the DNA may itself comprise a promoter that is active in mammalian cells, or may utilize a promoter already present in the genome that is the transformation target. Such promoters include the CMV promoter, as well as the SV40 late promoter and retroviral LTRs (long terminal repeat elements), although many other promoter elements well known to the art may be employed in the practice of the invention.

Other elements functional in the host cells, such as introns, enhancers, polyadenylation sequences and the like, may also be a part of the DNA. Such elements may or may not be necessary for the function of the DNA, but may provide improved expression of the DNA by affecting transcription, stability of the mRNA, or the like. Such elements may be included in the DNA as desired to obtain the optimal performance of the transforming DNA in the cell.

“Control sequences” is defined to mean DNA sequences necessary for the expression of an operably linked coding sequence in a particular host organism. The control sequences that are suitable for prokaryotic cells, for example, include a promoter, and optionally an operator sequence, and a ribosome binding site. Eukaryotic cells are known to utilize promoters, polyadenylation signals, and enhancers.

“Operably linked” is defined to mean that the nucleic acids are placed in a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is operably linked to DNA for a peptide or polypeptide if it is expressed as a preprotein that participates in the secretion of the peptide or polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, “operably linked” means that the DNA sequences being linked are contiguous and, in the case of a secretory leader, contiguous and in reading phase. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, the synthetic oligonucleotide adaptors or linkers are used in accord with conventional practice.

The DNA to be introduced into the cells further will generally contain either a selectable marker gene or a reporter gene or both to facilitate identification and selection of transformed cells from the population of cells sought to be transformed. Alternatively, the selectable marker may be carried on a separate piece of DNA and used in a co-transformation procedure. Both selectable markers and reporter genes may be flanked with appropriate regulatory sequences to enable expression in the host cells. Useful selectable markers are well known in the art and include, for example, antibiotic and herbicide-resistance genes, such as *neo*, *hpt*, *dhfr*, *bar*, *aroA*, *dapA* and the like. See also, the genes listed on Table 1. of Lundquist et al. (U.S. Pat. No. 5,848,956).

Reporter genes are used for identifying potentially transformed cells and for evaluating the functionality of regulatory sequences. Reporter genes which encode for easily assayable proteins are well known in the art. In general, a reporter gene is a gene which is not present in or expressed by the recipient organism or tissue and which encodes a protein whose expression is manifested by some easily detectable property, e.g., enzymatic activity. Preferred genes include the chloramphenicol acetyl transferase gene (*cat*) from Tn9 of *E. coli*, the beta-glucuronidase gene (*gus*) of the *uidA* locus of *E. coli*, and the luciferase *luc* gene from firefly *Photinus pyralis*. Expression of the reporter gene is assayed at a suitable time after the DNA has been introduced into the recipient cells.

The general methods for constructing recombinant DNA which can transform target cells are well known to those skilled in the art, and the same compositions and methods of construction may be utilized to produce the DNA useful herein. For example, J. Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press (2d ed., 1989), provides suitable methods of construction.

### C. Transformation into Host Cells

The recombinant DNA can be readily introduced into the host cells, e.g., mammalian, bacterial, yeast or insect cells by

transfection with an expression vector comprising DNA encoding TLR4 or its complement, by any procedure useful for the introduction into a particular cell, e.g., physical or biological methods, to yield a transformed cell having the recombinant DNA stably integrated into its genome, so that the DNA molecules, sequences, or segments, of the present invention are expressed by the host cell.

Physical methods to introduce a DNA into a host cell include calcium phosphate precipitation, lipofection, particle bombardment, microinjection, electroporation, and the like. Biological methods to introduce the DNA of interest into a host cell include the use of DNA and RNA viral vectors. The main advantage of physical methods is that they are not associated with pathological or oncogenic processes of viruses. However, they are less precise, often resulting in multiple copy insertions, random integration, disruption of foreign and endogenous gene sequences, and unpredictable expression. Viral vectors, useful to introduce genes to mammalian cells include, but are not limited to, poxvirus vectors, herpes simplex virus I vectors, adenovirus vectors, adeno-associated virus vectors, and the like.

As used herein, the term “cell line” or “host cell” is intended to refer to well-characterized homogenous, biologically pure populations of cells. These cells may be eukaryotic cells that are neoplastic or which have been “immortalized” in vitro by methods known in the art, as well as primary cells, or prokaryotic cells. The cell line or host cell is preferably of mammalian origin, but cell lines or host cells of non-mammalian origin may be employed, including plant, insect, yeast, fungal or bacterial sources. Generally, the DNA sequence is related to a DNA sequence which is resident in the genome of the host cell but is not expressed, or not highly expressed, or, alternatively, overexpressed.

“Transfected” or “transformed” is used herein to include any host cell or cell line, the genome of which has been altered or augmented by the presence of at least one DNA sequence, which DNA is also referred to in the art of genetic engineering as “heterologous DNA,” “recombinant DNA,” “exogenous DNA,” “genetically engineered,” “non-native,” or “foreign DNA,” wherein said DNA was isolated and introduced into the genome of the host cell or cell line by the process of genetic engineering. The host cells of the present invention are typically produced by transfection with a DNA sequence in a plasmid expression vector, a viral expression vector, or as an isolated linear DNA sequence.

To confirm the presence of the DNA sequence in the host cell, a variety of assays may be performed. Such assays include, for example, “molecular biological” assays well known to those of skill in the art, such as Southern and Northern blotting, RT-PCR and PCR; “biochemical” assays, such as detecting the presence or absence of a particular TLR4, e.g., by immunological means (ELISAs and Western blots) or by assays described herein.

To detect and quantitate RNA produced from introduced DNA segments, RT-PCR may be employed. In this application of PCR, it is first necessary to reverse transcribe RNA into DNA, using enzymes such as reverse transcriptase, and then through the use of conventional PCR techniques amplify the DNA. In most instances PCR techniques, while useful, will not demonstrate integrity of the RNA product. Further information about the nature of the RNA product may be

obtained by Northern blotting. This technique demonstrates the presence of an RNA species and gives information about the integrity of that RNA. The presence or absence of an RNA species can also be determined using dot or slot blot Northern hybridizations. These techniques are modifications of Northern blotting and only demonstrate the presence or absence of an RNA species.

While Southern blotting and PCR may be used to detect the DNA segment in question, they do not provide information as to whether the DNA segment is being expressed. Expression may be evaluated by specifically identifying the polypeptide products of the introduced DNA sequences or evaluating the phenotypic changes brought about by the expression of the introduced DNA segment in the host cell.

#### D. TLR4 Polypeptides, Variants, and Derivatives Thereof

The present isolated, purified TLR4 polypeptides, variants or derivatives thereof, can be synthesized in vitro, e.g., by the solid phase peptide synthetic method or by recombinant DNA approaches (see above). The solid phase peptide synthetic method is an established and widely used method, which is described in the following references: Stewart et al., *Solid Phase Peptide Synthesis*, W. H. Freeman Co., San Francisco (1969); Merrifield, *J. Am. Chem. Soc.*, 85 2149 (1963); Meienhofer in "Hormonal Proteins and Peptides," ed.; C. H. Li, Vol. 2 (Academic Press, 1973), pp. 48-267; Bavaay and Merrifield, "The Peptides," eds. E. Gross and F. Meienhofer, Vol. 2 (Academic Press, 1980) pp. 3-285; and Clark-Lewis et al., *Meth. Enzymol.*, 287, 233 (1997). These peptides can be further purified by fractionation on immunoaffinity or ion-exchange columns; ethanol precipitation; reverse phase HPLC; chromatography on silica or on an anion-exchange resin such as DEAE; chromatofocusing; SDS-PAGE; ammonium sulfate precipitation; gel filtration using, for example, Sephadex G-75; or ligand affinity chromatography.

Once isolated and characterized, derivatives, e.g., chemically derived derivatives, of a given TLR4 can be readily prepared. For example, amides of the TLR4 or TLR4 variants of the present invention may also be prepared by techniques well known in the art for converting a carboxylic acid group or precursor, to an amide. A preferred method for amide formation at the C-terminal carboxyl group is to cleave the peptide from a solid support with an appropriate amine, or to cleave in the presence of an alcohol, yielding an ester, followed by aminolysis with the desired amine.

Salts of carboxyl groups of a peptide or peptide variant of the invention may be prepared in the usual manner by contacting the peptide with one or more equivalents of a desired base such as, for example, a metallic hydroxide base, e.g., sodium hydroxide; a metal carbonate or bicarbonate base such as, for example, sodium carbonate or sodium bicarbonate; or an amine base such as, for example, triethylamine, triethanolamine, and the like.

N-acyl derivatives of an amino group of the TLR4 or variant thereof may be prepared by utilizing an N-acyl protected amino acid for the final condensation, or by acylating a protected or unprotected peptide. O-acyl derivatives may be prepared, for example, by acylation of a free hydroxy peptide or peptide resin. Either acylation may be carried out using standard acylating reagents such as acyl halides, anhydrides, acyl

imidazoles, and the like. Both N- and O-acylation may be carried out together, if desired.

Formyl-methionine, pyroglutamine and trimethyl-alanine may be substituted at the N-terminal residue of the peptide or peptide variant. Other amino-terminal modifications include aminooxypentane modifications (see Simmons et al., *Science*, 276, 276 (1997)).

In addition, the amino acid sequence of TLR4 can be modified so as to result in a variant TLR4. The modification includes the substitution of at least one amino acid residue in the polypeptide for another amino acid residue, including substitutions which utilize the D rather than L form, as well as other well known amino acid analogs, e.g., unnatural amino acids such as  $\alpha$ ,  $\alpha$ -disubstituted amino acids, N-alkyl amino acids, lactic acid, and the like. These analogs include phosphoserine, phosphothreonine, phosphotyrosine, hydroxyproline, gamma-carboxyglutamate; hippuric acid, octahydroindole-2-carboxylic acid, statine, 1,2,3,4-tetrahydroisoquinoline-3-carboxylic acid, penicillamine, ornithine, citrulline,  $\alpha$ -methyl-alanine, para-benzoyl-phenylalanine, phenylglycine, propargylglycine, sarcosine,  $\epsilon$ -N,N,N-trimethyllysine,  $\epsilon$ -N-acetyllysine, N-acetylserine, N-formylmethionine, 3-methylhistidine, 5-hydroxylysine,  $\omega$ -N-methylarginine, and other similar amino acids and imino acids and tert-butylglycine.

Conservative amino acid substitutions include aspartic-glutamic as acidic amino acids; lysine/arginine/histidine as basic amino acids; leucine/isoleucine, methionine/valine, alanine/valine as hydrophobic amino acids; serine/glycine/alanine/threonine as hydrophilic amino acids. Conservative amino acid substitution also includes groupings based on side chains. For example, a group of amino acids having aliphatic side chains is glycine, alanine, valine, leucine, and isoleucine; a group of amino acids having aliphatic-hydroxyl side chains is serine and threonine; a group of amino acids having amide-containing side chains is asparagine and glutamine; a group of amino acids having aromatic side chains is phenylalanine, tyrosine, and tryptophan; a group of amino acids having basic side chains is lysine, arginine, and histidine; and a group of amino acids having sulfur-containing side chains is cysteine and methionine. For example, it is reasonable to expect that replacement of a leucine with an isoleucine or valine, an aspartate with a glutamate, a threonine with a serine, or a similar replacement of an amino acid with a structurally related amino acid will not have a major effect on the properties of the resulting variant polypeptide. Whether an amino acid change results in a functional polypeptide can readily be determined by assaying the activity of the polypeptide variant. Such assays are described herein.

Conservative substitutions are shown in FIG. 8 under the heading of exemplary substitutions. More preferred substitutions are under the heading of preferred substitutions. After the substitutions are introduced, the variants are screened for biological activity.

Amino acid substitutions falling within the scope of the invention, are, in general, accomplished by selecting substitutions that do not differ significantly in their effect on maintaining (a) the structure of the peptide backbone in the area of the substitution, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain.



Naturally occurring residues are divided into groups based on common side-chain properties:

- (1) hydrophobic: norleucine, met, ala, val, leu, ile;
- (2) neutral hydrophilic: cys, ser, thr;
- (3) acidic: asp, glu;
- (4) basic: asn, gln, his, lys, arg;
- (5) residues that influence chain orientation: gly, pro; and
- (6) aromatic; trp, tyr, phe.

The invention also envisions polypeptide variants with non-conservative substitutions. Non-conservative substitutions entail exchanging a member of one of the classes described above for another.

Acid addition salts of the polypeptide or variant polypeptide or of amino residues of the polypeptide or variant polypeptide may be prepared by contacting the polypeptide or amine with one or more equivalents of the desired inorganic or organic acid, such as, for example, hydrochloric acid. Esters of carboxyl groups of the polypeptides may also be prepared by any of the usual methods known in the art.

#### E. Methods of the Invention

The invention provides a method to diagnose individuals who are at a greater risk of deleterious consequences due to bacterial infection, e.g., the individuals may be more susceptible to infection by gram-negative bacteria, e.g., local gram-negative infection, more susceptible to sepsis induced by gram-negative bacteria, more susceptible to chronic airway disease, more susceptible to asthma, more susceptible to gall bladder disease, more susceptible to pyelonephritis, more susceptible to pneumonia, more susceptible to bronchitis, more susceptible to chronic obstructive pulmonary disease, more susceptible to arthritis, at higher risk for severe cystic fibrosis, and more susceptible to local and systemic inflammatory conditions such as systemic inflammatory response syndrome (SIRS), and acute respiratory distress syndrome (ARDS). The invention is also useful in the development of drugs that target the TLR4 gene product, e.g., increase or decrease the function of TLR4, especially the extracellular domain. These agents may thus be useful to prevent or ameliorate infection by gram-negative bacteria, prevent or ameliorate sepsis induced by gram-negative bacteria, prevent or ameliorate LPS-induced chronic airway disease in normal, cystic fibrosis and asthmatic populations, prevent or ameliorate arthritis, and prevent or ameliorate local and systemic inflammatory conditions such as SIRS and ARDS, particularly in individuals at risk for these indications or conditions.

The invention will be further described by the following non-limiting examples.

#### EXAMPLE 1

##### Methods

**Study Subjects.** The study population consisted of 83 healthy adult volunteers (31 men, 52 women) aged 18-50. Exclusion criteria included any history of allergies, tobacco use, cardiac or pulmonary disease. After written informed consent was obtained, all subjects were screened with spirometry, inhalation challenge with histamine, skin testing for common aeroallergens, chest x-ray, and electrocardiogram. All participants had normal screening studies (including histamine PC<sub>20</sub>>32 mg/ml), were on no medications (ex-

cept birth control), and had no significant acute or chronic cardiopulmonary disease or occupational exposures.

**Inhalation Challenge Protocol.** All subjects were exposed by inhalation challenge to buffered sterile saline (HBSS) followed by increasing concentrations of LPS. The solutions were delivered via a DeVilbiss 646 nebulizer powered by compressed air at 30 psi (DeVilbiss Co., Somerset, Pa.) and a Rosenthal dosimeter (Laboratory for Applied Immunology, Baltimore, Md.). After the HBSS, subsequent inhalations delivered in increasing doses of LPS according to the following schedule: 0.5 µg, 1.0 µg, 2.0 µg, 3.0 µg, 5.0 µg, 10 µg, and 20 µg. Thus, the entire protocol delivered a total of 41.5 µg of LPS.

**Incremental LPS inhalation challenge.** The incremental LPS inhalation challenge was performed as follows. The % decline in FEV<sub>1</sub>/µg LPS was calculated following administration of the cumulative LPS dose that either resulted in at least a 20% decline in FEV<sub>1</sub> or the decline in FEV<sub>1</sub> following a cumulative inhaled dose of 41.5 µg of LPS. Subjects above the x-axis (solid bars in FIG. 5) are homozygous for the wild type allele (WT/WT); subjects below the x-axis (open bars in FIG. 5) are either heterozygous or homozygous (\*) for the missense Asp299Gly allele. The data is replotted in the inset after the values for the dose-response slope were log normalized. P values are presented for the comparison of the % decline in FEV<sub>1</sub>/µg LPS between subjects with the WT/WT genotype (N=73) and those with at least one Asp299Gly allele (N=10) using absolute values (P=0.037) and log normalized values (P=0.026). Since the distribution of the dose-response slope (% decline FEV<sub>1</sub>/cumulative dose of inhaled LPS) was highly skewed, the two-sample Monte-Carlo permutation test based on 10,000 permutations was used to calculate P values (Fisher et al., 1993). To assess the allelic frequency of TLR4 sequence variants, a well-characterized Iowa population (Lidral et al., 1998) and the Centre d'Etude du Polymorphisme Humain (CEPH) population (NIH-CEPIT, 1992) were screened for specific sequence variants identified in the 83 study subjects.

**Endotoxin.** Solutions of endotoxin for inhalation were prepared according to a standard protocol using lyophilized *Escherichia coli* (serotype 0111:B4, Sigma Chemical Co., St. Louis, Mo.) LPS. These solutions of LPS were resuspended in sterile Hank's balanced salt solution (without calcium or magnesium) at a pH of 7.0 and filter sterilized. All solutions used for inhalation were tested for sterility (bacteria and fungi) and LPS content (*Limulus amoebocyte* lysate assay, QCL-1000; Whittaker Bioproducts, Walkersville, Md.) prior to separation into individual aliquots. These aliquots were stored immediately after preparation at -70° C. until used.

**Physiologic Measurements.** A Spirotech (Atlanta, Ga.) S600 spirometer was used to assess pulmonary function; spirometry was performed using standards established by the American Thoracic Society. Subjects were positioned upright in a chair and were using noseclips. Baseline spirometry was recorded after inhalation of saline, and then 1, 10, 20, and 30 minutes following inhalation of each dose of LPS, and compared with the post-saline baseline spirometry. If the study subject's FEV<sub>1</sub> was greater than 80% of the baseline measurement at the final assessment (30 minutes post-saline), the inhalation challenge was continued and the next does of LPS was administered. The challenge test was terminated when

any of the following criteria had been met: 1) the subject did not wish to continue for any reason; 2) the subject's FEV<sub>1</sub> decreased 20% or greater from baseline; or 3) a cumulative dose of 41.5 µg had been achieved. Of 84 subjects enrolled in the study, 1 subject withdrew prior to completion of the LPS inhalation challenge test, 52 subjects had at least a 20% decline in the FEV<sub>1</sub> during the LPS inhalation challenge test, and 31 subjects inhaled a cumulative dose of 41.5 µg of LPS and did not decrease their FEV<sub>1</sub> by 20%.

**Assignment of Phenotype.** Study subjects were categorized as either "responsive" or "hypo-responsive" to inhaled LPS. In the course of previous investigations, a large number of study subjects have been exposed to inhaled LPS (Jagiello et al., 1996; Deetz et al., 1997). In general, most healthy non-asthmatic study subjects develop airflow obstruction (FEV<sub>1</sub> ≤ 80% of the pre-exposure value) when challenged with approximately 40 µg of LPS. Based on this and a standard approach to the definition of airway hyperreactivity (FEV<sub>1</sub> decline of 20% from pre-exposure values) (Chai et al., 1975), subjects were categorized as "LPS responsive" if they decreased their FEV<sub>1</sub> by at least 20% at any point during the LPS inhalation challenge, or "hypo-responsive" if they had <20% decline in their FEV<sub>1</sub> after inhaling a cumulative dose of 41.5 µg of LPS.

**Isolation of Genomic Clone.** A human bacterial artificial chromosomal (BAC) library (Research Genetics; Huntsville, Ala.) was screened with two sets of primers (1F: 5'ATGGGGCATATCAGAGCCTA 3; SEQ ID NO:8, IR: 5'GTCCAATGGGGGAAGTTCTCT 3; SEQ ID NO:9, 2F: 5'TCATTGTCCTGCAGAAGGTG 3; SEQ ID NO:10, and 2R: 5'CAGGGCTTTTCTGAGTCGTC 3; SEQ ID NO:11) derived from the human TLR4 gene (Genbank Accession Nos. U88880 and U93091). These sets of primers amplified a 160 bp and 140 bp product, respectively. PCR reactions were prepared by combining the following components: 1 µl of PCR 10× buffer (100 mM Tris-HCl, pH 8.3, 500 mM KCl), 200 µM each of dCTP, dGTP, dATP, and dTTP, 0.25 µM of each primer, 0.2 U of AmpliTaq DNA polymerase (Perkin Elmer; Norwalk, Conn.) and 1 µl of the library sample in a final volume of 10 µl. Thermal cycling was performed with an initial denaturation at 94° C. for 3 minutes followed by 35 cycles comprising 94° C., 55° C., and 72° C. steps of 30 seconds each and a final extension of 72° C. for 5 minutes. PCR products were separated by electrophoresis on 2% agarose gels, stained with ethidium bromide and visualized under UV light.

**Mutation Detection.** Genomic DNA was isolated from whole blood obtained from the study subjects using a rapid salt isolation procedure (Laitinen et al., 1994). Overlapping primer sets were designed across the coding sequence such that products did not exceed 250 bp (Table 1). Primers were derived from flanking intronic sequences to include all splice sites. PCR reactions were prepared as described above except that 10-20 ng of genomic DNA was used as template. Amplification products were separated on non-denaturing, fan-cooled gels containing 5% acrylamide/bis (19:1), 0.5× TBE, and 2.5% glycerol for 3 hours at 20 W. A subset of PCR products were also run on MDE gels. The gels were subjected to silver staining and aberrant bands extracted from the gel, reamplified, and sequenced in both directions. To verify the sequence of the aberrant band, the same primers were used to

amplify and sequence genomic DNA from each subject. At least one individual without the aberrant band was also sequenced for comparison. Amplification products derived from the extracted DNA were purified using spin columns (Qiagen; Santa Clarita, Calif.) and DNA concentration determined by spectrophotometry. The DNA sequence was determined with a Model 377 automated DNA sequencer (Perkin Elmer; Norwalk, Conn.).

TABLE 1

Exon	5' NT <sup>a</sup>	Primers for amplification of TLR4 coding region.		5' NT <sup>a</sup>	SIZE (bp)
		FORWARD PRIMER (5'-3')	REVERSE PRIMER (5'-3')		
1	24	CTGCTCGGTCAAACGG TGAT <sup>1</sup>	+33 AGTCAGTCTCCTCTG AACTTTC <sup>2</sup>	206	
2	-106	CAGCAAGCACGATATT GGAT <sup>3, b</sup>	+18 CGTGCTCAGCATTTC TAACCT <sup>4, b</sup>	244	
3	-53	GAGTTGGGAGACCATG CAGT <sup>5</sup>	+41 GAACACCTCACCTT GTGCAG <sup>6</sup>	262	
4	-52	GGTTCTTATTCAGCAG AAAT <sup>7</sup>	483 CTTGATAGTCCAGA AAAGGCT <sup>8</sup>	172	
4	496	GGTGGCTGTGGAGACA AATC <sup>9</sup>	691 CCGCAAGTCTGTGC AATAAA <sup>10</sup>	195	
4	639	ACTTGGACTTTCCAG CAAC <sup>11</sup>	798 GTCAGTATGAAG CCTAATTTCT <sup>12</sup>	159	
4	751	CTTTATCCAACCAGGT GCA <sup>13</sup>	952 CAAATGCACAGGC CCTCTAGA <sup>14</sup>	201	
4	882	TGGGAGAATTTAGAAA TGAAGGA <sup>15</sup>	1080 AAAGTCTTTTACCC TTTCAATAGTCA <sup>16</sup>	198	
4	1032	TTTCTTCATTTCCCT GGTGA <sup>17</sup>	1161 AGAGATTTGAGTTT CAATGTGGG <sup>18</sup>	129	
4	1129	TGGACAGTTTCCACA TTGA <sup>19</sup>	1309 CTTAGGCTGGTTG TCCCAA <sup>20</sup>	180	
4	1262	TTCAAAGGTTGCTGTT CTCAAA <sup>21</sup>	1426 TGAAACTCACTCA TTTGTTTCAA <sup>22</sup>	164	
4	1349	TCAAACCTCTTGGGCT TAGAACA <sup>23</sup>	1526 CTTGAGACTGGAC AAGCCA <sup>24</sup>	177	
4	1480	CAGAGTTGCTTTCAAT GGCA <sup>25</sup>	1614 GAGAGTCCAGGAA GGTCAA <sup>26</sup>	134	
4	1551	TCCAGGAAAACCTCCT TCCA <sup>27</sup>	1746 ACCTGGAGGGAGTT CAGACA <sup>28</sup>	195	
4	1702	TTCATTGGATACGTTT CC <sup>29</sup>	1905 AAGAGTGCCTCTG GTCCTT <sup>30</sup>	203	
4	1863	ACCAGAGTTTCTGCA ATGG <sup>31</sup>	2032 TACAAGCACACTGA GGACCG <sup>32</sup>	169	
4	1956	TGCCTGTGCTGAGTTT GAAT <sup>33</sup>	2100 TTTATGCAGCCAGC AAGAAG <sup>34</sup>	144	
4	2013	CGGTCTCAGTGTGCT TGTA <sup>35</sup>	2208 GGAGGCACCCCTTC TTCTAA <sup>36</sup>	195	
4	2149	CCAGGATGAGGACTGG GTAA <sup>37</sup>	2338 GCGGCTCTGGATGA AGTGCT <sup>38</sup>	189	
4	2290	AAGCCGAAAGGTGATT GTTG <sup>39</sup>	2442 CTGAGCAGGGTCTT CTCCAC <sup>40</sup>	152	
4	2398	TATCATCTTCATTGTC CTGCAGA <sup>41</sup>	2617 AGATGTTGCTTCCT GCCAAT <sup>42</sup>	219	

TABLE 1-continued

Exon	Primers for amplification of TLR4 coding region.				SIZE (bp)
	5' NT <sup>a</sup>	FORWARD PRIMER (5'-3')	5' NT <sup>a</sup>	REVERSE PRIMER (5'-3')	
4	2406	TCATTGTCCTGCAGAA GGTG <sup>43</sup>	2548	CAGGGCTTTTCTGA GTCGTC <sup>44</sup>	142
4	2529	GACGACTCAGAAAAGC CCTG <sup>45</sup>	2683	TGAACAAGTGTGG ACCCAG <sup>46</sup>	154
4	2597	AATTGGCAGGAAGCAA CATC <sup>47</sup>	2772	GATTAGCAGCCCTG CATATCT <sup>48</sup>	175

<sup>a</sup>relative to published cDNA sequence of Medzhitov et al. (1997), Genbank accession #U93091

<sup>b</sup>relative to published cDNA sequence of Rock et al. (1997), Genbank Accession #U88880, since exon 2 is absent in #U93091.

A "-" indicates primer is within intron on the 5' side of amplified exon; a "+" indicates primer is on the 3' side.

<sup>1</sup>SEQ ID NO:12

<sup>2</sup>SEQ ID NO:36

<sup>3</sup>SEQ ID NO:13

<sup>4</sup>SEQ ID NO:37

<sup>5</sup>SEQ ID NO:14

<sup>6</sup>SEQ ID NO:38

<sup>7</sup>SEQ ID NO:15

<sup>8</sup>SEQ ID NO:39

<sup>9</sup>SEQ ID NO:16

<sup>10</sup>SEQ ID NO:40

<sup>11</sup>SEQ ID NO:17

<sup>12</sup>SEQ ID NO:41

<sup>13</sup>SEQ ID NO:18

<sup>14</sup>SEQ ID NO:42

<sup>15</sup>SEQ ID NO:19

<sup>16</sup>SEQ ID NO:43

<sup>17</sup>SEQ ID NO:20

<sup>18</sup>SEQ ID NO:44

<sup>19</sup>SEQ ID NO:21

<sup>20</sup>SEQ ID NO:45

<sup>21</sup>SEQ ID NO:22

<sup>22</sup>SEQ ID NO:46

<sup>23</sup>SEQ ID NO:23

<sup>24</sup>SEQ ID NO:47

<sup>25</sup>SEQ ID NO:24

<sup>26</sup>SEQ ID NO:48

<sup>27</sup>SEQ ID NO:25

<sup>28</sup>SEQ ID NO:49

<sup>29</sup>SEQ ID NO:26

<sup>30</sup>SEQ ID NO:50

<sup>31</sup>SEQ ID NO:27

<sup>32</sup>SEQ ID NO:51

<sup>33</sup>SEQ ID NO:28

<sup>34</sup>SEQ ID NO:52

<sup>35</sup>SEQ ID NO:29

<sup>36</sup>SEQ ID NO:53

<sup>37</sup>SEQ ID NO:30

<sup>38</sup>SEQ ID NO:54

<sup>39</sup>SEQ ID NO:31

<sup>40</sup>SEQ ID NO:55

<sup>41</sup>SEQ ID NO:32

<sup>42</sup>SEQ ID NO:56

<sup>43</sup>SEQ ID NO:33

<sup>44</sup>SEQ ID NO:57

<sup>45</sup>SEQ ID NO:34

<sup>46</sup>SEQ ID NO:58

<sup>47</sup>SEQ ID NO:35

<sup>48</sup>SEQ ID NO:59

Multi-tissue cDNA Expression Screen. Human adult (Clontech #K1420-1; Palo Alto, Calif.) and fetal (Clontech #K1425-1; Palo Alto, Calif.) multi-tissue cDNA panels were screened by PCR using primers derived from exon 1 (forward; 5'GCTCACAGAAGCAGTGAGGA 3'; SEQ ID NO:60) and exon 4 (reverse; 5'TAGGCTCTGATATGC-

CCCAT3'; SEQ ID NO:61) of the human TLR4 gene. These PCR experiments were performed in 10 µl reactions composed of 0.025 U/µl of DNA polymerase (BioXACT; Bio-Line; Reno, Nev.), 1.0 µM of each primer, 200 µM of each dNTP, 1 µl of 10× buffer supplied by the manufacturer, and 1 µl of the cDNA sample. PCR conditions were: 95° C. for 2 minutes; cycles of 0.5 minutes at 94° C., 0.5 minutes at 55° C., and 1 minute at 68° C.; followed by a final extension step for 10 minutes at 68° C. The amplified products were separated on a 2% agarose gel, stained with ethidium bromide, and visualized under UV light.

Statistical Analysis. The statistical analysis was designed to determine whether specific mutations in the TLR4 gene were associated with the airway hyporesponsiveness to inhaled LPS. A one-tailed test of statistical significance was employed (Fleiss, 1986). In a 2×2 analysis, a Fisher's one-tailed exact test was used to determine whether specific mutations of the TLR4 gene occurred more frequently in study subjects who were considered LPS hyporesponsive compared to those with a normal airway response to inhaled LPS. In addition, because of the nonparametric distribution of dose-response slope (percent decline FEV<sub>1</sub>/cumulative dose of inhaled LPS), this outcome log was transformed and the Student's one-tailed T test (assuming unequal variances) was used to determine whether the dose-response slope was significantly less in subjects with a specific mutation of TLR4 compared to subjects with the common TLR4 allele.

Bioassays. Cells were maintained in F12 media supplemented with 10% fetal calf serum (Gibco, Rockville, Md.), 2 mM L-glutamine, and 10,000 units penicillin/streptomycin, with 1 µg/ml of G418 as selective antibiotic. Cells were transfected with a mixture of 4 µg of DNA and 10 µl of Superfect (Qiagen, Valencia, Calif.) for a 35 mm<sup>2</sup> dish. The DNA mix consisted of 2 µg of DNA and 10 µg NFκB reporter plasmid encoding for the luciferase gene (Clontech, Palo Alto, Calif.) and 1 µg of each TLR4 (Medzhitov et al., 1997; Genbank #U93091) expression plasmid (WT or Asp299Gly). If only one expression plasmid was used, empty vector pcDNA3.1<sup>+</sup> was added to keep the DNA concentration constant. Twenty-four hours later the cells were stimulated with 100 ng/ml of LPS for 6 hours. Total luciferase activity was measured using a commercially available method (Tropix, Bedford, Mass.). Briefly, after rising with PBS, cells were removed from filters by incubation with 120 µl lysis buffer (25 mM Tris-phosphate, pH 7.8; 2 mM DTT; 2 mM 1,2-diaminocyclohexane-N,N,N',N'-tetraacetic acid; 10% glycerol; and 1% Triton X-100) for 15 minutes. Light emission was quantified in a luminometer (Analytical Luminescence Laboratory, San Diego, Calif.).

Airway epithelial cells were obtained from trachea and bronchi of lungs removed for organ donation. Cell were genotyped for TLR4 and isolated by enzyme digestion as previously described (Zabner et al., 1996). Freshly isolated cells were seeded at a density of 5×10<sup>5</sup> cells/cm<sup>2</sup> onto collagen-coated, 0.6 cm<sup>2</sup> diameter millicell polycarbonate filters (Milipore Corp., Bedford, Mass.). The cells were maintained at 37° C. in a humidified atmosphere of 5% CO<sub>2</sub> and air. Twenty-four hours after plating, the mucosal media was removed and the cells were allowed to grow at the air-liquid interface. The culture media consisted of a 1:1 mix of DMEM/Ham's F12, 5% Ultraser G (Biopre SA, Cedex, France), 100 U/ml penicillin, 100 µg/ml streptomycin, 1% nonessential amino acids, and 0.12 U/ml insulin. Epithelia were tested for transepithelial resistance, and for morphology by scanning electron microscopy. Fourteen days after seeding, the basal release of IL-1α was measured in WT/WT (12 specimens from 4 individuals) and WT/Asp299Gly (24 speci-

mens from 4 individuals) epithelia by collecting the basolateral conditioned media after 24 hours (FIG. 6*b*). The epithelia were then exposed to 100 ng/ml of LPS on the apical side for 6 hours, and the basolateral media was collected after 24 hours. IL-1 $\alpha$  was measured using a commercially available ELISA (R&D; Minneapolis, Minn.).

To rescue the LPS hyporesponsive phenotype, heterozygote (WT/Asp299Gly) airway epithelia (FIG. 6*c*) or homozygote (Asp299Gly/Asp299Gly) alveolar macrophages (FIG. 6*d*) were transfected with a recombinant adenovirus vector expressing TLR4 (Rock et al. 1998; Genbank #U88880) that was prepared as described previously (Dandson et al., 1994) by the University of Iowa Gene Transfer Vector Core at titers of about 10<sup>10</sup> infectious units (I.U.)/ml. Briefly TLR4 plasmid was blunt ended and cloned into the shuttle vector pAd5/CMV $\kappa$ -NpA using the EcoRV site. Fourteen days after seeding of the airway epithelia (20 specimens from 4 individuals), 50 MOI of the recombinant viruses (Ad/TLR4 and Ad/eGFP in phosphate buffered saline) were added to the basolateral surface of the epithelia for 30 minutes (Walters et al., 1999). After infection, the epithelia were incubated at 37 $^{\circ}$  C. for an additional 48 hours before the LPS stimulation assay. To assay for basal LPS response, the media was changed with fresh 500  $\mu$ l of Ultrosor G, and collected after 24 hours to measure the basal IL-1 $\alpha$  secretion to the basolateral side. After collecting the basal specimen, the epithelia were exposed to 100 ng/ml of LPS on the apical side for 6 hours, and the media was collected after 24 hours. To assay for gene transfer efficiency, the epithelia was associated with 0.05% trypsin and 0.53 mM EDTA. Fluorescence from 50,000 individual cells was analyzed using fluorescence-activated cell analysis (FACScan, Lysys II software, Becton Dickinson, San Jose, Calif.). The percentage of GFP positive cells ranged between 52% and 76%. Human alveolar macrophages were collected by BAL from a homozygote (Asp299Gly/Asp299Gly) study subject as previously described (Deetz et al., 1967). Macrophages were seeded onto a 96 well plate at a density of 10<sup>5</sup> cells per well. The cells were infected after 4 hours with Ad/TLR4 in a CaPi coprecipitate (Iasbinder et al., 1998) at an MOI of 50, and a C<sup>++</sup> a concentration of 16.8 mM. Sixteen hours after infection, the cells were exposed to LPS in 1% serum for 6 hours, and the media was collected. TNF- $\alpha$  was measured using a commercially available ELISA (R&D; Minneapolis, Minn.).

## Results

**Genomic Structure of the TLR4 Gene.** To determine the genomic structure of the TLR4 gene, a bacterial artificial chromosome (BAC) library was screened by PCR with primers derived from the 5' and 3' ends of the cDNA sequence. Human BAC clone 439F3 was identified with both sets of primers and sequenced to identify exon/intro splice sites using the two published cDNA sequences (Medzhitov et al., 1997; Rock et al., 1998). The TLR4 cDNA published by Medzhitov et al. (1997) (Genbank Accession No. U93091) was missing a 120 bp sequence beginning at nucleotide 136 of the other published cDNA (Rock et al., 1998) (Genbank Accession No. U88880). Assuming this missing sequence represented an alternatively spliced exon, DNA sequencing primers were designed to amplify across the predicted splice junctions. When the genomic sequence (FIG. 9) was compared to the cDNA sequences, the exon/intron junctions were revealed (FIG. 1). The human TLR4 gene is organized into 4 exons that span about 11 kb of genomic DNA. As predicted, the 120 bp sequence missing in the cDNA from Medzhitov et al. (1997) corresponds to exon 2.

To determine whether other exons may be revealed by additional splice forms, a panel of cDNAs from adult (FIG. 2*a*) and fetal (FIG. 2*b*) tissues were examined with PCR primers designed to amplify all nucleotides between exons 1 and 4. In all adult and fetal tissues, three products were amplified (453, 333, and 167 bp), although their relative amounts varied. In addition to the 3' end of exon 1 and the 5' end of exon 4, the 453 bp product contained both exons 2 and 3, the 333 bp product was missing exon 2, and the 167 bp product was missing both exons 2 and 3 (FIG. 2*c*). The 453 and 333 bp products are identical to the published cDNA sequences (Medzhitov et al., 1997; Rock et al., 1998), and confirm that no other exons are present for the human TLR4 gene. The 167 bp product depicts alternative splicing of exons 1 and 4, and represents a novel TLR4 product. The nucleotide position of alleles are relative to the TLR4 cDNA sequence published by Medzhitov et al. (1997).

**Mutation Analysis.** SSCV was employed to detect sequence variants across the entire coding region of TLR4 gene in the 83 unrelated probands who completed the incremental LPS inhalation challenge test. When band shifts were detected on SSCV analysis, the bands were sequenced to identify and confirm the sequence variants. The SSCV and sequence analysis were performed blinded to the LPS response phenotype of the study subjects. A band variant was detected by SSCV in 10 (12%) of the 83 subjects, and direct sequencing detected an A to G substitution at nucleotide 896 of the published TLR4 cDNA (FIG. 3*a*) (Genbank Accession No. U88880). To confirm these findings, the 83 unrelated probands were sequenced in the forward and reverse directions with primers designed to amplify the 896 nucleotide; the same 10 individuals were found to have the A896G substitution and the remaining individuals were confirmed to have the common TLR4 allele. Importantly, one of the 10 individuals with the A896G substitution was homozygous for this mutation and the remaining 9 had a single mutant allele. The allelic frequency of the A896G substitution was 6.6% in the study population, 7.9% in a normal control population from Iowa (Lidral et al., 1998), and 3.3% in the parental chromosomes of the CEPH population (NIH-CEPH, 1992).

The A896G substitution results in replacement of a conserved aspartic acid (A) residue with glycine (G) at amino acid 299 (FIG. 4). This missense mutation (Asp299Gly) is in the fourth exon of TLR4 and is present in the extracellular domain of this receptor. The region surrounding amino acid 299 appears to be in an alpha helical conformation (Gibrat et al., 1987). Replacement of the conserved aspartic acid with glycine at position 299 causes disruption of the alpha helical protein structure resulting in the formation of an extended beta strand (Gibrat et al., 1987).

Two other SSCV variants were identified in a single proband; a T to C change at nucleotide 479 (FIG. 3*b*) and a deletion of a thymine nucleotide at position -11 in the third intron. The T479C variant did not alter the amino acid composition and is therefore considered a silent mutation. The T-11 variant did not alter the acceptor splice site, so the significance of this deletion is unknown. The subject with these two sequence variants was responsive to inhaled LPS.

An additional missense mutation was identified within exon 4 of the TLR4 gene. This mutation results in the replacement of a threonine with an isoleucine residue at amino acid 399, which is on the 3' side of the previously identified Asp299Gly mutation. Both mutations are present within the extracellular domain of the TLR4 protein. When the genotypes were compared with the phenotypes associated with endotoxin responsiveness, one individual who carried the Asp299Gly mutation did not carry the Thr399Ile mutation.

Likewise, one individual who had the Thr399Ile mutation did not have the Asp299Gly mutation. Nine of the 83 volunteers carry both mutations. There was a strong correlation between both of the TLR4 mutations and endotoxin hyporesponsiveness.

**Phenotype/Genotype Analysis.** Of the 83 unrelated study subjects who completed the LPS inhalation challenge test, 52 (63%) were responsive to inhaled LPS and 31 (37%) were hyporesponsive to inhaled LPS. When the genotypes of these individuals were examined, the Asp299Gly sequence variant occurred in 3 LPS responsive (5.8%) and 7 LPS hyporesponsive (22.6%) study subjects ( $P=0.029$ ). Among the subjects with the common TLR4 allele ( $N=73$ ), the dose-response slope (percent decline  $FEV_1$ /cumulative dose of inhaled LPS) averaged 1.86% decline in  $FEV_1/\mu\text{g}$  inhaled LPS (range 0.01%-19.78%), while the dose-response slope for the subjects with the Asp299Gly allele ( $N=10$ ) was significantly less ( $P=0.037$ ), averaging 0.59% decline in  $FEV_1/\mu\text{g}$  inhaled LPS (range 0.00%-1.59%) (FIG. 5). This group difference in the dose-response slope also exists on the log scale ( $P=0.026$ ) where  $\log(1+\Delta FEV_1)$  was used since one individual has a  $\Delta FEV_1$  of 0.0. The subject who was homozygous for the Asp299Gly allele was hyporesponsive to inhaled LPS with a 0.28% decline in  $FEV_1/\mu\text{g}$  of inhaled LPS. This homozygote Asp299Gly subject is one of a monozygote twin pair; her twin sister was subsequently phenotyped and was also found to be hyporesponsive to inhaled LPS with a 0.34% decline in  $FEV_1/\mu\text{g}$  of inhaled LPS.

The biological significance of the Asp299Gly mutation was evaluated in several ways. First, transfection of CHO cells with either the WT or the mutant TLR4 gene demonstrated that the mutant allele does not respond normally to LPS stimulation (FIG. 6a). Second, airway epithelia obtained from heterozygote (WT/Asp299Gly) individuals do not respond to LPS stimulation (FIG. 6b). Third, the wild-type allele of TLR4 clearly restored LPS responsiveness in either airway epithelial cells (FIG. 6c) or alveolar macrophages (FIG. 6d) obtained from individuals with the TLR4 mutation.

#### Discussion

The results described herein provide the first direct evidence to indicate that a sequence polymorphism in the TLR4 gene is associated with a hyporesponsive LPS phenotype in humans that interrupts LPS signaling. This conclusion is supported by the following findings: 1) unrelated subjects with the Asp299Gly substitution were significantly less responsive to inhaled LPS than those homozygous for the common TLR4 allele; 2) an individual who was homozygous for the Asp299Gly substitution was hyporesponsive to inhaled LPS; 3) a monozygous twin pair, homozygous for the Asp299Gly substitution, demonstrated a very similar response to inhaled LPS; and 4) in vitro studies demonstrate that mutant TLR4 does not respond to LPS stimulation. This conclusion is indirectly supported by the finding in C3H/HeJ mice where a point mutation in intracellular domain of Tlr4 is also associated with LPS hyporesponsiveness (Poltorak et al., 1998; Qureshi et al., 1999). The identified Asp299Gly missense mutation is in the fourth exon of the TLR4 gene and, unlike the C3H/HeJ mutation, is located in the extracellular domain of this receptor. The findings described herein suggest that a specific region in the extracellular domain of TLR4 may play an important role in receptor function and regulation of the innate immune response to LPS in humans.

This discovery may have important ramifications for a broad spectrum of human diseases. First, the Asp299Gly TLR4 substitution is a relatively common mutation, potentially affecting the responsiveness of a substantial portion of

the population to LPS. The allelic frequency of the Asp299Gly TLR4 substitution in the three populations screened was between 3.3% and 7.9%, and homozygous individuals were identified in each of the populations. Second, the results described herein provide further support for the role of TLR4 in LPS signaling. A fundamental understanding of LPS signaling will undoubtedly alter approaches to gram-negative sepsis, as well as other diseases thought to be mediated by endotoxin, such as the systemic inflammatory response syndrome (Wang et al., 1995), acute respiratory distress syndrome (Brighan et al., 1986), and asthma or other forms of airway disease caused (Rylander et al., 1989; Schwartz et al., 1995) or exacerbated (Schwartz et al., 1995; Michel et al., 1996) by endotoxin. Identification of the essential components of LPS signaling also provide new therapeutic targets for endotoxin mediated conditions. Third, the specific Asp299Gly TLR4 substitution may provide a simple screening mechanism to risk stratify a population. Identification of this mutation may lead to a better appreciation of the role of LPS responsiveness in a broad range of acquired and genetic disorders. For example, mutations of the TLR4 gene may explain why a minority of patients with gram-negative sepsis develop ARDS (Hudson et al., 1995) or why patients with cystic fibrosis and the identical CFTR mutation have different courses of disease (Veeze et al., 1994). Finally, it must be acknowledged that normal responsiveness to endotoxin is an important component of innate immunity, and the Asp299Gly TLR4 substitution may prove to be associated with negative clinical outcomes. In fact, the C3H/HeJ mouse is more susceptible to *Salmonella typhimurium* (O'Brien et al., 1980), despite its resistance to LPS (Sultzter et al., 1968). Although individuals with the Asp299Gly TLR4 substitution may be more resistant to localized forms of endotoxin-induced inflammation, these individuals may prove to be more susceptible to a systemic inflammatory response initiated or exacerbated by endotoxin.

The findings described herein demonstrate that a specific region in the extracellular domain of TLR4 plays an important role in receptor function and regulation of the innate immune response to LPS in humans. The sequence variants reported thus far for mammalian, i.e., murine, TLR4 are located in the cytoplasmic domain (Poltorak et al., 1998; Qureshi et al., 1999). While it is logical to hypothesize that mutations in the cytoplasmic domain of TLR4 disrupt the signaling pathway leading to activation of NF- $\kappa$ B and AP-1 (Medhitov et al., 1998), the potential mechanisms that are altered by mutations in extracellular domain of TLR4 are less obvious.

TLR4 is a transmembrane protein and it is therefore not surprising to find a putative signal peptide at the N-terminus, presumably responsible for proper trafficking to the cell membrane (Medzhitov et al., 1997). This finding raises the possibility that sequence variants in the extracellular domain of TLR4 can disrupt trafficking of this receptor to the cell membrane and enhance proteolysis. There are several examples in the literature which show that the presence of a single point mutation at a conserved amino acid can disrupt protein folding and specifically affect protein trafficking (Jackson et al., 1998; DeFranco et al., 1998). Mutations of this kind could give rise to a range of phenotypes from almost normal to completely unresponsive, depending on the severity of the trafficking defect. Another possible mechanism that may explain the association between a mutation in the extracellular domain of TLR4 and LPS responsiveness is impairment of ligand binding to the extracellular domain. The extracellular domains of the toll proteins are clearly involved in the cellular response to LPS (Medzhitov et al., 1997; Yang et al.,

1998). Yet, so far no convincing evidence has been found that the toll proteins are directly interacting with LPS (Wright, 1999). The discovery of an association between a mutation in the extracellular domain of TLR4 and a discernable LPS phenotype in humans should aid in clarifying whether the extracellular domain of this receptor is important for proper membrane localization of TLR4 and if LPS directly interacts with TLR4. A final possibility is that the missense mutation of TLR4 affects its interaction with either an extracellular ligand or an intracellular protein (Wright, 1999). The extracellular domain of TLR4, and specifically the Asp299Gly amino acid change may provide a key therapeutic target to modulate LPS signaling.

TLR4 may be one of several factors that may regulate the airway response to inhaled LPS. Among the subjects with the Asp299Gly substitution, 7 were hyporesponsive to LPS and 3 developed airflow obstruction during the LPS inhalation challenge test. None of the subjects with this mutation were in the most sensitive quartile of the LPS responders. Although asthmatic patients were included because of their increased airway responsiveness to inhaled LPS (Michel et al., 1989), it remains possible that the 3 subjects with the Asp299Gly substitution who were responsive to inhaled LPS had some type of airway inflammation (e.g., early viral illness) that enhanced their response to inhaled LPS. Moreover, since 22.6% of the study subjects who were hyporesponsive to LPS had at least one copy of the Asp299Gly TLR4 allele, other genes (or possibly acquired conditions) may prove to play a role in modulating the biological response to LPS. For instance, TNF- $\alpha$  is one of the primary cytokines that mediates the toxic effects of LPS (Beutler et al., 1985). A polymorphism at -308 in the TNF- $\alpha$  promoter region results in higher constitutive and inducible levels of TNF- $\alpha$  (Wilson et al., 1997), and this polymorphism has been shown to result in a worse outcome in children with either meningococcal disease (Nadel et al., 1996) or cerebral malaria (McGuire et al., 1994). Likewise, allelic variants of TLR4 receptors and their ligands appear to delay the progression of disease in patients with HIV infection (Mummidi et al., 1995). Thus, TLR4 appears to represent only one of a number of genetic variants that may modulate the pathophysiological response to LPS

#### EXAMPLE 2

##### Methods

To genotype patients for TLR4, a PCR based RFLP assay was employed. For example, to detect point mutations in codon 299 such as those that result in an amino acid substitutions at amino acid 299 (Asp (GAT) to Gly (GGT) change) and/or in codon 399 (Thr (ACC) to Ile (ATC) change), a primer spanning that codon and a second primer 3' or 5' to that primer are prepared. Primers useful to detect a nucleotide change at codon 299 are: 299Forward primer (5' GATTAG-CATACTTAGACTACTACCTCCATG 3'; SEQ ID NO:66) and 299Reverse primer (5' GATCAACTTCTGAAAAAG-CATTCCCAC; SEQ ID NO:67). The underlined base in the forward primer fortuitously creates a NcoI (CCATGG) restriction site in the context of a mutation which is specific for carriers of the 299 mutation Asp to Gly. Primers useful to detect a nucleotide change at codon 399 include: 399Forward primer (5' GGTTGCTGTTCTCAAAGTGATTTGGG AGA A; SEQ ID NO:68) and 399Reverse Primer (5' ACCT-GAAGACTGGAGAGTGAGTTAAATGCT; SEQ ID NO:69). The underlined base in the 399 Forward primer fortuitously creates a HinI restriction site (GANTC) in the context of a mutation which is specific for carriers of the 399

mutation Thr to Ile. The reverse primers were chosen at random based solely on such parameters as annealing temperature, expected product size and the like. Additional residues or fewer residues at the 5' end of the primers having the nucleotide change will not affect the outcome as long as the amplification conditions and reverse primers are chosen accordingly.

A MJ Tetrad PTC-225 Thermo cycler was used with both primer sets. For the 299 assay, primers at 10-100 pmole were mixed with 20 ng of genomic DNA, and Perkin Elmer Taq polymerase, deoxyribonucleotides and buffer according to the manufacturer's directions. Cycling conditions included an initial denaturation of 95° C., for 3 to 4 minutes, then 30 cycles of denaturation at 95° C. for 30 seconds; annealing at 55° C. for 30 seconds; and extension at 72° C. for 30 seconds.

For the 399 assay, a Clontech Advantage amplification kit was used according to the manufacturer's directions. The cycling conditions included an initial denaturation at 95° C. for 3 to 4 minutes, then 30 cycles of denaturation at 95° C. for 30 seconds; annealing at 55° C. for 30 seconds; and extension at 72° C. for 30 seconds.

##### Results

Septic shock remains a significant health concern worldwide and despite progress in understanding the physiologic and molecular basis of septic shock, the high mortality rate of septic shock patients remains unchanged. A common polymorphism in TLR4 was identified that is associated with hyporesponsiveness to inhaled endotoxin or lipopolysaccharide (LPS) in humans (Example 1). Since TLR4 is a major receptor for LPS in mammals and Gram-negative bacteria are the prevalent pathogen associated with septic shock, these specific TLR4 alleles may be associated with a predisposition to or a more severe disease outcome for septic shock patients. Ninety-one septic shock patients as well as 83 healthy blood donor controls were genotyped for the presence of the TLR4 Asp299Gly and TLR4 Thr399Ile mutations. The TLR4 Asp299Gly allele was found exclusively in septic shock patients (P=0.05). Furthermore, septic shock patients with the TLR4 Asp299Gly allele had a significantly higher prevalence of Gram-negative infections. These findings suggest that the TLR4 Asp299Gly allele predisposes people to develop septic shock with Gram-negative organisms.

In another study in a German population, the following observations were made: patients with TLR4 mutation(s) tended to have a longer ICU stay; patients with TLR4 mutation(s) were more prone to sepsis; and sepsis in patients with TLR4 mutation(s) was more severe than in patients that were wild-type for TLR4.

In a third study in a Finnish population, it was observed that mothers of pre-term infants and pre-term infants had a higher frequency of TLR4 mutation(s) than term infants. Also, infants with respiratory distress syndrome (RDS), particularly those born very prematurely, tended to have a higher frequency of mutation(s) than premature infants without respiratory distress syndrome.

Thus, generally, the 299 mutation is slightly more prevalent than the 399 mutation, and in 90% of the cases, both mutations were observed. When only one mutation was found, it was more frequently the 299 mutation.

Thus, the presence of TLR4 mutations is associated with predisposition to septic shock, severity of sepsis, pre-term delivery, and RDS in pre-term infants.

##### REFERENCES

- Beutler et al., *Science*, 229:869 (1985).  
Bonner et al., *Am. J. Respir. Cell Mol. Biol.*, 19:672 (1998).

Brigham et al., *Am. Rev. Respir. Dis.* 133:913 (1986).  
 Chai et al., *J. Allergy Clin. Immunol.*, 56:323 (1975).  
 Chow et al., *J. Biol. Chem.*, 274:10689 (1999).  
 Coutinho et al., *Immunogenetics*, 7:17 (1978).  
 Davidson et al., *Exp. Neurol.*, 125:258 (1994).  
 Deetz et al., *Am. J. Respir. Crit. Care Med.*, 155:254 (1997).  
 DeFranco et al., *Prog. Clin. Biol. Res.*, 397:119 (1998).  
 De Franco et al., *J. Steroid Biochem. Mol. Biol.*, 65:51 (1998).  
 Fasbender et al., *J. Clin. Invest.*, 102:184 (1998).  
 Favorite et al., *J. Clin. Invest.*, 21:589 (1942).  
 Fisher et al., *Wiley*, New York (1993).  
 Fleiss, J. C. The design and analysis of clinical experiments. New York: *Wiley*, 1986.  
 Gibrat et al., *J. Mol. Biol.*, 198:425 (1987).  
 Hoshino et al., *J. Immunol.*, 162:3749 (1999).  
 Hudson et al., *Am. J. Respir. Crit. Care Med.*, 151:293 (1995).  
 Jackson et al., *Thromb. Haemost.*, 80:42 (1998).  
 Jagielo et al., *Chest*, 110:263 (1996).  
 Kuhns et al., *J. Immunol.*, 158:3959 (1997).  
 Laitinen et al., *Biotechniques*, 17:316 (1994).  
 Lidral et al., *Am. J. Hum. Genet.*, 63:557 (1998).  
 McGuire et al., *Nature*, 371:508 (1994).  
 Medzhitov et al., *Cell*, 91:295 (1997).  
 Medzhitov et al., *Nature*, 388:394 (1997).  
 Medzhitov et al., *Mol. Cell.*, 2:253 (1998).  
 Michel et al., *J. Appl. Physiol.*, 66:1059 (1989).  
 Michel et al., *Rev. Respir. Dis.*, 146:352 (1992).  
 Michel et al., *Am. J. Respir. Crit. Care Med.*, 154:1641 (1996).  
 Mummidi et al., *Nat. Med.*, 4:786 (1998).  
 Nadel et al., *J. Infect. Dis.*, 174:878 (1996).

NIH-CEPH (Collaborative Mapping Group), *Science* 258:67 (1992).  
 O'Brien et al., *Nature*, 287:440 (1980).  
 Poltorak et al., *Blood Cells Mol. Dis.*, 24:340 (1998a).  
 Poltorak et al., *Science*, 282:2085 (1998).  
 Qureshi et al., *J. Exp. Med.*, 189:615 (1999).  
 Rock et al., *Proc. Natl. Acad. Sci. USA*, 95:588 (1998).  
 Rylander et al., *Am. Rev. Respir. Dis.*, 140:981 (1989).  
 Santamaria et al., *J. Immunol.*, 143:913 (1989).  
 Schwartz et al., *Am. J. Respir. Crit. Care Med.*, 151:47 (1995a).  
 Schwartz et al., *Am. J. Respir. Crit. Care Med.*, 152:603 (1995b).  
 Sigmund et al., *Hypertension*, 22:599 (1993).  
 Sultzter, *Nature*, 219:1253 (1968).  
 Vogel et al., *J. Immunol.*, 122:619 (1979).  
 Veeze et al., *J. Clin. Invest.*, 93:461 (1994).  
 Walters et al., *J. Biol. Chem.*, 274:10219 (1999).  
 Wang et al., *Clin. Nucl. Med.*, 20:494 (1995).  
 Wilson et al., *Proc. Natl. Acad. Sci. USA*, 94:3195 (1997).  
 Wright et al., *Science*, 249:1431 (1990).  
 Wright et al., *J. Exp. Med.*, 189:605 (1999).  
 Yang et al., *Nature*, 395:284 (1998).  
 Zabner et al., *J. Virol.*, 70:6694 (1996).  
 All publications, patents and patent applications are incorporated herein by reference. While in the foregoing specification, this invention has been described in relation to certain preferred embodiments thereof, and many details have been set forth for purposes of illustration, it will be apparent to those skilled in the art that the invention is susceptible to additional embodiments and that certain of the details herein may be varied considerably without departing from the basic principles of the invention.

## SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 72

<210> SEQ ID NO 1

<211> LENGTH: 28

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

gcgtggaggt atgtggctgg agtcagct

28

<210> SEQ ID NO 2

<211> LENGTH: 28

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

tcacggaggt tagaatgctg agcacgta

28

<210> SEQ ID NO 3

<211> LENGTH: 28

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 3

ttatccaggt aatgaatcca cttttaca

28

-continued

---

<210> SEQ ID NO 4  
 <211> LENGTH: 21  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 4

Leu Ala Tyr Leu Asp Tyr Tyr Leu Asp Asp Ile Ile Asp Leu Phe Asn  
 1 5 10 15

Cys Leu Thr Asn Val  
 20

<210> SEQ ID NO 5  
 <211> LENGTH: 20  
 <212> TYPE: PRT  
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 5

Leu Thr Tyr Thr Asn Asp Phe Ser Asp Asp Ile Val Lys Phe His Cys  
 1 5 10 15

Leu Ala Asn Val  
 20

<210> SEQ ID NO 6  
 <211> LENGTH: 20  
 <212> TYPE: PRT  
 <213> ORGANISM: Rattus norvegicus

<400> SEQUENCE: 6

Leu Thr Tyr Ile Asn His Phe Ser Asp Asp Ile Tyr Asn Leu Asn Cys  
 1 5 10 15

Leu Ala Asn Ile  
 20

<210> SEQ ID NO 7  
 <211> LENGTH: 20  
 <212> TYPE: PRT  
 <213> ORGANISM: Cricetulus griseus

<400> SEQUENCE: 7

Phe Thr Tyr Ala Asn Glu Phe Ser Glu Asp Ile Thr Asp Phe Asp Cys  
 1 5 10 15

Leu Ala Asn Val  
 20

<210> SEQ ID NO 8  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 8

atggggcata tcagagccta 20

<210> SEQ ID NO 9  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 9

gtccaatggg gaagttctct 20

<210> SEQ ID NO 10  
 <211> LENGTH: 20  
 <212> TYPE: DNA



-continued

---

```

<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 10
tcattgtcct gcagaagggtg                20

<210> SEQ ID NO 11
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 11
cagggccttt ctgagtcgtc                20

<210> SEQ ID NO 12
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 12
ctgctcggtc aaacggtgat                20

<210> SEQ ID NO 13
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 13
cagcaagcac gatattggat                20

<210> SEQ ID NO 14
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 14
gagttgggag accatgcagt                20

<210> SEQ ID NO 15
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 15
ggttcttatt cagcagaaat                20

<210> SEQ ID NO 16
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 16
ggtggctgty gagacaaatc                20

<210> SEQ ID NO 17
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 17
acttgacct ttccagcaac                20

<210> SEQ ID NO 18
<211> LENGTH: 19

```

-continued

---

```

<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 18
ctttatccaa ccaggtgca 19

<210> SEQ ID NO 19
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 19
tgggagaatt tagaaatgaa gga 23

<210> SEQ ID NO 20
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 20
tttcttcatt ttccctggtg a 21

<210> SEQ ID NO 21
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 21
tggacagttt cccacattga 20

<210> SEQ ID NO 22
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 22
ttcaaagggt gctgttctca aa 22

<210> SEQ ID NO 23
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 23
tcaaacttct tgggcttaga aca 23

<210> SEQ ID NO 24
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 24
cagagttgct ttcaatggca 20

<210> SEQ ID NO 25
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 25
tccagaaaa cttccttcca 20

<210> SEQ ID NO 26

```

-continued

---

```

<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 26

ttcattggat acgtttcc 18

<210> SEQ ID NO 27
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 27

accagagttt cctgcaatgg 20

<210> SEQ ID NO 28
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 28

tgccgtgtgct gagtttgaat 20

<210> SEQ ID NO 29
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 29

cggtcctcag tgtgcttgta 20

<210> SEQ ID NO 30
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 30

ccaggatgag gactgggtaa 20

<210> SEQ ID NO 31
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 31

aagccgaaag gtgattgttg 20

<210> SEQ ID NO 32
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 32

tatcatcttc attgtcctgc aga 23

<210> SEQ ID NO 33
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 33

tcattgtcct gcagaaggtg 20

```

-continued

---

```

<210> SEQ ID NO 34
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 34

gacgactcag aaaagccctg                20

<210> SEQ ID NO 35
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 35

aattggcagg aagcaacatc                20

<210> SEQ ID NO 36
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 36

agtcagctcc tctgaacttt c            21

<210> SEQ ID NO 37
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 37

cgtgctcagc attctaacct                20

<210> SEQ ID NO 38
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 38

gaacacctca cttgtgcag                20

<210> SEQ ID NO 39
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 39

cttgatagtc cagaaaaggc t            21

<210> SEQ ID NO 40
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 40

ccgcaagtct gtgcaataaa                20

<210> SEQ ID NO 41
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 41

gtcagcttat gaagcctaat ttct        24

```

---

-continued

---

<210> SEQ ID NO 42  
<211> LENGTH: 22  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens  
  
<400> SEQUENCE: 42  
caaattgcac aggccctcta ga 22

<210> SEQ ID NO 43  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens  
  
<400> SEQUENCE: 43  
aaagtctttt accctttcaa tagtca 26

<210> SEQ ID NO 44  
<211> LENGTH: 23  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens  
  
<400> SEQUENCE: 44  
agagattga gtttcaatgt ggg 23

<210> SEQ ID NO 45  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens  
  
<400> SEQUENCE: 45  
ctttaggctg gttgccc aa 20

<210> SEQ ID NO 46  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens  
  
<400> SEQUENCE: 46  
tgaaaactca ctcatttggt tcaa 24

<210> SEQ ID NO 47  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens  
  
<400> SEQUENCE: 47  
cttcgagact ggacaagcca 20

<210> SEQ ID NO 48  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens  
  
<400> SEQUENCE: 48  
gagaggtcca ggaaggtcaa 20

<210> SEQ ID NO 49  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens  
  
<400> SEQUENCE: 49  
acctggaggg agttcagaca 20

---

-continued

---

<210> SEQ ID NO 50  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens  
  
<400> SEQUENCE: 50  
aagagctgcc tctggtcctt 20

<210> SEQ ID NO 51  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens  
  
<400> SEQUENCE: 51  
tacaagcaca ctgaggaccg 20

<210> SEQ ID NO 52  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens  
  
<400> SEQUENCE: 52  
tttatgcagc cagcaagaag 20

<210> SEQ ID NO 53  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens  
  
<400> SEQUENCE: 53  
ggaggcacc cttcttctaa 20

<210> SEQ ID NO 54  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens  
  
<400> SEQUENCE: 54  
geggctctgg atgaagtgct 20

<210> SEQ ID NO 55  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens  
  
<400> SEQUENCE: 55  
ctgagcaggg tcttctccac 20

<210> SEQ ID NO 56  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens  
  
<400> SEQUENCE: 56  
agatgttct tctgccaat 20

<210> SEQ ID NO 57  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens  
  
<400> SEQUENCE: 57

-continued

---

```

cagggtttt ctgagtcgtc 20

<210> SEQ ID NO 58
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 58

tgaacaagtg ttggaccag 20

<210> SEQ ID NO 59
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 59

gattagcagc cctgcatatc t 21

<210> SEQ ID NO 60
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 60

gctcacagaa gcagtgagga 20

<210> SEQ ID NO 61
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 61

taggctctga tatgccccat 20

<210> SEQ ID NO 62
<211> LENGTH: 10665
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)...(10665)
<223> OTHER INFORMATION: n = A,T,C or G

<400> SEQUENCE: 62

aaaaactcc cttgcctcaa aaactgctcg gtcaaacggt gatagcaaac cacgcattca 60
cagggccact gctgctcaca aaaccagtga ggatgatgcc aggatgatgt ctgcctcgcg 120
cctggctggg actctgatcc cagccatggc cttcctctcc tgcgtgagac cagaaagctg 180
ggagccctgc gtggaggtat gtggctggag tcagctctc tgaactttcc ctcacttctg 240
cccagaactt ctcaactgtgt gccctggttt gtttatTTTT gcaaaaaaaaa aaagagttaa 300
attaccttaa agactcaaga agccacagag atcaaataat tcattgttac agggcactag 360
aggcagccat tgggggtttg ttccatttgg aaattttgag tgctaacagg ggcagtagat 420
aacatagatc tgcttaaggt ccctgctctg ctaccttctg gctctgtgaa gaaattatca 480
aacctgtctg agactagttt tgcactctgt aagagaatta taataccttc ttcactagag 540
agtaagcaga ctgcttcagt gtcatttctt cccactggtg gtctttacac tcagcttcaa 600
gcagtcaccc tgctccttcc aatctcagga aaaagatggc tttgtgtgtg tgtctctaga 660
gaaagaactt tctaagttgg tgcagacttc tgtatgcagt aatatagttt agtccagagg 720
atgaaaaaaaa taagagaatg aaaaaggaaa agagagagag agagaagaaa aaagcaagag 780

```

-continued

---

ggaaatagt	ataatgtcag	ctaattgcaac	agtttctttc	ttagtgaaat	accaatcagc	840
tggttggtaa	tcttattcat	gatggatctc	ttttgttttt	cccctgcgca	gacttcacag	900
ttgctttaga	aaccatagc	agagccgaac	agctaagaaa	atgatttaca	gtgaggcagg	960
gtcagaaact	caagagagaa	aaagccagct	gcagtcctga	agttgaggat	ataggagaaa	1020
atcaagtaat	attagcaaaa	gactaattca	ttatcttgaa	gccatccctt	ccctcaattc	1080
cctgcccata	gtcctcctcc	ttgtcctctt	ctctgnatcc	ctctgctggt	aggttaatgg	1140
agatagattt	tctaattang	ctcactgcga	gataaaacc	agcccatggt	tctattagnc	1200
aatattgtct	ttgaggtctc	atggcttgca	ncatttaagc	agacatacga	atgaagatct	1260
gcatgtttga	actctgactt	tgccgatatt	acttcatttc	tttgaatttc	cattttctct	1320
atctttaa	gcttatttga	agattaagtg	aaagtatata	acaacaaga	actatgcagg	1380
cgtatggtaa	gggattaatg	atagatgata	ataattaatg	ttgacatcta	ttgatcactt	1440
atactgtagc	gggcttttaa	ataaactctt	taaacacctt	atctcattta	atccttcaaa	1500
cattctattg	gtttcaaa	acagaaaact	acaattagct	ggcttctgca	aggaattttg	1560
ttggaggaaa	tgagagcatt	cagaaattag	atgggagcgt	tagagaatta	ggcttcaaaa	1620
gaatgtggga	aagtaggcta	gaaagcagtg	taaaaacaaa	gacagcataa	agcacttgac	1680
cttatttact	aggttccacc	atgggaatcc	atgcactcta	aagatttccc	cctatttcta	1740
catcactttg	ctcaagggtc	aatgagccaa	ggaaaagaat	gcagttgtca	aatctggggc	1800
catgactaag	gaaggtctgg	acatcttgac	tgccagacag	tctcccaat	gatatggagt	1860
attdagaatg	atactggata	ttttatttat	ttttgtatt	ttcaactttt	aagttcagag	1920
gcacatgtgc	agagcatgca	ggtttattac	ataagtaaat	gtgtccatg	gtgatttgct	1980
gcatagatca	tgaaaatag	gaacgcata	tggatttggt	tgatcctt	gtgcaggggc	2040
catgctcacc	ttctctgtat	ccttccaatt	ttagtatatg	tgctactgca	gcaagcacga	2100
tattggatat	tttattacct	acattttaca	tatgataaaa	tgaggctcac	tgaggttttt	2160
ctttgttctg	ttttattttg	ttttgttttt	aaagacttgg	ccctaaacca	cacagaagag	2220
ctggcatgaa	accagagact	ttcagactcc	ggagcctcag	cccttcccc	cgattccatt	2280
gcttcttctg	aaatgctgcc	gttttatcnc	ggaggttaga	atgctgagca	cgtagtaggt	2340
gctctttact	ttctaactca	gagtaagaca	attdataagc	atgaattgag	tgaatggatg	2400
gatggatata	tggatggaag	gatggacaga	tggatgaaag	gttgactgaa	ttttgtgctt	2460
gcacaaaaag	aggccccctc	ccaccatctc	tggtctagga	gaggggaggt	gggagaccat	2520
gcagtaaaga	tacttcatgt	catgtgtaat	cattgcaggt	ggtoctaata	attacttatc	2580
aatgcatgga	gctgaatttc	tacaaaatcc	cgcacaacct	ccccttctca	accaagaacc	2640
tggacctgag	ctttaatccc	ctgaggcatt	taggcagcta	tagcttcttc	agtttcccag	2700
aactgcaggt	gctggattta	tccaggtaat	gaatccactt	ttacatactg	cacaagggtga	2760
ggtgttcatt	gtcctatcat	ttcattattg	gactggaaag	cttggtttgt	ggagtctcat	2820
cttcattcac	ttattcattc	atacaacaga	tgtcttatta	actatataac	cttgagcaag	2880
ctacctctat	tctccaggtc	tcagttttct	aatctgtgaa	gtaggcaggt	ggctgagaca	2940
gcttctaagg	gcaattctaa	ttttagggtt	tcttttaaga	caggagagaa	aattagetta	3000
aattctttca	taagcagcta	tttattgact	acttgctata	tgttgtaaac	tctgcaagaa	3060
gacaggcata	tattgatata	taacacacag	cccctgttgt	taaggaggca	tatcttcttg	3120



-continued

---

aaagagtaa taccttaaag tccctgggtat ggtcctgggt acatagtata tagtcaacac	3180
atTTtaatta tgattttttg gatctggaaa ctgatataaa gatagcgaca tataacagta	3240
ggTgataaat tatgtttaaa ctaaaggtaa ctaattgtat ttttcagaag aggggccttc	3300
tctgtggTgg gtagtcaaga aagattcatg aactgcataa gattcaaaca atgtctagaa	3360
tattaaaact agtggTggca ggtgaaatgt catcttgata ttttagggga accaaaattct	3420
aaaagggttt tcatcatcgg ggccttattt gcaaatcgaa ctagataatg gatcatgttc	3480
tctgcaatgg tttgtaaaac atttcaaaac attttacata ttttttatta tagaaattat	3540
tgataaagac taaggtcaca gtataaaaat cctttttaga gcagacattt ctgtagaaga	3600
gtgaacatat gacctattat actctaattt ggatatagat aggatgtaac aaaggagtaa	3660
tgTggaacaat tcaaaggcag TggTatagTg catanagtcc TgtTggggTc agaagacctg	3720
agcccaagtt taccoccaac atttataacc catgtaacct tagcatatta ctTcatctcc	3780
cttaatcctt agtttcatat ctgatcaatg gaaatgatga aacttattct gctggattaa	3840
atgtgataat aaatattaat atgctgtata tatttaaatt tttataaaat atattttata	3900
agcataaagt attcttacag aatttcatta ggtttttaaa ataatttcaa cttttatttt	3960
Tgattcaggg atttcatagg ttatattgTg taatgctgag gtgtagggta caatcgatac	4020
catcactcag gtagtgagca tagtaccocaa tagttagttt tTcaaccctt gctgctttct	4080
ctctatcccc tctctagtaa tccccagggt ctatttttTg catctttatg tccatgtgta	4140
ctccatgttt ggatcctact tataaagtga gaactcatgg tattTggctt tctgTncctt	4200
TgtTngctaa tttgcttagg ataatggcta ctagctgcat ctatgccatt atgttctaaa	4260
ttTcantTnc ctgcatgaaa attttgtcaa gtactctatt aaggtagacc acctctccct	4320
ttttttttca aacaagaagt agnttttccc aaacaatgcc cttatggaat TnatctTca	4380
tccnnggata cccaataact Tgccccaaan ccttaatctg nctTcagag aggccacctt	4440
ccttctgtaa cccataggag attTggattg gtaagaatgc tttgtgatag cccagcagcc	4500
ttctttcccc tatagaaata tatatatant ctttttatag gtgaggaaact gaagctTgaa	4560
taattTaaat gacttatata catnatcatt gctTgttagc cacagaccag agattTaaT	4620
Tncatctcc agaatccaac TtaaagtTt tctTgtctt aactctctac ttctTaaag	4680
Tgattatcac caatgtaatg atatagagnc acagcaagac cctttccttc TcaocTaatg	4740
tatagagcaa Tgcagagata gaatgatggg ctataacaat catataattg aaagaaagaa	4800
ctTcaaaaat aatcaagTtc agctgtTtga tttataaatg Tgataactaa aacctagaga	4860
ggaaaagagg tactcaagat cacacagtag gagaggactg cagaaacacc aaocccaagc	4920
tctttTgtcc actctTccag cgttctTtct actatactgc ctatcctTta tctagTtacc	4980
aaTaaataac aaaagctTgg accacaatgc ttttattgTc taggaaactc ctgaagaagc	5040
Taaataaaat gggTggggaa tattgTaaat gtaattcagg ctggattaaT aaagaactT	5100
ttTgacattg TaaTgacaa gcacctgcaa TgctgaaagT aattttTcat Tggentgctg	5160
ttTgctgggc TgcatcaaaT cccTgtctct aggcacatgTc tctgaacatt gTgtgtagca	5220
TggctTtcat ttctTtttagg ataaaattca aaaccctTta tctggTtggT aaacctctgc	5280
ctaaTtggga acctctTtTc tccacaactc catattgtac actccaattt catctctgtt	5340
ctccaacct ggaagctatt Tgtcatgatt cctcctTgtg TcattttTtt tctgTcaacc	5400
Ttggggcttt TgtgtTgtct gTtcaactca cctcctTtTta TtgTaaactt ctactcatct	5460
TtcaattTtTc aactTaaTgT ttctcagaga aacctactTt gattttctTg gTccnaacg	5520

-continued

---

gttctctgga tgtgaactct tatagcacat aattttcact tttttccaca aaactcgctc 5580  
 ctatcacctg ttacaagcat ttacctctga taacaagaac tttcaaatat cttagctgtca 5640  
 tgtaagcact tttcataaac attaagagta tctgtgacac ttatgtgtaa tgtttcgtat 5700  
 ctctgaaatt gatatttacc agtcatttat cttggctacc aactaacaac tatccatatt 5760  
 atctgtacca atcagatgta taatcacaat tttgtgtgac agaaaatggc taaacttgat 5820  
 ccaaggctat tacatgcttt atcaactgca caatctttat atatgtcaat tattgatctt 5880  
 taactgattt ccttcttatg gattttctcc tctgcttacc atgtatgcct aacatgacaa 5940  
 aaaaagagcct atcattgcag ccagtatgat aatactcagt ctgtggggct tcttatttgc 6000  
 ttattccatc atcatctgtc ctgcttgatg tctttgccta tgcacaatca tatgacctat 6060  
 cacatctgta tgaagagctg gatgactagg attaatatc tatttttaggt tcttattcag 6120  
 cagaaatatt agataatcaa tgtcttttta ttctgttagg tgtgaaatcc agacaattga 6180  
 agatggggca tatcagagcc taagccacct ctctacctta atattgacag gaaaccccat 6240  
 ccagagttaa gccctgggag ccttttctgg actatcaagt ttacagaagc tgggtgctgt 6300  
 ggagacaaat cttagcatctc tagagaactt cccattgga catctcaaaa ctttgaaga 6360  
 acttaatgtg gctcacaatc ttatccaatc tttcaaatc cctgagtatt tttctaatct 6420  
 gaccaatcta gagcacttgg acctttccag caacaagatt caaagtattt attgcacaga 6480  
 cttgctgggt ctacatcaaa tgccctact caatctctct tttagacctgt cctgaaccc 6540  
 tatgaacttt atccaaccag gtgcatttaa agaaattagg cttcataagc tgactttaag 6600  
 aaataatatt gatagtttaa atgtaatgaa aacttgatt caaggctctg ctggtttaga 6660  
 agtccatcgt ttggttctgg gagaatttag aaatgaagga aacttgaaa agtttgacaa 6720  
 atctgtctca gagggcctgt gcaatttgac cattgaagaa ttcccgatta gcatacttag 6780  
 actactacct cgatgatatt attgacttat ttaattggtt gacaaatggt tcttcatttt 6840  
 ccctggtgag tgtgactatt gaaagggtaa aagacttttc ttataatttc ggatggcaac 6900  
 atttagaatt agttaactgt aaatttgac agtttccac attgaaactc aaatctctca 6960  
 aaaggcttac tttcacttcc aacaaagggt ggaatgcttt ttcagaagtt gatctacca 7020  
 gccttgagtt tctagatctc agtagaaatg gcttgagttt caaagggtgc tgtctcaaa 7080  
 gtgattttgg gacaaccagc cttaaagtatt tagatctgag cttcaatggt gttattacca 7140  
 tgagttcaaa cttcttgggc ttagaacaac tagaacatct ggatttccag cattccaatt 7200  
 tgaacaaaat gagtgagttt tcagtattcc tatcactcag aaacctcatt taccttgaca 7260  
 tttctcatac tcacaccaga gttgctttca atggcatctt caatggcttg tccagtctcg 7320  
 aagtcttgaa aatggctggc aattctttcc aggaaaactt ccttccagat atcttcacag 7380  
 agctgagaaa cttgaccttc ctggaacctc ctcaagtgtca actggagcag ttgtctccaa 7440  
 cagcatttaa ctcaactctc agtcttcagg tactaaatat gagccacaac aacttctttt 7500  
 cattggatac gtttctttat aagtgtctga actccttcca ggttcttgat tacagtctca 7560  
 atcacataat gacttccaaa aacaggaac tacagcattt tccaagtagt cttagctttct 7620  
 taaactttac tcagaatgac tttgcttgta cttgtgaaca ccagagtttc ctgcaatgga 7680  
 tcaaggacca gaggcagctc ttggtggaag ttgaacgaat ggaatgtgca acacctcag 7740  
 ataaagcagg catgcctgtg ctgagtttga atatcacctg tcagatgaat aagaccatca 7800  
 ttggtgtgtc ggtcctcagt gtgcttgtag tatctggtgt agcagttctg gtctataagt 7860

-continued

---

tctattttca cctgatgctt cttgctggct gcataaagta tggtagaggt gaaaacatct	7920
atgatgcctt tghtatctac tcaagccagg atgaggactg ggtaaggaat gagctagtaa	7980
agaatttaga agaaggggtg cctccatttc agctctgcct tctactacaga gactttattc	8040
ccggtgtggc cattgctgcc aacatcatcc atgaaggttt ccataaaagc cgaaggtga	8100
ttgttgggt gtcccagcac ttcattccaga gccgctgggt tatctttgaa tatgagattg	8160
ctcagacctg gcagtttctg agcagtcgtg ctggtatcat cttcattgtc ctgcagaagg	8220
tggagaagac cctgctcagg cagcaggtgg agctgtaccg ccttctcagc aggaacactt	8280
acctggagtg ggaggacagt gtcctggggc ggcacatctt ctggagacga ctcagaaaag	8340
ccctgctgga tggtaaatca tggaaatccag aaggaacagt gggtagcagga tgcaattggc	8400
aggaagcaac atctatctga agaggaaaa taaaaacctc ctgaggcatt tcttgcccag	8460
ctgggtccaa cacttgttca gttataaagt attaaatgct gccacatgtc aggccttatg	8520
ctaaggggta gtaattccat ggtgcactag atatgcaggg ctgctaactc caaggagctt	8580
ccagtgcaga gggaataaat gctagactaa aatacagagt cttccagggt ggcatttcaa	8640
ccaactcagt caaggaacct atgacaaaaga aagtcatttc aactcttacc tcatcaagtt	8700
gaataaagac agagaaaaca gaaagagaca ttgttctttt cctgagtcct ttgaatggaa	8760
attgtattat gttatagcca tcataaaacc attttggtag ttttgactga actgggtgtt	8820
cactttttcc tttttgattg aatacaattt aaattctact tgatgactgc agtcgtcaag	8880
gggctcctga tgcaagatgc cccttcatt ttaagtctgt ctccttacag aggttaaagt	8940
ctagtggcta attcctaagg aaacctgatt aacacatgct cacaaccatc ctggtcattc	9000
tcgagcatgt tctatttttt aactaatcac ccctgatata tttttatttt tatatatcca	9060
gttttcaatt ttttactgct tgcctataag ctaatatcat aaataagggt gtttaagacg	9120
tgcttcaaat atccatatta accactattt ttcaaggaag tatggaaaag tacactctgt	9180
cactttgtca ctcgatgtca ttccaaagtt attgcctact aagtaatgac tgcatgaaa	9240
gcagcattga aataatttgt ttaaaggggg cactctttta aacgggaaga aaatttccgc	9300
ttcttggtct tatcatggac aatttgggct ataggcatga aggaagtggg attacctcag	9360
gaagtcacct tttcttgatt ccagaaacat atgggctgat aaaccggggg tgacctcatg	9420
aaatgagttg cagcagatgt ttatttttt cagaacaagt gatgtttgat ggacctatga	9480
atctatttag ggagacacag atggctggga tcctctccct gtaccttct cactgccagg	9540
agaactacgt gtgaaggat tcaaggcagg gagtatacat tgctgtttcc tgttgggcaa	9600
tgctccttga ccacattttg ggaagagtgg atgttatcat tgagaaaaa atgtgtctgg	9660
aattaatggg gttcttataa agaaggttcc cagaaaagaa tgttcattcc agcttcttca	9720
ggaaacagga acattcaagg aaaaggacaa tcaggatgtc atcagggaaa tgaataaaa	9780
aaccacaatg agatatcacc ttataccagg tagatggcta ctataaaaa atgaagtgtc	9840
atcaaggata tagagaaatt ggaaccttc ttcactgctg gagggaatgg aaaatggtgt	9900
agcgttatg aaaaacagta cggaggttcc tcaaaaatta aaaatagaac tgctatatga	9960
tccagcaate tcaactctgt atatatacc aaaataattg aaatcagaat tccaagaaaa	10020
tatttacact cccatgttca ttgtggcact cttcacaatc actgtttcca aagttatgga	10080
aacaacccaa atttccattg gaaaataaat ggacaaagga aatgtgcata taacgtacaa	10140
tggggatatt attcagccta aaaaagggg ggatcctggt atttatgaca acatgaataa	10200
accggaggc cattatgcta tgtaaatga gcaagtaaca gaaagacaaa tactgcctga	10260

-continued

---

```

tttcatttat atgaggttct aaaatagtca aactcataga agcagagaat agaacagtgg 10320
ttcttaggga aaaggaggaa gggagaaatg aggaaatagg gagttgtcta attggtataa 10380
aattatagta tgcaagatga attagctcta aagatcagct gtatagcaga gttcgtataa 10440
tgaacaatac tgtattatgc acttaacatt ttgttaagag ggtacctctc atgttaagtg 10500
ttcttaccat atacatatac acaaggaagc ttttgagggt gatggatata tttattacct 10560
tgattgtggt gatggtttga caggtatgtg actatgtcta aactcatcaa attgtatata 10620
ttaaataat gtagttttat aatatcaaaa aaaaaaaaaa aaaaa 10665

```

```

<210> SEQ ID NO 63
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 63

```

```

attttgtttt gtttttaaag acttggcc 28

```

```

<210> SEQ ID NO 64
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 64

```

```

gtcatgtgta atcattgcag gtggttcc 28

```

```

<210> SEQ ID NO 65
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 65

```

```

atgtcttttt attcctgtag gtgtgaaa 28

```

```

<210> SEQ ID NO 66
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: A primer

```

```

<400> SEQUENCE: 66

```

```

gattagcata cttagactac tacctocatg 30

```

```

<210> SEQ ID NO 67
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 67

```

```

gatcaacttc tgaaaagca ttcccac 27

```

```

<210> SEQ ID NO 68
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: A primer

```

```

<400> SEQUENCE: 68

```

```

ggttgctggt ctcaaagtga ttttgggaga a 31

```

-continued

<210> SEQ ID NO 69  
 <211> LENGTH: 30  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 69

acctgaagac tggagagtga gttaaatgct 30

<210> SEQ ID NO 70  
 <211> LENGTH: 1360  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (1)...(1360)  
 <223> OTHER INFORMATION: n = A,T,C or G

<400> SEQUENCE: 70

ttccacttct aagagctgcc tagagtagtc aagattatag agacaaaagt agtgcataga 60  
 ttcaagggcc tagggaaagg ggaaatgggg agttatttat taatgaatag tggatgatgat 120  
 tgtacaaaaa tatgaacata attaatgcca ctaaattgtn cacatacaaa tggcaagat 180  
 aataaatttt atgttatgtc atgttatggt atgtgatttt accataatac agaaaatgaa 240  
 aaaagaaaag aaagaaagta aagcttagcg gtttncatga cttgnccaat gcctcaaagc 300  
 catgagtcga cccagctgag atctganctt cagtatatc cattctgaaa tcccagactt 360  
 ttcccaatct tcttgtactt ttcaaactgt gtttcagttg aggtttattt tcagttttgt 420  
 atgtgagttt cttcgaaga agggcgggcc aaattgtgtc ctgcacaaac ctacatatcg 480  
 aagtcctaac ccctctacct cagactatga ctgtatatgg agagagagcc ttgaaagagg 540  
 tatgtaagggt agaagaggt cattatggtg ggccctaac caacataact ggtgtcctta 600  
 taagaagggg agattagaat tcagacacac ttgctgacac cttgagttca gactggaagc 660  
 ctctagaatt gtgagaaaat gaatgtctgt tgtttaagcc acccagctcg tggatatttc 720  
 ttatggcagc cccagcaaac taatacaaat agtgtttcca cagctgaaac aaaattggaa 780  
 aatcaccgct atcctagaga gttacaaggg ctattttaat agaacctgat tgttttctca 840  
 aattcaccaa gccagcgagc aggtcagatg actaattggg ataaaagcca actagcttcc 900  
 tcttctgtgt tctttagcca ctggtctgca ggcgttttct tcttctaact tectctctg 960  
 tgacaaaaga gataactatt agagaacaa aagtcagaa tgctaagggt gccgctttca 1020  
 ctctctctca ccctttagcc cagaactgct ttgaatacac caattgctgt ggggcccgtc 1080  
 gaggaagaga agacaccagt gcctcagaaa ctgctcggtc agacgggtgat agcgagccac 1140  
 gcattcacag gccactgct gctcacagaa gcagtgagga tgatgccagg atgatgtctg 1200  
 cctcgcgcct ggtgggact ctgatcccag ccatggcctt cctctcctgc gtgagaccag 1260  
 aaagctggga gccctgctgt gaggtatgtg gctggagtca gctctctga actttcctc 1320  
 acttctgccc agaacttctc actgtgtgcc ctggtttgtt 1360

<210> SEQ ID NO 71  
 <211> LENGTH: 1333  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 71

cgcacatgg atttgtgtgt catcctgtg caggggccc gctcatcttc tctgtatcct 60  
 tccaatttta gtatagtgtc tactgacgca agcagcatat tggatatttt attacctaca 120

-continued

---

```

ttttacatat gataaaatga ggctcactga ggtttttctt ttgttcgttt tattttgttt 180
tgtttttaa gacttggccc taaaccacac agaagagctg gcatgaaacc cagagctttc 240
agactccgga gcctcagccc ttcaccccga ttccattgct tcttgctaaa tgctgcccgt 300
ttatcacgga ggtagaatg ctgagcacgt agtaggtgct ctttactttc taatctagag 360
taagacaatt tataagcatg aattgagtga atggatggat ggatataatg atggaaggat 420
ggacagatgg atgaaagggt gactgaattt tgtgcttgca caaaaagagg cccctctcca 480
ccatctctgg tctaggagag gggagttggg agaccatgca gtaaagatac ttcattgcat 540
gtgtaatcat tgcaggtggg tcctaataat acttatcaat gcatggagct gaatttctac 600
aaaatccccg acaacctccc cttctcaacc aagaacctgg acctgagctt taatccccctg 660
aggcatttag gcagctatag cttcttcagt tcccagaac tgcaggtgct ggatttatcc 720
aggtaatgaa tccacttta catactgcac aaggtgaggt gttcattgct ctatcatttc 780
attattggac tggaaagctt ggtttgtgga gtctcatctt cttcactta ttcattcata 840
caacagatgt cttattaact atataacct gagcaagcta cctctattct ccaggctctca 900
gttttcta ctgtgaagta ggcagttggc tgagacagct tctaagggca attctaattt 960
taggttttct ttaagacag gagagaaaat tagcttaaat tctttcataa gcagctattt 1020
attgactact tgctatatgt tgtacactct gcaagaagac aggcataat t gatatataa 1080
cacacagccc ctgttgtaa ggagcatat cttcttgaaa gagttaatac cttaaagtcc 1140
tgggataggt cctgggtaca tagtatatag tcaacacatt ttaattatga tttttggat 1200
ctggaaactg atataaagat agcgacatat aacagtaggt gataaattat gtttaacta 1260
aaggtaacta attgtatttt tcagaagagg ggccttctct gtggtgggta gtcaagaaag 1320
attcatgaac tgc 1333

```

```

<210> SEQ ID NO 72
<211> LENGTH: 6786
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(6786)
<223> OTHER INFORMATION: n = A,T,C or G

<400> SEQUENCE: 72

```

```

ggtaagaatg ctttgtgata gccagcagc cttctttccc ctatagaaat atatatatan 60
tctttttata ggtgaggaac tgaagcttga ataatttaa tgacttatat acatnatcat 120
tgcttgtag ccacagacca gagatttaag ttcncatctc cagaatccaa cttaaatggt 180
ttctttgtct taatactcta cttctctaaa gtgattatca ccaatgtaat gatatagagn 240
cacagcaaga ccctttcctt ctcacctaat gtatagagca atgcagagat agaatgatgg 300
gctataacaa tcatataaatt gaaagaaaga acttcaaaaa taatcaagtt cagctgtttg 360
atttataaat gtgataacta aaacctagag aggaaaagag gtactcaaga tcacacagta 420
ggagaggact gcagaacac caaacccaag ctcttttgtc cactcttcca gcgttctttc 480
tactatactg cctatccttt atctagttac caataataa caaaagcttg gaccacaatg 540
cttttattgt ctaggaaact cctgaagaag ctaataaaa tgggtgggga atattgtaa 600
tgtaattcag gctggattaa gaaagaactt atttgacatt gtaactgaca agcacctgca 660
atgctgaaag gaatttttca ttggentgct gtttgctggg ctgcatcaaa gcctgtctc 720

```

-continued

taggcacatgt	ctctgaacat	tgtgtgtagc	atggctttca	tttcttttag	gataaaatc	780
aaaacccttt	atctggttgg	taaacctctg	cctaattggg	aaccttcttt	ctccacaact	840
ccatattgta	cactccaatt	tcatctctgt	tctccaacca	tggaagctat	ttgtcatgat	900
tctctcttgt	gtcatttttt	ttctgtcaac	cttggggctt	ttgtgtttgc	tgttcacttc	960
acctcctttt	atgtttaact	tctactcatc	tttcaatttt	caacttaagt	gttctcagag	1020
aaacctactt	tgattttctt	ggtccanaac	ggttctctgg	atgtgaactc	ttatagcaca	1080
taattttcac	ttttttccac	aaaactcgct	cctatcacct	gttacaagca	tttacctctg	1140
ataacaagaa	ctttcaaata	tctagctgtc	atgtaagcac	ttttcataaa	cattaagagt	1200
atctgtgaca	cttatgtgta	atgtttcgta	tctctgaaat	tgatatttac	cagtcattta	1260
tcttggtctc	caactaacia	ctatccatat	tatctgtacc	aatcagatgt	ataatcacia	1320
ttttgtgtga	cagaaaatgg	ctaaaactga	tccaaggcta	ttacatgctt	tatcaactgc	1380
acaatcttta	tatatgtcaa	ttattgatct	ttanctgatt	tccttcttat	ggattttctc	1440
ctctgcttat	catgatgtcc	taacatgaca	aaaaagagcc	tatcattgca	gccagtatga	1500
taataactcag	tctgtggggc	ttcttatttg	cttattccat	catcatctgt	cctgcttgat	1560
gtctttgcct	atgcacaatc	atatgacca	tcacatctgt	atgaagagct	ggatgactag	1620
gattaatatt	ctatttttag	ttcttattca	gcagaaatat	tagataatca	atgtcttttt	1680
attcctgtag	gtgtgaaatc	cagacaatgt	aagatggggc	atcagagagc	ctaagccacc	1740
tctctacctt	aatattgaca	ggaaacccca	tccagagttt	agccctggga	gccttttctg	1800
gactatcaag	tttacagaag	ctgggtggctg	tggagacaaa	tctagcatct	ctagagaact	1860
tccccattgg	acatctcaaa	actttgaaag	aacttaatgt	ggctcacaat	cttatccaat	1920
ctttcaaatt	acctgagtat	ttttctaact	tgaccaatct	agagcacttg	gacctttcca	1980
gcaacaagat	tcaaagtatt	tattgcacag	acttgccggg	tctacatcaa	atgcccttac	2040
tcaatctctc	tttagacctg	tccctgaacc	ctatgaactt	tatccaacca	ggtgcattta	2100
aagaaattag	gcttcataag	ctgactttaa	gaaataatct	tgatagttaa	aatgtaatga	2160
aaacttgat	tcaaggtctg	gctggtttag	aagtccatcg	tttggttctg	ggagaattta	2220
gaaatgaagg	aaacttgtaa	aagtttgaca	aatctgctct	agagggcctg	tgcaatttga	2280
ccattgaaga	attccgatta	gcatacttag	actactacct	cgatgatatt	attgacttat	2340
ttaattgttt	gacaaatggt	tcttcatttt	ccttgggtgag	tgtgactatt	gaaagggtaa	2400
aagacttttc	ttataatttc	ggatggcaac	atttagaatt	agttaactgt	aaatttggac	2460
agtttccacc	attgaaactc	aaatctctca	aaaggcttac	tttcaacttc	aacaaagggtg	2520
ggaatgcttt	ttcagaagtt	gatctaccaa	gccttgagtt	tctagatctc	agtagaatg	2580
gcttgagttt	caaaggttgc	tgttctcaaa	gtgattttgg	gacaaccagc	ctaaagtatt	2640
tagatctgag	cttcaatggt	gttattacca	tgagttcaaa	cttcttgggc	ttagaacaa	2700
tagaacatct	ggattttccag	cattccaatt	tgaacaaat	gagtgagttt	tcagtattcc	2760
tatcactcag	aaacctcatt	taccttgaca	tttctcatac	tcacaccaga	gttgctttca	2820
atggcatctt	caatggcttg	tccagtctcg	aagtcttgaa	aatggctggc	aattctttcc	2880
aggaaaactt	ccttccagat	atcttcacag	agctgagaaa	cttgaccttc	ctggacctct	2940
ctcagtgta	actggagcag	ttgtctccaa	cagcatttaa	ctcactctcc	agtcttcagg	3000
tactaaat	gagccacaac	aacttctttt	cattggatc	gtttccttat	aagtgtctga	3060
actccctcca	ggttcttgat	tacagtctca	atcacataat	gacttccaaa	aaacaggaac	3120

-continued

---

tacagcattt	tccaagtagt	ctagctttct	taaatcttac	tcagaatgac	tttgcttgta	3180
cttgtaaca	ccagagtttc	ctgcaatgga	tcaaggacca	gaggcagctc	ttggtggaag	3240
ttgaacgaat	ggaatgtgca	acaccttcag	ataagcaggg	catgcctgtg	ctgagtttga	3300
atatcacctg	tcagatgaat	aagaccatca	ttggtgtgtc	ggtcctcagt	gtgctttag	3360
tatctgttgt	agcagttctg	gtctataagt	tctattttca	cctgatgctt	cttgctggct	3420
gcataaagta	tggtagaggt	gaaaacatct	atgatgcctt	tggtatctac	tcaagccagg	3480
atgaggactg	ggtaaggaat	gagctagtaa	agaattttaga	agaaggggtg	cctccatttc	3540
agctctgctc	tactacaga	gactttattc	ccggtgtggc	cattgctgcc	aacatcatcc	3600
atgaaggttt	ccataaaagc	cgaaagggtga	ttgttgggt	gtcccagcac	ttcatccaga	3660
gccgctgggt	tatctttgaa	tatgagattg	ctcagacctg	gcagtttctg	agcagtcgtg	3720
ctggtatcat	cttcattgtc	ctgcagaagg	tggagaagac	cctgctcagg	cagcaggtgg	3780
agctgtaccg	ccttctcagc	aggaacactt	acctggagtg	ggaggacagt	gtcctggggc	3840
ggcacatctt	ctggagacga	ctcagaaaag	ccctgctgga	tggtaaatca	tggaatccag	3900
aaggaacagt	gggtacagga	tgcaattggc	aggaagcaac	atctatctga	agaggaaaaa	3960
taaaaacctc	ctgaggcatt	tcttgcccag	ctgggtccaa	cacttgttca	gttaataagt	4020
attaaatgct	gccacatgtc	aggccttatg	ctaaggggtga	gtaattccat	ggtgcactag	4080
atatgcaggg	ctgctaactc	caaggagctt	ccagtgcaga	gggaataaat	gctagactaa	4140
aatacagagt	cttccagggtg	ggcatttcaa	ccaactcagt	caaggaacc	atgacaaaga	4200
aagtcatctc	aactcttacc	tcatcaagtt	gaataaagac	agagaaaaa	gaaagagaca	4260
ttgttctttt	cctgagtctt	ttgaatggaa	attgtattat	gttatagcca	tcataaaacc	4320
atthtggtag	ttttgactga	actgggtgtt	cactttttcc	tttttgattg	aatacaattt	4380
aaattctact	tgatgactgc	agtcgtcaag	gggctcctga	tgcaagatgc	cccttccatt	4440
ttaagtctgt	ctccttacag	aggttaaagt	ctagtggcta	attcctaagg	aaacctgatt	4500
aacacatgct	cacaaccatc	ctggctcatc	tcgagcatgt	tctatthttt	aactaatcac	4560
ccctgatata	tttttatttt	tatatatcca	gttttcattt	ttttacgtct	tgctataag	4620
ctaataatcat	aaataaggtt	gtttaagacg	tgettcaaat	atccatatta	accactattt	4680
ttcaaggaag	tatgaaaaag	tacactctgt	cactttgtca	ctcgatgtca	ttccaaagtt	4740
attgcctact	aagtaatgac	tgatcatgaa	gcagcattga	aataatttgt	ttaaaggggg	4800
cactctttta	aacgggaaga	aaatttccgc	ttcctggtct	tatcatggac	aatttgggct	4860
agaggcagga	aggaagtggg	atgacctcag	gaggtcacct	tttcttgatt	ccagaaacat	4920
atgggctgat	aaaccggggg	tgacctcatg	aaatgagttg	cagcagaagt	ttatthtttt	4980
cagaacaagt	gatgtttgat	ggacctctga	atctcttttag	ggagacacag	atggctggga	5040
tcctctccct	gtacccttct	cactgccagg	agaactacgt	gtgaaggat	tcaaggcagg	5100
gagtatacat	tgtcttttcc	tgttgggcaa	tgctccttga	ccacattttg	ggaagagtgg	5160
atgttatcat	tgagaaaaa	atgtgtctgg	aattaatggg	gttcttataa	agaaggttcc	5220
cagaaaaaaa	tgttcatcca	gcctcctcag	aaacagaaca	ttcaagaaaa	ggacaatcag	5280
gatgtcatca	gggaaatgaa	aataaaaaacc	acaatgagat	atcaccttat	accaggtaga	5340
atggctacta	taaaaaatg	aagtgtcatc	aaggatatag	agaatttggg	acccttcttc	5400
actgctggag	ggaatggaaa	atggtgtagc	cgttatgaaa	aacagtacgg	aggtttctca	5460



-continued

---

aaaattaaaa atagaactgc tatatgatcc agcaatctca cttctgtata tatacccaaa	5520
ataattgaaa tcagaatttc aagaaaatat ttactactccc atgttcattg tggcactctt	5580
cacaatcact gtttccaaag ttatggaaac aaccctaaatt tccattgaaa aataaatgga	5640
caagaaaaat gtgcatatac gtacaatggg atattattca gcctaaaaaa aggggggnatc	5700
ctgttattta tgacaacatg aataaaccoc gagccattat gctatgtaaa atgagcaagt	5760
aacagaaaga caaatactgc ctgatttcat ttatatgagg ttctaaaaata gtcaaaactca	5820
tagaagcaga gaatagaaca gtggttccta gggaaaagga ggaagggaga aatgaggaaa	5880
tagggagttg tctaattggt ataaaattat agtatgcaag atgaattagc tctaaagatc	5940
agctgtatag cagagtctgt ataatagaaca atactgtatt atgcacttaa cattttgtta	6000
agaggttacc tctcatgta agtgttctta ccatatacat atacacaagg aagcttttgg	6060
aggtgatgga tatatttatt accttgattg tggatgatgg ttgacaggta tgtgactatg	6120
tctaaactca tcaaatgtga tacattaaat atatgcagtt ttataatata aattatgtct	6180
gaatgaagct ataaaaaga aaagacaaca aaattcagtt gtcaaaactg gaaatatgac	6240
cacagtcaga agtgtttgtt actgagtgtt tcagagtgtg tttggttga gcaggcttag	6300
ggtgattgaa catccctggg tgtgtttcca tgtctcatgt actagtgtgaa gtagatgtgt	6360
gcatttgtgc acatatacct atgtatcctc atcagggctg tgtgtatttg aaagtgtgtg	6420
tgtccgcatg atcatatctg tatagaagag agtgtgatta tatttcttga agaatacatc	6480
catttgaat ggatgtctat ggctgtttga gatgagttct ctactcttgt gcttgtacag	6540
tagtctcccc ttatccctta tgcttgggtg atacgttctt agaccccaag tggatctctg	6600
agaccgcaga tggtagcaaa cctcatatata gcaatatttt ttctatata taaataccta	6660
agataaagtt catcttctga attaggcaca gtaagagatt aacaataact aacaataaaa	6720
ttgaatagtt ataataatat attgtaataa aagttatgtg aatgtgatct ctttcttttc	6780
tctctc	6786

---

What is claimed is:

1. An isolated and purified DNA molecule comprising a 45 TLR4 human genomic DNA segment comprising exon 4, wherein the codon for the residue at position 299 in TLR4 encodes a naturally occurring amino acid residue other than aspartic acid, or wherein the codon for the residue at position 399 in TLR4 encodes a naturally occurring amino acid residue other than threonine.
2. The DNA molecule of claim 1 wherein the TLR4 comprises a glycine residue at position 299 or an isoleucine residue at position 399.
3. An isolated host cell, comprising the DNA molecule of 55 claim 1.
4. The DNA molecule of claim 1 wherein the DNA segment comprises SEQ ID NO:62.
5. The DNA molecule of claim 1 wherein the DNA segment 60 comprises SEQ ID NO:70.
6. The DNA molecule of claim 1 wherein the DNA segment comprises SEQ ID NO:71.
7. The DNA molecule of claim 1 wherein the DNA segment comprises SEQ ID NO:72.
8. The DNA molecule of claim 1 wherein the DNA segment 65 comprises GGT at the codon for residue 299.
9. The DNA molecule of claim 1 wherein the DNA segment comprises ATC at the codon for residue 399.
10. An isolated and purified nucleic acid molecule comprising a human nucleic acid segment encoding a variant TLR4, wherein the codon for the residue at position 299 in TLR4 encodes a naturally occurring amino acid residue other than aspartic acid, or wherein the codon for the residue at position 399 in TLR4 encodes a naturally occurring amino acid residue other than threonine.
11. The nucleic acid molecule of claim 10 wherein the variant comprises a glycine residue at position 299 or an isoleucine residue at position 399.
12. A method to prepare a variant TLR4, comprising: culturing a host cell transformed with the nucleic acid molecule of claim 10 operably linked to a promoter, so that said host cell expresses the variant TLR4.
13. The method of claim 12 further comprising isolating TLR4 from the host cell.
14. The nucleic acid molecule of claim 10 wherein the human nucleic acid segment comprises GGT at the codon for residue 299.
15. The nucleic acid molecule of claim 10 wherein the human nucleic acid segment comprises ATC at the codon for residue 399.

**67**

**16.** An expression cassette comprising a promoter operably linked to a human DNA segment encoding a variant TLR4, wherein the codon for the residue at position 299 in TLR4 encodes a naturally occurring amino acid residue other than aspartic acid or wherein the codon for the residue at position 399 in TLR4 encodes a naturally occurring amino acid residue other than threonine.

**68**

**17.** The expression cassette of claim **16** wherein the variant TLR4 comprises a glycine residue at position 299 or an isoleucine residue at position 399.

**18.** An isolated host cell comprising the expression cassette of claim **16**.

\* \* \* \* \*

UNITED STATES PATENT AND TRADEMARK OFFICE  
**CERTIFICATE OF CORRECTION**

PATENT NO. : 7,498,151 B2  
APPLICATION NO. : 11/517172  
DATED : March 3, 2009  
INVENTOR(S) : Schwartz et al.

Page 1 of 1

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

On the first page, in field (56), under "Other Publications", in column 1, line 13, delete "May" and insert -- May, --, therefor.

On the first page, in field (56), under "Other Publications", in column 2, line 23, delete "Tlr4" and insert -- TLR4 --, therefor.

On the first page, in field (56), under "Other Publications", in column 2, line 26, delete "(tlr4)" and insert -- (TLR4) --, therefor.

In column 3, line 12, delete "conimon" and insert -- common --, therefor.

In column 13, line 40, delete "Table 1." and insert -- Table 1 --, therefor.

In column 22, line 62, delete "U/ml" and insert -- U/ml --, therefor.

In column 23, line 40, delete "(Iasbinder)" and insert -- (Iasbinder --, therefor.

In column 24, line 67, delete "Thr399Ile" and insert -- Thr399Ile --, therefor.


In column 25, line 1, delete "Thr399Ile" and insert -- Thr399Ile --, therefor.

In column 25, line 55, delete "Tlr4" and insert -- TLR4 --, therefor.

In column 27, line 41, after "LPS" insert -- . --.

Signed and Sealed this

Thirtieth Day of June, 2009



JOHN DOLL  
*Acting Director of the United States Patent and Trademark Office*