



US007888327B2

(12) **United States Patent**
Krieg et al.

(10) **Patent No.:** **US 7,888,327 B2**
(45) **Date of Patent:** ***Feb. 15, 2011**

(54) **METHODS OF USING
IMMUNOSTIMULATORY NUCLEIC ACID
MOLECULES TO TREAT ALLERGIC
CONDITIONS**

(75) Inventors: **Arthur M. Krieg**, Wellesley, MA (US);
Joel N. Kline, Iowa City, IA (US)

(73) Assignee: **University of Iowa Research
Foundation**, Iowa City, IA (US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

This patent is subject to a terminal dis-
claimer.

(21) Appl. No.: **12/383,824**

(22) Filed: **Mar. 25, 2009**

(65) **Prior Publication Data**

US 2010/0125101 A1 May 20, 2010

Related U.S. Application Data

(60) Continuation of application No. 10/894,657, filed on
Jul. 16, 2004, now abandoned, which is a continuation
of application No. 09/818,918, filed on Mar. 27, 2001,
now abandoned, which is a division of application No.
08/738,652, filed on Oct. 30, 1996, now Pat. No. 6,207,
646, which is a continuation-in-part of application No.
08/386,063, filed on Feb. 7, 1995, now Pat. No. 6,194,
388, which is a continuation-in-part of application No.
08/276,358, filed on Jul. 15, 1994, now abandoned.

(51) **Int. Cl.**
A61K 48/00 (2006.01)

(52) **U.S. Cl.** **514/44 R**

(58) **Field of Classification Search** None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

- 3,906,092 A 9/1975 Hilleman et al.
- 5,059,519 A 10/1991 Owerbach
- 5,112,605 A 5/1992 Jardieu et al.
- 5,248,670 A 9/1993 Draper et al.
- 5,498,410 A 3/1996 Gleich
- 5,585,479 A 12/1996 Hoke et al.
- 5,663,153 A 9/1997 Hutcherson et al.
- 5,679,647 A 10/1997 Carson et al.
- 5,681,555 A 10/1997 Gleich
- 5,723,335 A 3/1998 Hutcherson et al.
- 5,726,160 A 3/1998 McMichael
- 5,780,448 A 7/1998 Davis
- 5,786,189 A 7/1998 Loch et al.
- 5,804,566 A 9/1998 Carson et al.
- 5,830,877 A 11/1998 Carson et al.
- 5,849,719 A 12/1998 Carson et al.
- 5,908,620 A 6/1999 Tu et al.
- 5,955,059 A 9/1999 Gilchrest et al.
- 5,955,442 A 9/1999 McMichael
- 5,985,847 A 11/1999 Carson et al.

- 5,994,315 A 11/1999 Nyce et al.
- 6,025,339 A 2/2000 Nyce et al.
- 6,030,955 A 2/2000 Stein et al.
- 6,040,296 A 3/2000 Nyce
- 6,086,898 A 7/2000 DeKruyff et al.
- 6,090,791 A 7/2000 Sato et al.
- 6,096,721 A 8/2000 McMichael
- 6,100,244 A 8/2000 McMichael
- 6,174,872 B1 1/2001 Carson et al.
- 6,194,388 B1 2/2001 Krieg et al.
- 6,207,646 B1 3/2001 Krieg et al.
- 6,214,806 B1 4/2001 Krieg et al.
- 6,218,371 B1 4/2001 Krieg et al.
- 6,221,882 B1 4/2001 Macfarlane
- 6,225,292 B1 5/2001 Raz et al.
- 6,239,116 B1 5/2001 Krieg et al.
- 6,339,068 B1 1/2002 Krieg et al.
- 6,399,630 B1 6/2002 Macfarlane
- 6,406,705 B1 6/2002 Davis et al.
- 6,426,336 B1 7/2002 Carson et al.
- 6,429,199 B1 8/2002 Krieg et al.
- 6,479,504 B1 11/2002 Macfarlane et al.
- 6,498,148 B1 12/2002 Raz
- 6,514,948 B1 2/2003 Raz et al.
- 6,521,637 B2 2/2003 Macfarlane
- 6,558,670 B1 5/2003 Friede et al.
- 6,562,798 B1 5/2003 Schwartz
- 6,589,940 B1 7/2003 Raz et al.
- 6,610,308 B1 8/2003 Haensler
- 6,610,661 B1 8/2003 Carson et al.
- 6,653,292 B1 11/2003 Krieg et al.
- 6,727,230 B1 4/2004 Hutcherson et al.
- 6,787,524 B2 9/2004 Chang et al.
- 6,797,276 B1 9/2004 Glenn et al.
- 6,815,429 B2 11/2004 Agrawal
- 6,821,957 B2 11/2004 Krieg et al.

(Continued)

FOREIGN PATENT DOCUMENTS

EP 0 302 758 2/1989

(Continued)

OTHER PUBLICATIONS

[No Author Listed] National Institute of Health, Publication No.
97-4051, Jul. 1997.

(Continued)

Primary Examiner—N. M Minnifield
(74) *Attorney, Agent, or Firm*—Wolf, Greenfield & Sacks,
P.C.; Gregg C. Benson

(57) **ABSTRACT**

Nucleic acids containing unmethylated CpG dinucleotides
and therapeutic utilities based on their ability to stimulate an
immune response and to redirect a Th2 response to a Th1
response in a subject are disclosed. Methods for treating
atopic diseases, including atopic dermatitis, are disclosed.

9 Claims, 19 Drawing Sheets

U.S. PATENT DOCUMENTS							
6,835,395	B1	12/2004	Semple et al.	2003/0191079	A1	10/2003	Krieg et al.
6,843,992	B2	1/2005	Diamond	2003/0203861	A1	10/2003	Carson et al.
6,887,464	B1	5/2005	Coleman et al.	2003/0212026	A1	11/2003	Krieg et al.
6,936,261	B2	8/2005	Granoff et al.	2003/0224010	A1	12/2003	Davis et al.
6,943,240	B2	9/2005	Bauer et al.	2003/0232074	A1	12/2003	Lipford et al.
6,949,520	B1	9/2005	Hartmann et al.	2003/0232780	A1	12/2003	Carson et al.
6,951,845	B2	10/2005	Carson et al.	2003/0232856	A1	12/2003	Macfarlane
7,001,890	B1	2/2006	Wagner et al.	2004/0006010	A1	1/2004	Carson et al.
7,034,007	B1	4/2006	Nyce et al.	2004/0006034	A1	1/2004	Raz et al.
7,208,478	B2	4/2007	Carson et al.	2004/0009949	A1	1/2004	Krieg
7,223,398	B1	5/2007	Tuck et al.	2004/0030118	A1	2/2004	Wagner et al.
7,223,741	B2	5/2007	Krieg	2004/0038922	A1	2/2004	Haensler et al.
7,271,156	B2	9/2007	Krieg et al.	2004/0047869	A1	3/2004	Garcon et al.
7,354,711	B2	4/2008	Macfarlane	2004/0053880	A1	3/2004	Krieg
7,402,572	B2	7/2008	Krieg et al.	2004/0064064	A1	4/2004	Zhou et al.
7,410,975	B2	8/2008	Lipford et al.	2004/0067902	A9	4/2004	Bratzler et al.
7,488,490	B2	2/2009	Davis et al.	2004/0067905	A1	4/2004	Krieg
7,517,861	B2	4/2009	Krieg et al.	2004/0087534	A1	5/2004	Krieg et al.
7,524,828	B2 *	4/2009	Krieg et al. 514/44 R	2004/0087538	A1	5/2004	Krieg et al.
7,534,772	B2	5/2009	Weiner et al.	2004/0092468	A1	5/2004	Schwartz
7,566,703	B2	7/2009	Krieg et al.	2004/0092472	A1	5/2004	Krieg
7,569,553	B2	8/2009	Krieg	2004/0106568	A1	6/2004	Krieg et al.
7,576,066	B2	8/2009	Krieg	2004/0131628	A1	7/2004	Bratzler et al.
7,585,847	B2 *	9/2009	Bratzler et al. 514/44 R	2004/0132685	A1	7/2004	Krieg et al.
7,605,138	B2 *	10/2009	Krieg 514/44 R	2004/0142469	A1	7/2004	Krieg et al.
7,615,227	B2	11/2009	Klinman et al.	2004/0143112	A1	7/2004	Krieg et al.
7,615,539	B2	11/2009	Krieg et al.	2004/0147468	A1	7/2004	Krieg et al.
7,666,674	B2	2/2010	Klinman et al.	2004/0152649	A1	8/2004	Krieg
7,674,777	B2 *	3/2010	Krieg et al. 514/44 R	2004/0152656	A1	8/2004	Krieg et al.
7,713,529	B2	5/2010	Krieg et al.	2004/0152657	A1	8/2004	Krieg et al.
7,723,022	B2	5/2010	Krieg et al.	2004/0162258	A1	8/2004	Krieg et al.
7,723,500	B2	5/2010	Krieg et al.	2004/0162262	A1	8/2004	Krieg et al.
2001/0044416	A1	11/2001	McCluskie et al.	2004/0167089	A1	8/2004	Krieg et al.
2002/0064515	A1	5/2002	Krieg et al.	2004/0171150	A1	9/2004	Krieg et al.
2002/0065236	A1	5/2002	Yew et al.	2004/0171571	A1	9/2004	Krieg et al.
2002/0086839	A1	7/2002	Raz et al.	2004/0181045	A1	9/2004	Krieg et al.
2002/0091097	A1	7/2002	Bratzler et al.	2004/0198680	A1	10/2004	Krieg
2002/0102255	A1	8/2002	Chang	2004/0198688	A1	10/2004	Krieg et al.
2002/0156033	A1	10/2002	Bratzler et al.	2004/0229835	A1	11/2004	Krieg et al.
2002/0164341	A1	11/2002	Davis et al.	2004/0234512	A1	11/2004	Wagner et al.
2002/0165178	A1	11/2002	Schetter et al.	2004/0235770	A1	11/2004	Davis et al.
2002/0192184	A1	12/2002	Carpentier et al.	2004/0235774	A1	11/2004	Bratzler et al.
2002/0198165	A1	12/2002	Bratzler et al.	2004/0235777	A1	11/2004	Wagner et al.
2003/0022852	A1	1/2003	Van Nest et al.	2004/0235778	A1	11/2004	Wagner et al.
2003/0026782	A1	2/2003	Krieg	2004/0247662	A1	12/2004	Dow et al.
2003/0026801	A1	2/2003	Weiner et al.	2004/0248837	A1	12/2004	Raz et al.
2003/0027782	A1	2/2003	Carson et al.	2004/0266719	A1	12/2004	McCluskie et al.
2003/0049266	A1	3/2003	Fearon et al.	2005/0004061	A1	1/2005	Krieg et al.
2003/0050261	A1	3/2003	Krieg et al.	2005/0004062	A1	1/2005	Krieg et al.
2003/0050263	A1	3/2003	Krieg et al.	2005/0004144	A1	1/2005	Carson et al.
2003/0050268	A1	3/2003	Krieg et al.	2005/0009774	A1	1/2005	Krieg et al.
2003/0055014	A1	3/2003	Bratzler	2005/0013812	A1	1/2005	Dow et al.
2003/0059773	A1	3/2003	Van Nest et al.	2005/0031638	A1	2/2005	Dalemans et al.
2003/0064064	A1	4/2003	Dina et al.	2005/0032734	A1	2/2005	Davis et al.
2003/0078223	A1	4/2003	Raz et al.	2005/0032736	A1	2/2005	Krieg et al.
2003/0087848	A1	5/2003	Bratzler et al.	2005/0037403	A1	2/2005	Krieg et al.
2003/0091599	A1	5/2003	Davis et al.	2005/0037985	A1	2/2005	Krieg et al.
2003/0092663	A1	5/2003	Raz et al.	2005/0042203	A1	2/2005	Davis et al.
2003/0100527	A1	5/2003	Krieg et al.	2005/0043529	A1	2/2005	Davis et al.
2003/0104523	A1	6/2003	Bauer et al.	2005/0049215	A1	3/2005	Krieg et al.
2003/0109469	A1	6/2003	Carson et al.	2005/0049216	A1	3/2005	Krieg et al.
2003/0119773	A1	6/2003	Raz et al.	2005/0054601	A1	3/2005	Wagner et al.
2003/0125279	A1	7/2003	Junghans et al.	2005/0054602	A1	3/2005	Krieg et al.
2003/0133988	A1	7/2003	Fearon et al.	2005/0059619	A1	3/2005	Krieg et al.
2003/0139364	A1	7/2003	Krieg et al.	2005/0059625	A1	3/2005	Krieg et al.
2003/0148316	A1	8/2003	Lipford et al.	2005/0070491	A1	3/2005	Krieg et al.
2003/0148976	A1	8/2003	Krieg et al.	2005/0075302	A1	4/2005	Hutcherson et al.
2003/0166001	A1	9/2003	Lipford	2005/0079152	A1	4/2005	Bot et al.
2003/0176389	A1	9/2003	Raz et al.	2005/0100983	A1	5/2005	Bauer et al.
2003/0181406	A1	9/2003	Schetter et al.	2005/0101554	A1	5/2005	Krieg et al.
2003/0186921	A1	10/2003	Carson et al.	2005/0101557	A1	5/2005	Krieg et al.
				2005/0119273	A1	6/2005	Lipford et al.
				2005/0123523	A1	6/2005	Krieg et al.

2005/0130911 A1	6/2005	Uhlmann et al.	2009/0155307 A1	6/2009	Davis et al.
2005/0148537 A1	7/2005	Krieg et al.	2009/0191188 A1	7/2009	Krieg et al.
2005/0169888 A1	8/2005	Hartmann et al.	2009/0202575 A1	8/2009	Krieg et al.
2005/0171047 A1	8/2005	Krieg et al.	2009/0214578 A1	8/2009	Bauer
2005/0181422 A1	8/2005	Bauer et al.	2009/0306177 A1	12/2009	Uhlmann et al.
2005/0182017 A1	8/2005	Krieg	2009/0311277 A1	12/2009	Krieg et al.
2005/0197314 A1	9/2005	Krieg et al.	2010/0004319 A1*	1/2010	Von Stein et al. 514/44 R
2005/0209183 A1	9/2005	Kippenberger et al.	2010/0125101 A1*	5/2010	Krieg et al. 514/44 R
2005/0209184 A1	9/2005	Klinman et al.	2010/0144846 A1*	6/2010	Jurk et al. 514/44 R
2005/0215500 A1	9/2005	Krieg et al.			
2005/0215501 A1	9/2005	Lipford et al.			
2005/0233995 A1	10/2005	Krieg et al.			
2005/0233999 A1	10/2005	Krieg et al.			
2005/0239732 A1	10/2005	Krieg et al.			
2005/0239733 A1	10/2005	Jurk et al.			
2005/0239734 A1	10/2005	Uhlmann et al.			
2005/0239736 A1	10/2005	Krieg et al.			
2005/0244379 A1	11/2005	Krieg et al.			
2005/0244380 A1	11/2005	Krieg et al.			
2005/0245477 A1	11/2005	Krieg et al.			
2005/0250726 A1	11/2005	Krieg et al.			
2005/0256073 A1	11/2005	Lipford et al.			
2005/0267057 A1	12/2005	Krieg			
2005/0267064 A1	12/2005	Krieg et al.			
2005/0277604 A1	12/2005	Krieg et al.			
2005/0277609 A1	12/2005	Krieg et al.			
2006/0003955 A1	1/2006	Krieg et al.			
2006/0003962 A1	1/2006	Ahluwalia et al.			
2006/0019239 A1	1/2006	Ivins et al.			
2006/0019909 A1	1/2006	Agrawal et al.			
2006/0019916 A1	1/2006	Krieg et al.			
2006/0019923 A1	1/2006	Davis et al.			
2006/0058251 A1	3/2006	Krieg et al.			
2006/0089326 A1	4/2006	Krieg et al.			
2006/0094683 A1	5/2006	Krieg et al.			
2006/0140875 A1	6/2006	Krieg et al.			
2006/0154890 A1	7/2006	Bratzler et al.			
2006/0171968 A1	8/2006	Brimnes et al.			
2006/0172966 A1	8/2006	Lipford et al.			
2006/0188913 A1	8/2006	Krieg et al.			
2006/0211639 A1	9/2006	Bratzler et al.			
2006/0211644 A1	9/2006	Krieg et al.			
2006/0229271 A1	10/2006	Krieg et al.			
2006/0241076 A1	10/2006	Uhlmann et al.			
2006/0246035 A1	11/2006	Ahluwalia et al.			
2006/0286070 A1	12/2006	Hartmann et al.			
2006/0287263 A1	12/2006	Davis et al.			
2007/0009482 A9	1/2007	Krieg et al.			
2007/0010470 A9	1/2007	Krieg et al.			
2007/0037767 A1	2/2007	Bratzler et al.			
2007/0065467 A1	3/2007	Krieg et al.			
2007/0066553 A1	3/2007	Krieg et al.			
2007/0066554 A1	3/2007	Krieg et al.			
2007/0078104 A1	4/2007	Krieg et al.			
2007/0129320 A9	6/2007	Davis et al.			
2007/0142315 A1	6/2007	Forsbach et al.			
2007/0184465 A1	8/2007	Wagner et al.			
2007/0202128 A1	8/2007	Krieg et al.			
2007/0224210 A1	9/2007	Krieg et al.			
2007/0232622 A1	10/2007	Lipford et al.			
2008/0009455 A9	1/2008	Krieg et al.			
2008/0026011 A1	1/2008	Krieg et al.			
2008/0031936 A1	2/2008	Krieg et al.			
2008/0045473 A1	2/2008	Uhlmann et al.			
2008/0113929 A1	5/2008	Lipford et al.			
2008/0226649 A1	9/2008	Schetter et al.			
2009/0017021 A1	1/2009	Davis et al.			
2009/0060927 A1	3/2009	Wagner et al.			
2009/0062224 A1	3/2009	Kim et al.			
2009/0117132 A1	5/2009	Readett et al.			
2009/0137519 A1	5/2009	Krieg et al.			
2009/0142362 A1	6/2009	Krieg et al.			
2009/0155212 A1	6/2009	Bratzler et al.			

FOREIGN PATENT DOCUMENTS

EP	0 468 520	1/1992
EP	1 671 646 A2	6/2006
WO	WO 91/12811 A1	9/1991
WO	WO 92/03456 A1	3/1992
WO	WO 92/18522 A1	10/1992
WO	WO 92/21353 A1	12/1992
WO	WO 93/09789 A1	5/1993
WO	WO 93/10138 A1	5/1993
WO	WO 94/19945 A1	9/1994
WO	WO 94/29461 A1	12/1994
WO	WO 95/05853 A1	3/1995
WO	WO 95/26204 A1	10/1995
WO	WO 96/02555 A1	2/1996
WO	WO 96/32138	10/1996
WO	WO 96/35782 A1	11/1996
WO	WO 96/40162	12/1996
WO	WO 97/28259 A1	8/1997
WO	WO 98/14210 A1	4/1998
WO	WO 98/16247 A1	4/1998
WO	WO 98/18810 A1	5/1998
WO	WO 98/37919 A1	9/1998
WO	WO 98/40100 A1	9/1998
WO	WO 98/52581 A1	11/1998
WO	WO 98/55495 A2	12/1998
WO	WO 99/11275 A2	3/1999
WO	WO 99/52549 A1	10/1999
WO	WO 98/49288 A1	11/1999
WO	WO 99/56755 A1	11/1999
WO	WO 99/58118 A2	11/1999
WO	WO 99/62923 A2	12/1999
WO	WO 00/06588 A1	2/2000
WO	WO 00/14217 A2	3/2000
WO	WO 00/15256 A2	3/2000
WO	WO 00/16804 A1	3/2000
WO	WO 00/20039 A1	4/2000
WO	WO 00/54803 A2	9/2000
WO	WO 00/61151 A2	10/2000
WO	WO 00/62787 A1	10/2000
WO	WO 00/67023 A1	11/2000
WO	WO 00/67787 A2	11/2000
WO	WO 01/02007 A1	1/2001
WO	WO 01/12223 A2	2/2001
WO	WO 01/35991 A2	5/2001
WO	WO 01/45750 A1	6/2001
WO	WO 01/68144 A2	9/2001
WO	WO 02/28428 A2	4/2002
WO	WO 03/000232 A2	1/2003
WO	WO 03/043572 A2	5/2003
WO	WO 03/094963 A2	11/2003
WO	WO 03/101375 A2	12/2003
WO	WO 2004/007743 A2	1/2004
WO	WO 2004/026888 A2	4/2004
WO	WO 2004-039829 A2	5/2004
WO	WO 2004/041183 A2	5/2004
WO	WO 2004/094671 A2	11/2004
WO	WO 2005/004910 A2	1/2005
WO	WO 2005/023289 A1	3/2005
WO	WO 2006/080946	8/2006
WO	WO 2006/096497	9/2006
WO	WO 2007/031877	3/2007
WO	WO 2007/038720	4/2007

WO WO 2008/030455 A2 3/2008
 WO WO 2008/033432 A2 3/2008
 WO WO 2008/039538 A2 4/2008
 WO WO 2008/068638 A2 6/2008
 WO WO 2008/139262 A2 11/2008

OTHER PUBLICATIONS

[No Author Listed] CpG oligonucleotide adjuvants modulate allergic response in mouse model. *Allergy Medicine, NewsRx.com*. Jan. 16, 2000. (Jahnschmid).

Press Release, Jan. 2007, "Coley Pharmaceutical Group Updates Hepatitis C Drug Development Strategy".

Press Release, Jun. 2007, "Coley Pharmaceutical Group Announces Pfizer's Discontinuation of Clinical Trials for PF-3512676 Combined with Cytotoxic Chemotherapy in Advanced Non Small Cell Lung Cancer".

Press Release, Jun. 2006, "Cytos Biotechnology updates on development of allergy vaccine".

Agrawal et al., Medicinal chemistry and therapeutic potential of CpG DNA. *Trends Mol Med*. Mar. 2002;8(3):114-21.

Agrawal et al., Chapter 19: Pharmacokinetics and bioavailability of antisense oligonucleotides following oral and colorectal administrations in experimental animals. 1998: 525-43.

Ahluwalia et al., Immunostimulatory profiles from two classes of CpG ODN administered subcutaneously to healthy subjects. *ICI FOCUS 2004*. Poster.

Anderson et al., TH2 and 'TH2-like' cells in allergy and asthma: pharmacological perspectives. *Trends Pharmacol Sci*. Sep. 1994;15(9):324-32.

Anitescu et al., Interleukin-10 functions in vitro and in vivo to inhibit bacterial DNA-induced secretion of interleukin-12. *J Interferon Cytokine Res*. Dec. 1997;17(12):781-8.

Arora et al., Immunomodulation by liposome entrapped allergen. *Mol Cell Biochem*. Sep. 21, 1990;97(2):173-9. Abstract Only.

Askenase et al., Gee whiz: CpG DNA allergy therapy! *J Allergy Clin Immunol*. Jul. 2000;106(1 Pt 1):37-40.

Askew et al., CpG DNA induces maturation of dendritic cells with distinct effects on nascent and recycling MHC-II antigen-processing mechanisms. *J Immunol*. Dec. 15, 2000;165(12):6889-95.

Ballas et al., Induction of NK activity in murine and human cells by CpG motifs in oligodeoxynucleotides and bacterial DNA. *J Immunol*. Sep. 1, 1996;157(5):1840-5.

Baral et al., Immunostimulatory CpG oligonucleotides enhance the immune response of anti-idiotype vaccine that mimics carcinoembryonic antigen. *Cancer Immunol Immunother*. May 2003;52(5):317-27.

Barnes et al., New treatments for asthma. *European J Internal Medicine*. 2000;11:9-20. Abstract Only.

Bauer et al., Human TLR9 confers responsiveness to bacterial DNA via species-specific CpG motif recognition. *Proc Natl Acad Sci U S A*. Jul. 31, 2001;98(16):9237-42.

Bauer et al., DNA activates human immune cells through a CpG sequence-dependent manner. *Immunology*. Aug. 1999;97(4):699-705.

Blazar et al., Synthetic unmethylated cytosine-phosphate-guanosine oligodeoxynucleotides are potent stimulators of antileukemia responses in naïve and bone marrow transplant recipients. *Blood*. Aug. 15, 2001;98(4):1217-25.

Boggs et al., Characterization and modulation of immune stimulation by modified oligonucleotides. *Antisense Nucleic Acid Drug Dev*. Oct. 1997;7(5):461-71.

Bohle et al., Oligodeoxynucleotides containing CpG motifs induce IL-12, IL-18 and IFN-gamma production in cells from allergic individuals and inhibit IgE synthesis in vitro. *Eur J Immunol*. Jul. 1999;29(7):2344-53.

Brazolot Millan et al., CpG DNA can induce strong Th 1 humoral and cell-mediated immune responses against hepatitis B surface antigen in young mice. *Proc Natl Acad Sci U S A*. Dec. 22, 1998;95(26):15553-8.

Broide et al., DNA-Based immunization for asthma. *Int Arch Allergy Immunol*. Feb.-Apr. 1999;118(2-4):453-6.

Broide et al., Immunostimulatory DNA sequences inhibit IL-5, eosinophilic inflammation, and airway hyperresponsiveness in mice. *J Immunol*. Dec. 15, 1998;161(12):7054-62.

Broide et al., Modulation of asthmatic response by immunostimulatory DNA sequences. *Springer Semin Immunopathol*. 2000;22(1-2):117-24.

Brunner et al., Enhanced dendritic cell maturation by TNF-alpha or cytidine-phosphate-guanosine DNA drives T cell activation in vitro and therapeutic anti-tumor immune responses in vivo. *J Immunol*. Dec. 1, 2000;165(11):6278-86.

Campbell et al., Allergen immunotherapy: novel approaches in the management of allergic diseases and asthma. *Clin Immunol*. Dec. 2000;97(3):193-202.

Carpentier et al., Oligodeoxynucleotides containing CpG motifs can induce rejection of a neuroblastoma in mice. *Cancer Res*. Nov. 1, 1999;59(21):5429-32.

Carson et al., Oligonucleotide adjuvants for T helper 1 (Th1)-specific vaccination. *J Exp Med*. Nov. 17, 1997;186(10):1621-2.

Cella et al., Plasmacytoid dendritic cells activated by influenza virus and CD40L drive a potent TH1 polarization. *Nat Immunol*. Oct. 2000;1(4):305-10.

Cella et al., Plasmacytoid monocytes migrate to inflamed lymph nodes and produce large amounts of type I interferon. *Nat Med*. Aug. 1999;5(8):919-23.

Chace et al., Bacterial DNA-induced NK cell IFN-gamma production is dependent on macrophage secretion of IL-12. *Clin Immunol Immunopathol*. Aug. 1997;84(2):185-93.

Chan et al., CpG-A and CpG-B oligodeoxynucleotides differentially affect the cytokine profile, chemokine receptor expression and T-cell priming function of human plasmacytoid dendritic cells. *Blood*. 2002;100:50b. Abstract #3666.

Chisholm et al., Airway peptidoglycan and immunostimulatory DNA exposures have divergent effects on the development of airway allergen hypersensitivities. *J Allergy Clin Immunol*. Mar. 2004;113(3):448-54.

Cho et al., Immunostimulatory DNA sequences inhibit respiratory syncytial viral load, airway inflammation, and mucus secretion. *J Allergy Clin Immunol*. Nov. 2001;108(5):697-702.

Cho et al., Immunostimulatory DNA inhibits transforming growth factor-beta expression and airway remodeling. *Am J Respir Cell Mol Biol*. May 2004;30(5):651-61. Epub Nov. 14, 2003.

Choi et al., The level of protection against rotavirus shedding in mice following immunization with a chimeric VP6 protein is dependent on the route and the coadministered adjuvant. *Vaccine*. Mar. 15, 2002;20(13-14):1733-40.

Chu et al., CpG oligodeoxynucleotides act as adjuvants that switch on T helper 1 (Th1) immunity. *J Exp Med*. Nov. 17, 1997;186(10):1623-31.

Cooper et al., Safety and immunogenicity of CPG 7909 injection as an adjuvant to Fluarix influenza vaccine. *Vaccine*. Aug. 13, 2004;22(23-24):3136-43.

Cowdery et al., Bacterial DNA induces NK cells to produce IFN-gamma in vivo and increases the toxicity of lipopolysaccharides. *J Immunol*. Jun. 15, 1996;156(12):4570-5.

Creticos et al., New approaches in immunotherapy: allergen vaccination with immunostimulatory DNA. *Immunol Allergy Clin North Am*. Nov. 2004;24(4):569-81.

Creticos et al., Immunotherapy with immunostimulatory oligonucleotides linked to purified ragweed Amb a 1 allergen: effects on antibody production, nasal allergen provocation, and ragweed seasonal rhinitis. *J Allergy Clin Immunol*. 2002;109(4):741-3.

Daftarian et al., Two distinct pathways of immuno-modulation improve potency of p53 immunization in rejecting established tumors. *Cancer Res*. Aug. 1, 2004;64(15):5407-14.

Dar et al., Attenuated cytokine responses in porcine lymph node cells stimulated with CpG DNA are associated with low frequency of IFN-alpha-producing cells and TLR9 mRNA expression. *Vet Immunol Immunopathol*. Jun. 15, 2008;123(3-4):324-36. Epub Feb. 17, 2008.

Davila et al., Generation of antitumor immunity by cytotoxic T lymphocyte epitope peptide vaccination, CpG-oligodeoxynucleotide adjuvant, and CTLA-4 blockade. *Cancer Res*. Jun. 15, 2003;63(12):3281-8.

- Davis, Use of CpG DNA for enhancing specific immune responses. *Curr Top Microbiol Immunol.* 2000;247:171-83.
- Davis et al., CpG ODN is safe and highly effective in humans as adjuvant to HBV vaccine: Preliminary results of Phase I trial with CpG ODN 7909. Third Annual Conference on Vaccine Res. 2000. Abstract s25, No. 47.
- Durham et al., Immunotherapy and allergic inflammation. *Clin Exp Allergy.* Jan. 1991;21 Suppl 1:206-10.
- Dziedzic et al., Handbook of Experimental Pharmacology, Pharmacology and Therapeutics of Asthma and COPD. 2004;161:273-85.
- Erb et al., Novel vaccines protecting against the development of allergic disorders: a double-edged sword? *Curr Opin in Immunol.* 2002;14:633-43.
- Fanucchi et al., Immunostimulatory oligonucleotides attenuate airways remodeling in allergic monkeys. *Am J Respir Crit Care Med.* Dec. 1, 2004;170(11):1153-7. Epub Aug. 11, 2004.
- Farkas et al., Plasmacytoid dendritic cells activate allergen-specific TH2 memory cells: modulation by CpG oligodeoxynucleotides. *J Allergy Clin Immunol.* Aug. 2004;114(2):436-43.
- Ferreira et al., Customized antigens for desensitizing allergic patients. *Adv Immunol.* 2004;84:79-129. Abstract only.
- Fonseca et al., Use of CpG oligonucleotides in treatment of asthma and allergic disease. *Adv Drug Deliv Rev.* Jan. 9, 2009. [Epub ahead of print]. 7 pages.
- Fornadley, Allergy immunotherapy. *Otolaryngol Clin North Am.* Feb. 1998;31(1):111-27. Abstract Only.
- Gallichan et al., Intranasal immunization with CpG oligodeoxynucleotides as an adjuvant dramatically increases IgA and protection against herpes simplex virus-2 in the genital tract. *J Immunol.* Mar. 1, 2001;166(5):3451-7.
- Gao et al., Bacterial DNA and lipopolysaccharide induce synergistic production of TNF-alpha through a post-transcriptional mechanism. *J Immunol.* Jun. 1, 2001;166(11):6855-60.
- Garbi et al., CpG motifs as proinflammatory factors render autochthonous tumors permissive for infiltration and destruction. *J Immunol.* May 15, 2004;172(10):5861-9.
- Gavett et al., Interleukin 12 inhibits antigen-induced airway hyperresponsiveness, inflammation, and Th2 cytokine expression in mice. *J Exp Med.* Nov. 1, 1995;182(5):1527-36.
- Goldberg et al., Beyond danger: unmethylated CpG dinucleotides and the immunopathogenesis of disease. *Immunol Lett.* Jul. 3, 2000;73(1):13-8.
- Gouttefangeas et al., Problem solving for tumor immunotherapy. *Nat Biotechnol.* May 2000;18(5):491-2.
- Grossmann et al., Avoiding tolerance against prostatic antigens with subdominant peptide epitopes. *J Immunother.* May-Jun. 2001;24(3):237-41.
- Hafner et al., Antimetastatic effect of CpG DNA mediated by type I IFN. *Cancer Res.* Jul. 15, 2001;61(14):5523-8.
- Halpern et al., Bacterial DNA induces murine interferon-gamma production by stimulation of interleukin-12 and tumor necrosis factor-alpha. *Cell Immunol.* Jan. 10, 1996;167(1):72-8.
- Hancock et al., CpG containing oligodeoxynucleotides are potent adjuvants for parenteral vaccination with the fusion (F) protein of respiratory syncytial virus (RSV). *Vaccine.* Sep. 14, 2001;19(32):4874-82.
- Hartmann et al., CpG DNA and LPS induce distinct patterns of activation in human monocytes. *Gene Ther.* May 1999;6(5):893-903.
- Hartmann et al., Mechanism and function of a newly identified CpG DNA motif in human primary B cells. *J Immunol.* Jan. 15, 2000;164(2):944-53.
- Hartmann et al., Delineation of a CpG phosphorothioate oligodeoxynucleotide for activating primate immune responses in vitro and in vivo. *J Immunol.* Feb. 1, 2000;164(3):1617-24.
- Hartmann et al., CpG DNA: a potent signal for growth, activation, and maturation of human dendritic cells. *Proc Natl Acad Sci U S A.* Aug. 3, 1999;96(16):9305-10.
- Hawkes et al., Times of London News International. 4M:18. Printed on Sep. 18, 1999.
- Hayashi et al., TLR9-based immunotherapy for allergic disease. *Am J Med.* Oct. 2006;119(10):897.e1-6.
- Heeg et al., CpG DNA as a Th1 trigger. *Int Arch Allergy Immunol.* Feb. 2000;121(2):87-97.
- Hessel et al., Immunostimulatory oligonucleotides block allergic airway inflammation by inhibiting Th2 cell activation and IgE-mediated cytokine induction. *J Exp Med.* Dec. 5, 2005;202(11):1563-73.
- Higaki et al., Cell death and survival in response to immunostimulatory sequence oligodeoxynucleotides treatment in a murine model of allergic asthma. *J Allergy Clin. Immunol.* Feb. 2006;S156. Abstract #607.
- Hoffjan et al., Toll-like receptors and airway disease. *Drug Discov Today.* 2006;3(3):317-24.
- Hogg, The pathology of asthma. *APMIS.* Oct. 1997;105(10):735-45.
- Hopkin et al., Curbing the CpGs of Bacterial and Viral DNA. *BioMedNet.* Jun. 25, 1999; Issue 57.
- Horner et al., Optimized conjugation ratios lead to allergen immunostimulatory oligodeoxynucleotide conjugates with retained immunogenicity and minimal anaphylactogenicity. *J Allergy Clin Immunol.* Sep. 2002;110(3):413-20.
- Horner et al., Chapter 22: DNA-based immunotherapeutics for allergic disease. From: *Microbial DNA and Host Immunity.* 2002:279-87.
- Horner et al., Immunostimulatory sequence oligodeoxynucleotide-based vaccination and immunomodulation: two unique but complementary strategies for the treatment of allergic diseases. *J Allergy Clin Immunol.* Nov. 2002;110(5):706-12.
- Huang et al., Induction and regulation of Th1-inducing cytokines by bacterial DNA, lipopolysaccharide, and heat-inactivated bacteria. *Infect Immun.* Dec. 1999;67(12):6257-63.
- Hussain et al., CpG oligodeoxynucleotides: a novel therapeutic approach for atopic disorders. *Curr Drug Targets Inflamm Allergy.* Sep. 2003;2(3):199-205.
- Hussain et al., DNA, the immune system, and atopic disease. *J Invest Dermatol Symp Proc.* Jan. 2004;9(1):23-8.
- Hussain et al., Modulation of murine allergic rhinosinusitis by CpG oligodeoxynucleotides. *Laryngoscope.* Oct. 2002;112(10):1819-26.
- Iho et al., Oligodeoxynucleotides containing palindrome sequences with internal 5'-CpG-3' act directly on human NK and activated T cells to induce IFN-gamma production in vitro. *J Immunol.* Oct. 1, 1999;163(7):3642-52.
- Ikeda et al., Microbial DNA and Host Immunity. Chapter 23: Immunostimulatory DNA for allergic asthma. 2002: 289-99.
- Iliev et al., Immunostimulatory oligodeoxynucleotide containing TTTCGTTT motif from *Lactobacillus rhamnosus* GG DNA potentially suppresses OVA-specific IgE production in mice. *Scand J Immunol.* Apr. 2008;67(4):370-6. Epub Feb. 1, 2008.
- Jain et al., CpG DNA and immunotherapy of allergic airway diseases. *Clin Exp Allergy.* Oct. 2003;33(10):1330-5.
- Jain et al., Mucosal immunotherapy with CpG oligodeoxynucleotides reverses a murine model of chronic asthma induced by repeated antigen exposure. *Am J Physiol Lung Cell Mol Physiol.* Nov. 2003;285(5):L1137-46.
- Jain et al., CpG DNA: immunomodulation and remodelling of the asthmatic airway. *Expert Opin Biol Ther.* Sep. 2004;4(9):1533-40.
- Jain et al., CpG-oligodeoxynucleotides inhibit airway remodeling in a murine model of chronic asthma. *J Allergy Clin Immunol.* Dec. 2002;110(6):867-72.
- Jain et al., The promise of CpG DNA in the treatment of asthma. *Recent Res Develop Resp Crit Care Med.* 2002;2:7-18.
- Jakob et al., Activation of cutaneous dendritic cells by CpG-containing oligodeoxynucleotides: a role for dendritic cells in the augmentation of Th1 responses by immunostimulatory DNA. *J Immunol.* Sep. 15, 1998;161(6):3042-9.
- Jakob et al., Bacterial DNA and CpG-containing oligodeoxynucleotides activate cutaneous dendritic cells and induce IL-12 production: implications for the augmentation of Th1 responses. *Int Arch Allergy Immunol.* Feb.-Apr. 1999;118(2-4):457-61.
- Jiang et al., Enhancing immunogenicity by CpG DNA. *Curr Opin Mol Ther.* Apr. 2003;5(2):180-5.
- Jilek et al., Antigen-independent suppression of the allergic immune response to bee venom phospholipase A(2) by DNA vaccination in CBA/J mice. *J Immunol.* Mar. 1, 2001;166(5):3612-21.
- Juffermans et al., CpG oligodeoxynucleotides enhance host defense during murine tuberculosis. *Infect Immun.* Jan. 2002;70(1):147-52.

- Kataoka et al., Antitumor activity of synthetic oligonucleotides with sequences from cDNA encoding proteins of *Mycobacterium bovis* BCG. *Jpn J Cancer Res.* Mar. 1992;83(3):244-7.
- Kataoka et al., Immunotherapeutic potential in guinea-pig tumor model of deoxyribonucleic acid from *Mycobacterium bovis* BCG complexed with poly-L-lysine and carboxymethylcellulose. *Jpn J Med Sci Biol.* Oct. 1990;43(5):171-82.
- Kay et al., Allergy and allergic diseases. Second of two parts. *N Engl J Med.* Jan. 11, 2001;344(2):109-13.
- Kimura et al., Binding of oligonucleotides to scavenger receptors is required for oligonucleotides to augment NK cell activity and induce IFN. *J Biochem (Tokyo).* Nov. 1994;116(5):991-4.
- Kips et al., Interleukin-12 inhibits antigen-induced airway hyper-responsiveness in mice. *Am J Respir Crit Care Med.* Feb. 1996;153(2):535-9.
- Kitagaki et al., Oral administration of CpG-ODNs suppresses antigen-induced asthma in mice. *Clin Exp Immunol.* Feb. 2006;143(2):249-59.
- Kitagaki et al., CpG Oligodeoxynucleotides in Asthma. In: *Microbial DNA and Host Immunity.* 2002. Chapter 24: 301-14.
- Kitagaki et al., Immunomodulatory effects of CpG oligodeoxynucleotides on established Th2 responses. *Clin Diagn Lab Immunol.* Nov. 2002;9(6):1260-9.
- Kline et al., Modulation of airway inflammation by CpG oligodeoxynucleotides in a murine model of asthma. *J Immunol.* Mar. 15, 1998;160(6):2555-9.
- Kline et al., Induction of oral tolerance by CpG-ODNs in a murine model of asthma. *J Allergy Clin Immunol.* Feb. 2004;113(2):S254. Abstract 915.
- Kline et al., Treatment of established asthma in a murine model using CpG oligodeoxynucleotides. *Am J Physiol Lung Cell Mol Physiol.* Jul. 2002;283(1):L170-9.
- Kline et al., DNA therapy for asthma. *Curr Opin Allergy Clin Immunol.* Feb. 2002;2(1):69-73.
- Kline et al., The American Federation for Clinical Research, Mid-western section and Eastern section annual meetings. 1996. Abstracts. *J Investig Med.* Sep. 1996;44(7): 380A.
- Kline et al., Biomedicine '97. Medical research from bench to bedside. Washington, D.C., Apr. 25-27, 1997. Abstracts. *J Investig Med.* Mar. 1997;45(3): 282A.
- Kline et al., American Federation for Medical Research Midwestern Regional Meeting. Chicago, Illinois, Sep. 25-27, 1997. Abstracts. *J Investig Med.* Sep. 1997;45(7): 298A.
- Kline et al., CpG oligodeoxynucleotides do not require TH1 cytokines to prevent eosinophilic airway inflammation in a murine model of asthma. *J Allergy Clin Immunol.* Dec. 1999;104(6):1258-64.
- Kline et al., Effects of CpG DNA on Th1/Th2 balance in asthma. *Curr Top Microbiol Immunol.* 2000;247:211-25.
- Kline et al., Ch. 26: DNA Immunomodulation of asthma. In *Inflammatory Mechanisms in Allergic Diseases.* Ed., Zeiman et al. Marcel Dekker, Inc. 2002;551-64.
- Kline et al., Toll-like receptor 9 activation with CpG oligodeoxynucleotides for asthma therapy. *Drug News Perspect.* Oct. 2008;21(8):434-9.
- Kline, Immunotherapy of asthma using CpG oligodeoxynucleotides. *Immunol Res.* 2007;39(13):279-86. Review.
- Kline, J.N., Eat dirt: CpG DNA and immunomodulation of asthma. *Proc Am Thorac Soc.* Jul. 2007;4(3):283-8. Review.
- Klinman et al., CpG motifs present in bacteria DnA rapidly induce lymphocytes to secrete interleukin 6, interleukin 12, and interferon gamma. *Proc Natl Acad Sci U S A.* Apr. 2, 1996;93(7):2879-83.
- Klinman et al., Immunotherapeutic applications of CpG-containing oligodeoxynucleotides. *Drug News Perspect.* Jun. 2000;13(5):289-96.
- Klinman et al., Immunotherapeutic uses of CpG oligodeoxynucleotides. *Nat Rev Immunol.* Apr. 2004;4(4):249-58.
- Klinman et al., Immune recognition of foreign DNA: a cure for bioterrorism? *Immunity.* Aug. 1999;11(2):123-9.
- Klinman et al., Contribution of CpG motifs to the immunogenicity of DNA vaccines. *J Immunol.* Apr. 15, 1997;158(8):3635-9.
- Klinman et al., Synthetic oligonucleotides as modulators of inflammation. *J Leukoc Biol.* Oct. 2008;84(4):1-7. Epub Apr. 22, 2008.
- Kohama et al., Immunostimulatory oligodeoxynucleotide induces TH1 immune response and inhibition of IgE antibody production to cedar pollen allergens in mice. *J Allergy Clin Immunol.* Dec. 1999;104(6):1231-8.
- Kou et al., [Analysis and regulation of interferon-gamma production by peripheral blood lymphocytes from patients with bronchial asthma] *Aterugi.* Mar. 1994;43(3):482-91. Japanese. Abstract Only.
- Kovarik et al., CpG oligodeoxynucleotides can circumvent the Th2 polarization of neonatal responses to vaccines but may fail to fully redirect Th2 responses established by neonatal priming. *J Immunol.* Feb. 1, 1999;162(3):1611-7.
- Kranzer et al., CpG-oligodeoxynucleotides enhance T-cell receptor-triggered interferon-gamma production and up-regulation of CD69 via induction of antigen-presenting cell-derived interferon type I and interleukin-12. *Immunology.* Feb. 2000;99(2):170-8.
- Krieg et al., Immune effects and therapeutic applications of CpG motifs in bacterial DNA. *Immunopharmacology.* Jul. 25, 2000;48(3):303-5.
- Krieg et al., Leukocyte stimulation by oligodeoxynucleotides. In: *Applied Antisense Oligonucleotide Technology.* 1998:431-48.
- Krieg et al., The role of CpG dinucleotides in DNA vaccines. *Trends Microbiol.* Jan. 1998;6(1):23-7.
- Krieg et al., Applications of immune stimulatory CpG DNA for antigen-specific and antigen-nonspecific cancer immunotherapy. *Eur J Canc.* Oct. 1999; 35/Suppl14:S10. Abstract #14.
- Krieg et al., CpG motifs in bacterial DNA and their immune effects. *Annu Rev Immunol.* 2002;20:709-60.
- Krieg et al., Causing a commotion in the blood: immunotherapy progresses from bacteria to bacterial DNA. *Immunol Today.* Oct. 2000;21(10):521-6.
- Krieg et al., Chapter 8: Immune Stimulation by Oligonucleotides. In: *Antisense Research and Application.* Crooke, Ed. 1998:243-62.
- Krieg et al., Enhancing vaccines with immune stimulatory CpG DNA. *Curr Opin Mol Ther.* Feb. 2001;3(1):15-24.
- Krieg, Chapter 7: CpG oligonucleotides as immune adjuvants. *Ernst Schering Research Found Workshop* 2001; 30:105-18.
- Krieg et al., Chapter 17: Immune stimulation by oligonucleotides. In *Antisense Drug Tech.* 2001;1394:471-515.
- Krieg et al., Mechanisms and applications of immune stimulatory CpG oligodeoxynucleotides. *Biochim Biophys Acta.* Dec. 10, 1999;1489(1):107-16.
- Krieg et al., The CpG motif: Implications for clinical immunology. *BioDrugs.* Nov. 1, 1998;10(5):341-6.
- Krieg, The role of CpG motifs in innate immunity. *Curr Opin Immunol.* Feb. 2000;12(1):35-43.
- Krieg et al., Mechanisms and therapeutic applications of immune stimulatory CpG DNA. *Pharmacol Ther.* Nov. 1999;84(2):113-20.
- Krieg et al., Sequence motifs in adenoviral DNA block immune activation by stimulatory CpG motifs. *Proc Natl Acad Sci U S A.* Oct. 13, 1998;95(21):12631-6.
- Krieg et al., CpG DNA induces sustained IL-12 expression in vivo and resistance to *Listeria monocytogenes* challenge. *J Immunol.* Sep. 1, 1998;161(5):2428-34.
- Krieg et al., CpG DNA: a novel immunomodulator. *Trends Microbiol.* Feb. 1999;7(2):64-5.
- Krieg, Signal transduction induced by immunostimulatory CpG DNA. *Springer Semin Immunopathol.* 2000;22(1-2):97-105.
- Krieg et al., Mechanism of action of CpG DNA. *Curr Top Microbiol Immunol.* 2000;247:1-21.
- Krieg, Therapeutic potential of Toll-like receptor 9 activation. *Nat Rev Drug Discov.* Jun. 2006;5(6):471-84.
- Krieg et al., Induction of systemic TH1-like innate immunity in normal volunteers following subcutaneous but not intravenous administration of CPG 7909, a synthetic B-class CpG oligodeoxynucleotide TLR9 agonist. *J Immunother.* Nov.-Dec. 2004;27(6):460-71.
- Krieg, Now I know my CpGs. *Trends Microbiol.* Jun. 2001;9(6):249-52.
- Krieg et al., Lymphocyte activation mediated by oligodeoxynucleotides or DNA containing novel unmethylated CpG motifs. *American College of Rheumatology 58th National Scientific Meeting.* Minneapolis, Minnesota, Oct. 22, 1994. Abstracts. *Arthritis Rheum.* Sep. 1994;37(9 Suppl).

- Krieg et al., Oligodeoxynucleotide modifications determine the magnitude of B cell stimulation by CpG motifs. *Antisense Nucleic Acid Drug Dev.* 1996 Summer;6(2):133-9.
- Krieg et al., Phosphorothioate oligodeoxynucleotides: antisense or anti-protein? *Antisense Res Dev.* 1995 Winter;5(4):241.
- Krieg, CpG DNA: a pathogenic factor in systemic lupus erythematosus? *J Clin Immunol.* Nov. 1995;15(6):284-92.
- Krieg et al., CpG motifs in bacterial DNA trigger direct B-cell activation. *Nature.* Apr. 6, 1995;374(6522):546-9.
- Krieg, An innate immune defense mechanism based on the recognition of CpG motifs in microbial DNA. *J Lab Clin Med.* Aug. 1996;128(2):128-33.
- Krieg et al., Bacterial DNA or oligonucleotides containing CpG motifs protect mice from lethal *L. monocytogenes* challenge. 1996 Meeting on Molecular Approaches to the Control of Infectious Diseases. Cold Spring Harbor Laboratory, Sep. 9-13, 1996:116.
- Krieg et al., Infection. In: McGraw Hill Book. 1996:242-3.
- Krieg et al., Lymphocyte activation by CpG dinucleotide motifs in prokaryotic DNA. *Trends Microbiol.* Feb. 1996;4(2):73-6.
- Krieg, Immune effects and mechanisms of action of CpG motifs. *Vaccine.* Nov. 8, 2000;19(6):618-22.
- Krug et al., Toll-like receptor expression reveals CpG DNA as a unique microbial stimulus for plasmacytoid dendritic cells which synergizes with CD40 ligand to induce high amounts of IL-12. *Eur J Immunol.* Oct. 2001;31(10):3026-37.
- Kuramoto et al., Induction of T-cell-mediated immunity against MethA fibrosarcoma by intratumoral injections of a bacillus Calmette-Guerin nucleic acid fraction. *Cancer Immunol Immunother.* 1992;34(5):283-8.
- Kuramoto et al., Changes of host cell infiltration into Meth A fibrosarcoma tumor during the course of regression induced by injections of a BCG nucleic acid fraction. *Int J Immunopharmacol.* Jul. 1992;14(5):773-82.
- Kuramoto et al., Oligonucleotide sequences required for natural killer cell activation. *Jpn J Cancer Res.* Nov. 1992;83(11):1128-31.
- Kuramoto et al., In situ infiltration of natural killer-like cells induced by intradermal injection of the nucleic acid fraction from BCG. *Microbiol Immunol.* 1989;33(11):929-40.
- Kussebi et al., Advances in immunological treatment of allergy. *Curr Med Chem.* 2003;2:297-308.
- Leath et al., Novel and emerging therapies for asthma. *Drug Discov Today.* Dec. 2005;10(23-24):1647-55.
- LeClerc et al., The preferential induction of a Th1 immune response by DNA-based immunization is mediated by the immunostimulatory effect of plasmid DNA. *Cell Immunol.* Aug. 1, 1997;179(2):97-106.
- Lee et al., Immuno-stimulatory effects of bacterial-derived plasmids depend on the nature of the antigen in intramuscular DNA inoculations. *Immunology.* Jul. 1998;94(3):285-9.
- Lee et al., Immunostimulatory DNA inhibits allergen-induced peribronchial angiogenesis in mice. *J Allergy Clin Immunol.* Mar. 2006;117(3):597-603. Epub Jan. 27, 2006.
- Leibson et al., Role of gamma-interferon in antibody-producing responses. *Nature.* Jun. 28-Jul. 4, 1984;309(5971):799-801.
- Leonard et al., Interleukin-12: potential role in asthma therapy. *BioDrugs.* 2003;17(1):1-7.
- Lewis et al., Allergy immunotherapy and inhibition of Th2 immune responses: a sufficient strategy? *Curr Opin Immunol.* Oct. 2002;14(5):644-51.
- Lipford et al., CpG-containing synthetic oligonucleotides promote B and cytotoxic T cell responses to protein antigen: a new class of vaccine adjuvants. *Eur J Immunol.* Sep. 1997;27(9):2340-4.
- Lipford et al., Immunostimulatory DNA: sequence-dependent production of potentially harmful or useful cytokines. *Eur J Immunol.* Dec. 1997;27(12):3420-6.
- Lipford et al., Bacterial DNA as immune cell activator. *Trends Microbiol.* Dec. 1998;6(12):496-500.
- Liu et al., CpG directly induces T-bet expression and inhibits IgG1 and IgE switching in B cells. *Nat Immunol.* Jul. 2003;4(7):687-93. Epub May 25, 2003.
- Liu et al., CpG ODN is an effective adjuvant in immunization with tumor antigen. *J Invest Med.* Sep. 7, 1997;45(7):333A.
- Lonsdorf et al., Intratumor CpG-oligodeoxynucleotide injection induces protective antitumor T cell immunity. *J Immunol.* Oct. 15, 2003;171(8):3941-6.
- Lukacs et al., Interleukin-4-dependent pulmonary eosinophil infiltration in a murine model of asthma. *Am J Respir Cell Mol Biol.* May 1994;10(5):526-32.
- Lukacs et al., C-C chemokine-induced eosinophil chemotaxis during allergic airway inflammation. *J Leukoc Biol.* Nov. 1996;60(5):573-8.
- Marshall et al., Immunostimulatory sequence DNA linked to the Amb a 1 allergen promotes T(H)1 cytokine expression while downregulating T(H)2 cytokine expression in PBMCs from human patients with ragweed allergy. *J Allergy Clin Immunol.* Aug. 2001;108(2):191-7.
- Martin-Orozco et al., Enhancement of antigen-presenting cell surface molecules involved in cognate interactions by immunostimulatory DNA sequences. *Int Immunol.* Jul. 1999;11(7):1111-8.
- McCluskie et al., CpG DNA is a potent enhancer of systemic and mucosal immune responses against hepatitis B surface antigen with intranasal administration to mice. *J Immunol.* Nov. 1, 1998;161(9):4463-6.
- McCluskie et al., CpG DNA as mucosal adjuvant. *Vaccine.* 2000;18:231-7.
- McCluskie et al., Oral, intrarectal and intranasal immunizations using CpG and non-CpG oligodeoxynucleotides as adjuvants. *Vaccine.* Oct. 15, 2000;19(4-5):413-22.
- McCluskie et al., CpG DNA is an effective oral adjuvant to protein antigens in mice. *Vaccine.* Nov. 22, 2000;19(7-8):950-7.
- McCluskie et al., The use of CpG DNA as a mucosal vaccine adjuvant. *Curr Opin Investig Drugs.* Jan. 2001;2(1):35-9.
- McCluskie et al., The role of CpG in DNA vaccines. *Springer Semin Immunopathol.* 2000;22(1-2):125-32.
- Messina et al., The influence of DNA structure on the in vitro stimulation of murine lymphocytes by natural and synthetic polynucleotide antigens. *Cell Immunol.* Mar. 1993;147(1):148-57.
- Metzger et al., Oligonucleotide therapy of allergic asthma. *J Allergy Clin Immunol.* Aug. 1999;104(2 Pt 1):260-6.
- Miconnet et al., CpG are efficient adjuvants for specific CTL induction against tumor antigen-derived peptide. *J Immunol.* Feb. 1, 2002;168(3):1212-8.
- Milas et al., CpG oligodeoxynucleotide enhances tumor response to radiation. *Cancer Res.* Aug. 1, 2004;64(15):5074-7.
- Milgrom et al., Treatment of allergic asthma with monoclonal anti-IgE antibody. rhuMAB-E25 Study Group. *N Engl J Med.* Dec. 23, 1999;341(26):1966-73. Abstract Only.
- Mosmann et al., The expanding universe of T-cell subsets: Th1, Th2 and more. *Immunol Today.* Mar. 1996;17(3):138-46.
- Mutwiri et al., Biological activity of immunostimulatory CpG DNA motifs in domestic animals. *Vet Immunol Immunopathol.* Jan. 30, 2003;91(2):89-103.
- Norman et al., Immunotherapy: 1999-2004. *J Allergy Clin Immunol.* Jun. 2004;113(6):1013-23.
- Nyce et al., DNA antisense therapy for asthma in an animal model. *Nature.* Feb. 20, 1997;385(6618):721-5.
- Padrid et al., CTLA4Ig inhibits airway eosinophilia and hyperresponsiveness by regulating the development of Th1/Th2 subsets in a murine model of asthma. *Am J Respir Cell Mol Biol.* Apr. 1998;18(4):453-62.
- Park et al., The enhanced effect of a hexameric deoxyriboguanosine run conjugation to CpG oligodeoxynucleotides on protection against allergic asthma. *J Allergy Clin Immunol.* Oct. 2001;108(4):570-6.
- Parronchi et al., Phosphorothioate oligodeoxynucleotides promote the in vitro development of human allergen-specific CD4+ T cells into Th1 effectors. *J Immunol.* Dec. 1, 1999;163(11):5946-53.
- Pavlick et al., Novel therapeutic agents under investigation for malignant melanoma. *Expert Opin Investig Drugs.* Sep. 2003;12(9):1545-58.
- Payette et al., History of vaccines and positioning of current trends. *Curr Drug Targets Infect Disord.* Nov. 2001;1(3):241-7.
- Peng et al., CpG oligodeoxynucleotide vaccination suppresses IgE induction but may fail to down-regulate ongoing IgE responses in mice. *Int Immunol.* Jan. 2001;13(1):3-11.
- Pisetsky et al., The immunologic properties of DNA. *J Immunol.* Jan. 15, 1996;156(2):421-3.

- Pisetsky, Immunologic consequences of nucleic acid therapy. *Antisense Res Dev.* 1995 Fall;5(3):219-25.
- Pisetsky et al., Stimulation of in vitro proliferation of murine lymphocytes by synthetic oligodeoxynucleotides. *Mol Biol Rep.* Oct. 1993;18(3):217-21.
- Pisetsky, The influence of base sequence on the immunostimulatory properties of DNA. *Immunol Res.* 1999;19(1):35-46.
- Pisetsky et al., The influence of base sequence on the immunological properties of defined oligonucleotides. *Immunopharmacology.* Nov. 1998;40(3):199-208.
- Polanczyk et al., Immunostimulatory effects of DNA and CpG motifs. *Cent Eur J of Immunol.* 2000;25(3):160-6.
- Racila et al., Perspectives in asthma: molecular use of microbial products in asthma prevention and treatment. *J Allergy Clin Immunol.* Dec. 2005;116(6):1202-5.
- Rankin et al., CpG motif identification for veterinary and laboratory species demonstrates that sequence recognition is highly conserved. *Antisense Nucleic Acid Drug Dev.* Oct. 2001;11(5):333-40.
- Ray et al., Oral pretreatment of mice with immunostimulatory CpG DNA induces reduced susceptibility to *Listeria monocytogenes*. *Experimental Biology* 2001. Orlando, Florida, USA. Mar. 31-Apr. 4, 2001. Abstracts, part II. *FASEB J.* Mar. 8, 2001;15(5):A1007.
- Raz et al., Potential role of immunostimulatory DNA sequences (ISS) in genetic immunization and autoimmunity. *ACR Poster Session C: Cytokines and Inflammatory Mediators.* Oct. 20, 1996; Abstract 615.
- Raz et al., Preferential induction of a Th1 immune response and inhibition of specific IgE antibody formation by plasmid DNA immunization. *Proc Natl Acad Sci U S A.* May 14, 1996;93(10):5141-5.
- Readett et al., PF-3512676 (CPG7909) a Toll-like receptor 9 agonist—status of development for non-small cell lung cancer (NSCLC). Abstract PD3-1-6. *Pfizer.* Aug. 24, 2007. Poster.
- Revaz et al., The importance of mucosal immunity in defense against epithelial cancers. *Curr Opin Immunol.* Apr. 2005;17(2):175-9.
- Rhee et al., Allergen-independent immunostimulatory sequence oligodeoxynucleotide therapy attenuates experimental allergic rhinitis. *Immunology.* Sep. 2004;113(1):106-13.
- Ricci et al., T cells, cytokines, IgE and allergic airways inflammation. *J Invest Allergol Clin Immunol.* Sep.-Oct. 1994;4(5):214-20.
- Robinson et al., Predominant TH2-like bronchoalveolar T-lymphocyte population in atopic asthma. *N Engl J Med.* Jan. 30, 1992;326(5):298-304.
- Roman et al., Gene immunization for allergic disorders. *Springer Semin Immunopathol.* 1997;19(2):223-32.
- Roman et al., Immunostimulatory DNA sequences function as T helper-1-promoting adjuvants. *Nat Med.* Aug. 1997;3(8):849-54.
- Rothenfusser et al., Recent advances in immunostimulatory CpG oligonucleotides. *Curr Opin Mol Ther.* Apr. 2003;5(2):98-106.
- Roy et al., Bacterial DNA in house and farm barn dust. *J Allergy Clin Immunol.* Sep. 2003;112(3):571-8.
- Rudginsky et al., Antitumor activity of cationic lipid complexed with immunostimulatory DNA. *Mol Ther.* Oct. 2001;4(4):347-55.
- Saito et al., Allergen-induced murine upper airway inflammation: local and systemic changes in murine experimental allergic rhinitis. *Immunology.* Oct. 2001;104(2):226-34.
- Sandrasagra et al., Discovery and development of respirable antisense therapeutics for asthma. *Antisense Nucleic Acid Drug Dev.* Jun. 2002;12(3):177-81. Review.
- Santeliz et al., Amb a 1-linked CpG oligodeoxynucleotides reverse established airway hyperresponsiveness in a murine model of asthma. *J Allergy Clin Immunol.* Mar. 2002;109(3):455-62.
- Sato et al., Immunostimulatory DNA sequences necessary for effective intradermal gene immunization. *Science.* Jul. 19, 1996;273(5273):352-4.
- Satoh et al., The study of mechanisms in CpG oligodeoxynucleotides-induced aggravation in murine allergic contact dermatitis to 2,4-dinitrofluorobenzene. *Fukushima Igaku Zasshi.* 2002;52(3):237-50.
- Scheule, The role of CpG motifs in immunostimulation and gene therapy. *Adv Drug Deliv Rev.* Nov. 15, 2000;44(2-3):119-34.
- Schwartz et al., Bacterial DNA or oligonucleotides containing unmethylated CpG motifs can minimize lipopolysaccharide-induced inflammation in the lower respiratory tract through an IL-12-dependent pathway. *J Immunol.* Jul. 1, 1999;163(1):224-31.
- Schwartz et al., CpG motifs in bacterial DNA cause inflammation in the lower respiratory tract. *J Clin Invest.* Jul. 1, 1997;100(1):68-73.
- Serebrisky et al., CpG oligodeoxynucleotides can reverse Th2-associated allergic airway responses and alter the B7.1/B7.2 expression in a murine model of asthma. *J Immunol.* Nov. 15, 2000;165(10):5906-12.
- Sester et al., Phosphorothioate backbone modification modulates macrophage activation by CpG DNA. *J Immunol.* Oct. 15, 2000;165(8):4165-73.
- Shao et al., CpG-containing oligodeoxynucleotide 1826 converts the weak uveitogenic rat interphotoreceptor retinoid-binding protein peptide 1181-1191 into a strong uveitogen. *J Immunol.* Nov. 1, 2003;171(9):4780-5.
- Shiohara et al., Animal models for atopic dermatitis: are they relevant to human disease? *J Dermatol Sci.* Oct. 2004;36(1):1-9.
- Silverman et al., Immunostimulatory DNA for asthma: better than eating dirt. *Am J Respir Cell Mol Biol.* Jun. 2003;28(6):645-7.
- Simons et al., Selective immune redirection in humans with ragweed allergy by injecting Amb a 1 linked to immunostimulatory DNA. *J Allergy Clin Immunol.* Jun. 2004;113(6):1144-51.
- Sjölander et al., Iscoms containing purified *Quillaja saponins* upregulate both Th1-like and Th2-like immune responses. *Cell Immunol.* Apr. 10, 1997;177(1):69-76.
- Smit et al., Therapeutic treatment with heat-killed *Mycobacterium vaccae* (SRL172) in a mild and severe mouse model for allergic asthma. *Eur J Pharmacol.* Jun. 6, 2003;470(3):193-9.
- Sonehara et al., Hexamer palindromic oligonucleotides with 5'-CG-3' motif(s) induce production of interferon. *J Interferon Cytokine Res.* Oct. 1996;16(10):799-803.
- Sparwasser et al., Bacterial DNA causes septic shock. *Nature.* Mar. 27, 1997;386(6623):336-7.
- Sparwasser et al., Immunostimulatory CpG-oligodeoxynucleotides cause extramedullary murine hemopoiesis. *J Immunol.* Feb. 15, 1999;162(4):2368-74.
- Sparwasser et al., Macrophages sense pathogens via DNA motifs: induction of tumor necrosis factor-alpha-mediated shock. *Eur J Immunol.* Jul. 1997;27(7):1671-9.
- Spiegelberg et al., DNA-based approaches to the treatment of allergies. *Curr Opin Mol Ther.* Feb. 2002;4(1):64-71.
- Stein et al., Problems in interpretation of data derived from in vitro and in vivo use of antisense oligodeoxynucleotides. *Antisense Res Dev.* 1994 Summer;4(2):67-9.
- Stein et al., Non-antisense effects of oligodeoxynucleotides. *Antisense Technology.* 1997; ch11: 241-64.
- Stokes et al., Rationale for new treatments aimed at IgE immunomodulation. *Ann Allergy Asthma Immunol.* Sep. 2004;93(3):212-7. Abstract Only.
- Stuart et al., Marketplace strategies: The asthma challenge—Armed with a better understanding of the bad actors in the misguided immune response that causes allergies, companies hope to develop long-lasting treatments for asthma. *Start-up.* Apr. 1999; 12-20.
- Sun et al., Type I interferon-mediated stimulation of T cells by CpG DNA. *J Exp Med.* Dec. 21, 1998;188(12):2335-42.
- Sun et al., Multiple effects of immunostimulatory DNA on T cells and the role of type I interferons. *Springer Semin Immunopathol.* 2000;22(1-2):77-84.
- Sur et al., Long term prevention of allergic lung inflammation in a mouse model of asthma by CpG oligodeoxynucleotides. *J Immunol.* May 15, 1999;162(10):6284-93.
- Tam et al., Liposomal encapsulation enhances the activity of immunostimulatory oligonucleotides. *Future Lipidology.* Feb. 2006; 1(1): 35-46.
- Taube et al., Insights into the pathogenesis of asthma utilizing murine models. *Int Arch Allergy Immunol.* Oct. 2004;135(2):173-86.
- Tayyari et al., CpG-oligodeoxynucleotides inhibit RSV-enhanced allergic sensitization in guinea pigs. *Eur Respir J.* Feb. 2005;25(2):295-302.
- Threadgill et al., Mitogenic synthetic polynucleotides suppress the antibody response to a bacterial polysaccharide. *Vaccine.* Jan. 1998;16(1):76-82.
- Tighe et al., Conjugation of protein to immunostimulatory DNA results in a rapid, long-lasting and potent induction of cell-mediated and humoral immunity. *Eur J Immunol.* Jul. 2000;30(7):1939-47.

- Tighe et al., Conjugation of immunostimulatory DNA to the short ragweed allergen amb a 1 enhances its immunogenicity and reduces its allergenicity. *J Allergy Clin Immunol.* Jul. 2000;106(1 Pt 1):124-34.
- Tokunaga et al., A synthetic single-stranded DNA, poly(dG,dC), induces interferon-alpha/beta and -gamma, augments natural killer activity, and suppresses tumor growth. *Jpn J Cancer Res.* Jun. 1988;79(6):682-6.
- Tokunaga et al., Synthetic oligonucleotides with particular base sequences from the cDNA encoding proteins of *Mycobacterium bovis* BCG induce interferons and activate natural killer cells. *Microbiol Immunol.* 1992;36(1):55-66.
- Tokunaga, Response of the organism to DNA—With a focus on immunostimulatory DNA. *Kansen Ensho Meneki.* 2001 Autumn; 31(3): 1-12. Japanese.
- Tortora et al., Oral antisense that targets protein kinase a cooperates with taxol and inhibits tumor growth, angiogenesis, and growth factor production. *Clin Cancer Res.* Jun. 2000;6(6):2506-12.
- Tournoy et al., Is Th1 the solution for Th2 in asthma? *Clin Exp Allergy.* Jan. 2002;32(1):17-29.
- Uhlmann et al., Recent advances in the development of immunostimulatory oligonucleotides. *Curr Opin Drug Discov Devel.* Mar. 2003;6(2):204-17.
- Van Uden et al., Immunostimulatory DNA and applications to allergic disease. *J Allergy Clin Immunol.* Nov. 1999;104(5):902-10.
- Verthelyi et al., Human peripheral blood cells differentially recognize and respond to two distinct CPG motifs. *J Immunol.* Feb. 15, 2001;166(4):2372-7.
- Vicari et al., Reversal of tumor-induced dendritic cell paralysis by CpG immunostimulatory oligonucleotide and anti-interleukin 10 receptor antibody. *J Exp Med.* Aug. 19, 2002;196(4):541-9.
- Vollmer et al., Highly immunostimulatory CpG-free oligodeoxynucleotides for activation of human leukocytes. *Antisense Nucleic Acid Drug Dev.* Jun. 2002;12(3):165-75.
- Vollmer et al., Identification of a new class of CpG oligonucleotides capable of inducing both B cell proliferation and high IFN-alpha secretion from PBMC of HCV chronic carriers. *Antiv Ther.* 2002;7:L115.
- Vollmer et al., Characterization of three CpG oligodeoxynucleotide classes with distinct immunostimulatory activities. *Eur J Immunol.* Jan. 2004;34(1):251-62.
- Vollmer et al., Modulation of CpG oligodeoxynucleotide-mediated immune stimulation by locked nucleic acid (LNA) oligonucleotides. 2004 Spring;14(1):23-31.
- Vollmer, TLR9 in health and disease. *Int Rev Immunol.* May-Aug. 2006;25(3-4):155-81.
- Wagner et al., CpG motifs are efficient adjuvants for genetic vaccines to induce antigen-specific protective anti-tumor T cell responses. *Immunobiology* 2000;203:429. Abstract R46.
- Wagner, Interactions between bacterial CpG-DNA and TLR9 bridge innate and adaptive immunity. *Curr Opin Microbiol.* Feb. 2002;5(1):62-9.
- Wang et al., CpG oligodeoxynucleotides inhibit tumor growth and reverse the immunosuppression caused by the therapy with 5-fluorouracil in murine hepatoma. *World J Gastroenterol.* Feb. 28, 2005;11(8):1220-4.
- Weeratna et al., Reduction of antigen expression from DNA vaccines by coadministered oligodeoxynucleotides. *Antisense Nucleic Acid Drug Dev.* Aug. 1998;8(4):351-6.
- Weeratna et al., CpG ODN can re-direct the Th bias of established Th2 immune responses in adult and young mice. *FEMS Immunol Med Microbiol.* Dec. 2001;32(1):65-71.
- Weeratna et al., CpG DNA induces stronger immune responses with less toxicity than other adjuvants. *Vaccine.* Mar. 6, 2000;18(17):1755-62.
- Weiner et al., The immunobiology and clinical potential of immunostimulatory CpG oligodeoxynucleotides. *J Leukoc Biol.* Oct. 2000;68(4):455-63.
- Weiner et al., Immunostimulatory oligodeoxynucleotides containing the CpG motif are effective as immune adjuvants in tumor antigen immunization. *Proc Natl Acad Sci U S A.* Sep. 30, 1997;94(20):10833-7.
- Weiss et al., Design of protective and therapeutic DNA vaccines for the treatment of allergic diseases. *Curr Drug Targets Inflamm Allergy.* Oct. 2005;4(5):585-97.
- Wernette et al., CpG oligodeoxynucleotides stimulate canine and feline immune cell proliferation. *Vet Immunol Immunopathol.* Jan. 15, 2002;84(3-4):223-36.
- Whitmore et al., Systemic administration of LPD prepared with CpG oligonucleotides inhibits the growth of established pulmonary metastases by stimulating innate and acquired antitumor immune responses. *Canc Immun Immunother.* 2001;50:503-14.
- Whitmore et al., LPD lipopolyplex initiates a potent cytokine response and inhibits tumor growth. *Gene Ther.* 1999;6:1867-75.
- Wilson et al., Immune mechanisms and therapeutic potential of CpG oligodeoxynucleotides. *Int Rev Immunol.* May-Aug. 2006;25(3-4):183-213. Review.
- Wohlleben et al., Atopic disorders: a vaccine around the corner? *Trends Immunol.* Nov. 2001;22(11):618-26.
- Yamamoto et al., Lipofection of synthetic oligodeoxyribonucleotide having a palindromic sequence of AACGTT to murine splenocytes enhances interferon production and natural killer activity. *Microbiol Immunol.* 1994;38(10):831-6.
- Yamamoto et al., Unique palindromic sequences in synthetic oligonucleotides are required to induce IFN [correction of INF] and augment IFN-mediated [correction of INF] natural killer activity. *J Immunol.* Jun. 15, 1992;148(12):4072-6.
- Yamamoto et al., [Commemorative lecture of receiving Imamura Memorial Prize. II. Mode of action of oligonucleotide fraction extracted from *Mycobacterium bovis* BCG] *Kekkaku.* Sep. 1994;69(9):571-4. Japanese.
- Yamamoto et al., Ability of oligonucleotides with certain palindromes to induce interferon production and augment natural killer cell activity is associated with their base length. *Antisense Res Dev.* 1994 Summer;4(2):119-22.
- Yamamoto et al., Synthetic oligonucleotides with certain palindromes stimulate interferon production of human peripheral blood lymphocytes in vitro. *Jpn J Cancer Res.* Aug. 1994;85(8):775-9. Abstract Only.
- Yamamoto, Cytokine production inducing action of oligo DNA. *Rinsho Meneki.* 1997; 29(9): 1178-84. Japanese.
- Yi et al., Rapid immune activation by CpG motifs in bacterial DNA. Systemic induction of IL-6 transcription through an antioxidant-sensitive pathway. *J Immunol.* Dec. 15, 1996;157(12):5394-402.
- Yi et al., Rapid induction of mitogen-activated protein kinases by immune stimulatory CpG DNA. *J Immunol.* Nov. 1, 1998;161(9):4493-7.
- Yi et al., CpG oligodeoxyribonucleotides rescue mature spleen B cells from spontaneous apoptosis and promote cell cycle entry. *J Immunol.* Jun. 15, 1998;160(12):5898-906.
- Yi et al., IFN-gamma promotes IL-6 and IgM secretion in response to CpG motifs in bacterial DNA and oligodeoxynucleotides. *J Immunol.* Jan. 15, 1996;156(2):558-64.
- Youn et al., Immunostimulatory DNA reverses established allergen-induced airway remodeling. *J Immunol.* Dec. 15, 2004;173(12):7556-64.
- Zhao et al., Pattern and kinetics of cytokine production following administration of phosphorothioate oligonucleotides in mice. *Antisense Nucleic Acid Drug Dev.* Oct. 1997;7(5):495-502.
- Zhu et al., Modulation of ovalbumin-induced Th2 responses by second-generation immunomodulatory oligonucleotides in mice. *Int Immunopharmacol.* Jul. 2004;4(7):851-62.
- Zimmermann et al., CpG oligodeoxynucleotides trigger protective and curative Th1 responses in lethal murine leishmaniasis. *J Immunol.* Apr. 15, 1998;160(8):3627-30.
- Patent Interference No. 105,171. Iowa Preliminary Motion 3 (for judgment based on failure to comply with 35 U.S.C. 135(b)). (Electronically filed, unsigned). Jun. 7, 2004.
- Patent Interference No. 105,171. Iowa Preliminary Motion 4 (for judgment of no. interference in fact). (Electronically filed, unsigned). Jun. 7, 2004.
- Patent Interference No. 105,171. Iowa Preliminary Motion 5 (for judgment based on lack of enablement). (Electronically filed, unsigned). Jun. 7, 2004.

- Patent Interference No. 105,171. Iowa Preliminary Motion 6 (for judgment based on lack of adequate written description). (Electronically filed, unsigned). Jun. 7, 2004.
- Patent Interference No. 105,171. Iowa Preliminary Motion 7 (motion to redefine interference to designate claims as not corresponding to the Count). (Electronically filed, unsigned). Jun. 7, 2004.
- Patent Interference No. 105,171. Iowa Preliminary Motion 8 (contingent motion to redefine the Count). (Electronically filed, unsigned). Jun. 7, 2004.
- Patent Interference No. 105,171. Iowa Preliminary Motion 9 (motion for benefit of earlier application). (Electronically filed, unsigned). Jun. 7, 2004.
- Patent Interference No. 105,171. Iowa Preliminary Motion 10 (contingent motion to redefine the interference by adding a continuation application). (Electronically filed, unsigned). Jul. 2, 2004.
- Patent Interference No. 105,171. Regents of the University of California Opposition 3 (to Iowa Preliminary Motion 3 for judgment under 35 USC 135(b)). Sep. 9, 2004.
- Patent Interference No. 105,171. Regents of the University of California Opposition 4 (to Iowa Preliminary Motion 4 for judgment of no interference in fact). Sep. 9, 2004.
- Patent Interference No. 105,171. Regents of the University of California Opposition 5 (to Iowa Preliminary Motion 5 for judgment that UC's claim is not enabled). Sep. 9, 2004.
- Patent Interference No. 105,171. Regents of the University of California Opposition 6 (to Iowa Preliminary Motion 6 for judgment based on lack of adequate written description). Sep. 9, 2004.
- Patent Interference No. 105,171. Regents of the University of California Opposition 7 (to Iowa Preliminary Motion 7 to redefine the interference). Sep. 9, 2004.
- Patent Interference No. 105,171. Regents of the University of California Opposition 8 (to Iowa Preliminary Motion 8 to redefine the Count). Sep. 9, 2004.
- Patent Interference No. 105,171. Regents of the University of California Response 9 (to Iowa Contingent Motion 9 for benefit). Sep. 9, 2004.
- Patent Interference No. 105,171. Regents of the University of California Opposition 10 (to Iowa Contingent Motion 10 to redefine the interference). Sep. 9, 2004.
- Patent Interference No. 105,171. Regents of the University of California Opposition 11 (to Iowa Contingent Motion 11 to suppress). Oct. 15, 2004.
- Patent Interference No. 105,171. Iowa Reply 3 (in support of Iowa Preliminary Motion 3 for judgment under 35 U.S.C. §135(b)) (Electronically filed, unsigned). Oct. 15, 2004.
- Patent Interference No. 105,171. Iowa Reply 4 (in support of Iowa Preliminary Motion for judgment of no interference in fact) (Electronically filed, unsigned). Oct. 15, 2004.
- Patent Interference No. 105,171. Iowa Reply 5 (in support of Iowa Preliminary Motion 5 for judgment that UC's claim 205 is not enabled) (Electronically filed, unsigned). Oct. 15, 2004.
- Patent Interference No. 105,171. Iowa Reply 6 (in support of Iowa Preliminary Motion 6 for judgment based on lack of adequate written description) (Electronically filed, unsigned). Oct. 15, 2004.
- Patent Interference No. 105,171. Iowa Reply 7 (in support of Iowa Preliminary Motion 7 to redefine the interference) (Electronically filed, unsigned). Oct. 15, 2004.
- Patent Interference No. 105,171. Iowa Reply 8 (in support of Iowa Preliminary Motion 8 to redefine the count) (Electronically filed, unsigned). Oct. 15, 2004.
- Patent Interference No. 105,171. Iowa Reply 10 (in support of Iowa Preliminary Motion 10 to redefine the interference) (Electronically filed, unsigned). Oct. 15, 2004.
- Patent Interference No. 105,171. Iowa Reply 11 (in support of Iowa Miscellaneous Motion to suppress). (Electronically filed, unsigned). Oct. 18, 2004.
- Patent Interference No. 105,171. Regents of the University of California Preliminary Statement. Jun. 7, 2004.
- Patent Interference No. 105,171. Regents of the University of California Preliminary Motion 1 (to designate additional claims of Iowa patent as corresponding to the Count). Jun. 7, 2004.
- Patent Interference No. 105,171. Regents of the University of California Preliminary Motion 2 (for judgment based on lack of written description support and introducing new matter). Jun. 7, 2004.
- Patent Interference No. 105,171. Regents of the University of California Preliminary Motion 3 (for judgment based on anticipation). Jun. 7, 2004.
- Patent Interference No. 105,171. Regents of the University of California Preliminary Motion 4 (for judgment based on obviousness). Jun. 7, 2004.
- Patent Interference No. 105,171. Regents of the University of California Preliminary Motion 5 (for judgment based on anticipation). Jun. 7, 2004.
- Patent Interference No. 105,171. Regents of the University of California Preliminary Motion 6 (for judgment based on inequitable conduct). Jun. 7, 2004.
- Patent Interference No. 105,171. Regents of the University of California Contingent Preliminary Motion 7 (for benefit of an earlier application under 37 CFR 1.633(j)). Jul. 2, 2004.
- Patent Interference No. 105,171. Regents of the University of California Contingent Preliminary Motion 8 (to add additional claims under 37 CFR 1.633(c)(2) and (i)). Jul. 2, 2004.
- Amended Claims for U.S. Appl. No. 09/265,191, filed Mar. 10, 1999.
- Patent Interference No. 105,171. Iowa Opposition 1 (opposition to motion to designate additional claims as corresponding to the Count) (Electronically filed, unsigned). Sep. 9, 2004.
- Patent Interference No. 105,171. Iowa Opposition 2 (opposition to motion for judgment based on lack of written description support and introducing new matter) (Electronically filed, unsigned). Sep. 9, 2004.
- Patent Interference No. 105,171. Iowa Opposition 3 (opposition to motion for judgment based on anticipation) (Electronically filed, unsigned). Sep. 9, 2004.
- Patent Interference No. 105,171. Iowa Opposition 4 (opposition to motion for judgment based on obviousness) (Electronically filed, unsigned). Sep. 9, 2004.
- Patent Interference No. 105,171. Iowa Opposition 5 (opposition to motion for judgment based on anticipation) (Electronically filed, unsigned). Sep. 9, 2004.
- Patent Interference No. 105,171. Iowa Opposition 6 (opposition to motion for judgment based on inequitable conduct) (Electronically filed, unsigned). Sep. 9, 2004.
- Patent Interference No. 105,171. Iowa Opposition 7 (opposition to motion for benefit of an earlier application under 7 CFR 1.633(j)) (Electronically filed, unsigned). Sep. 9, 2004.
- Patent Interference No. 105,171. Iowa Opposition 8 (opposition to motion to add additional claims under 37 CFR 1.633 (2) and (i)) (Electronically filed, unsigned). Sep. 9, 2004.
- Patent Interference No. 105,171. Regents of the University of California Reply 1 (to Iowa's opposition to UC's motion to designate Iowa claims as corresponding to the Count). Oct. 15, 2004.
- Patent Interference No. 105,171. Regents of the University of California Reply 2 (to Iowa's opposition to UC Preliminary Motion 2 for Judgment). Oct. 15, 2004.
- Patent Interference No. 105,171. Regents of the University of California Reply 3 (to Iowa's Opposition to UC Preliminary Motion 3 for Judgment). Oct. 15, 2004.
- Patent Interference No. 105,171. Regents of the University of California Reply 4 (to Iowa's Opposition to UC Preliminary Motion 4 for Judgment). Oct. 15, 2004.
- Patent Interference No. 105,171. Regents of the University of California Reply 5 (to Iowa's Opposition to UC Preliminary Motion 5 for Judgment). Oct. 15, 2004.
- Patent Interference No. 105,171. Regents of the University of California Reply 6 (to Iowa's opposition to UC Preliminary Motion 6 for judgment). Oct. 15, 2004.
- Patent Interference No. 105,171. Regents of the University of California Reply 7 (to Iowa's Opposition to UC Preliminary Motion 7 for Benefit). Oct. 15, 2004.
- Patent Interference No. 105,171. Regents of the University of California Reply 8 (to Iowa's Opposition to UC Preliminary Motion 8 to add additional claims). Oct. 15, 2004.
- Patent Interference No. 105,171. Decision on Motion under 37 CFR §41.125. Mar. 10, 2005.

- Patent Interference No. 105,171. Judgment and Order. Mar. 10, 2005.
- Patent Interference No. 105,171. Regents of the University of California. Brief of Appellant. Jul. 5, 2005.
- Patent Interference No. 105,171. University of Iowa and Coley Pharmaceutical Group, Inc. Brief of Appellees. Aug. 17, 2005.
- Patent Interference No. 105,171. Regents of the University of California. Reply Brief of Appellant. Sep. 6, 2005.
- Patent Interference No. 105,171. Regents of the University of California. Decision of CAFC. Jul. 17, 2006.
- Patent Interference No. 105,526. Krieg Substantive Motion 1 (for unpatentability based on interference estoppel). (Electronically filed, unsigned).
- Patent Interference No. 105,526. Krieg Substantive Motion 2 (for judgment based on inadequate written description and/or enablement). (Electronically filed, unsigned). Jun. 18, 2007.
- Patent Interference No. 105,526. Krieg Contingent Responsive Motion (to add new claims 104 and 105). (Electronically filed, unsigned). Jul. 25, 2007.
- Patent Interference No. 105,526. Krieg Substantive Motion 3 (for judgment based on prior art). (Electronically filed, unsigned). Jun. 18, 2007.
- Patent Interference No. 105,526. Raz Motion 1 (Unpatentability of Krieg Claims under 35 U.S.C. § 112, First Paragraph). (Electronically filed, unsigned). Jun. 18, 2007.
- Patent Interference No. 105,526. Raz Motion 2 (Raising a Threshold Issue of No Interference-in-Fact). (Electronically filed, unsigned). Jun. 18, 2007.
- Patent Interference No. 105,526. Raz Motion 3 (Krieg's Claims are Unpatentable Over Prior Art Under 35 U.S.C. § 102(b)). (Electronically filed, unsigned). Jun. 18, 2007.
- Patent Interference No. 105,526. Raz Motion 4 (to Designate Krieg Claims 46 and 82-84 as Corresponding to Count 1). (Electronically filed, unsigned). Jun. 18, 2007.
- Patent Interference No. 105,526. Raz Responsive Miscellaneous Motion 5 (to revive the Raz Parent Application) (Electronically filed, unsigned) Jul. 25, 2007.
- Patent Interference No. 105,526. Raz Contingent Responsive Motion 6 (To Add a New Claim 58) (Electronically filed, unsigned) Jul. 25, 2007.
- Patent Interference No. 105,526. Krieg Opposition 1 (Opposition to Motion for Lack of Enablement and Written Description) (Electronically filed, unsigned) Sep. 10, 2007.
- Patent Interference No. 105,526. Krieg Opposition 2 (to Raz Motion 2) (Electronically filed, unsigned) Sep. 10, 2007.
- Patent Interference No. 105,526. Krieg Opposition 3 (To Raz Motion 3) (Electronically filed, unsigned) Sep. 10, 2007.
- Patent Interference No. 105,526. Krieg Opposition 4 (Opposition to Motion for Designating Claims 46 and 82-84 as Corresponding to the Court) (Electronically filed, unsigned) Sep. 10, 2007.
- Patent Interference No. 105,526. Krieg Opposition 6 (Opposition to Raz Contingent Responsive Motion 6) (Electronically filed, unsigned) Sep. 10, 2007.
- Patent Interference No. 105,526. Raz Opposition 1 (Opposing Krieg Substantive Motion 1) (Electronically filed, unsigned) Sep. 10, 2007.
- Patent Interference No. 105,526. Raz Opposition 2 (Opposing Krieg Substantive Motion 2) (Electronically filed, unsigned) Sep. 10, 2007.
- Patent Interference No. 105,526. Raz Opposition 4 (Opposing Krieg Contingent Responsive Motion to Add New Claims 104 and 105) (Electronically filed, unsigned) Sep. 10, 2007.
- Patent Interference No. 105,526. Krieg Reply 1 (Reply to Raz opposition 1) Oct. 5, 2007.
- Patent Interference No. 105,526. Krieg Reply 2 (Reply to Raz opposition 2) Oct. 5, 2007.
- Patent Interference No. 105,526. Krieg Reply 4 (Reply to Raz opposition 4) Oct. 5, 2007.
- Patent Interference No. 105,526. Raz Reply 1 (Reply to Krieg opposition 1) Oct. 5, 2007.
- Patent Interference No. 105,526. Raz Reply 2 (Reply to Krieg opposition 2) Oct. 5, 2007.
- Patent Interference No. 105,526. Raz Reply 3 (Reply to Krieg opposition 3) Oct. 5, 2007.
- Patent Interference No. 105,526. Raz Reply 4 (Reply to Krieg opposition 4) Oct. 5, 2007.
- Patent Interference No. 105,526. Raz Reply 6 (Reply to Krieg opposition 6) Oct. 5, 2007.
- Patent Interference No. 105,526. Krieg Miscellaneous Motion 5 (To exclude exhibits 2066, 2070, 2071, 2072, 2073, 2074, 2075, 2076 and 2078) Oct. 9, 2007.
- Patent Interference No. 105,526. Raz Opposition 5 (Opposing Krieg Miscellaneous Motion 5) Oct. 25, 2007.
- Patent Interference No. 105,526. Raz Miscellaneous Motion 7 (To exclude evidence) Oct. 19, 2007.
- Patent Interference No. 105,526. Krieg Opposition 7 (To Raz Miscellaneous Motion 7) Oct. 25, 2007.
- Patent Interference No. 105,526. Krieg Reply 5 (Reply to Raz opposition 5) Oct. 30, 2007.
- Patent Interference No. 105,526. Raz Reply 7 (Reply to Krieg opposition 7) Oct. 30, 2007.
- Patent Interference No. 105,526. Order—Bd.R. 104. Conference Call. Paper 211. Sep. 30, 2008.
- Patent Interference No. 105,526. Memorandum Opinion and Order (Decision on Motions) Dec. 1, 2008.
- Patent Interference No. 105,526. Judgement on Preliminary Motions under 37 C.F.R. §41.127 Dec. 1, 2008.
- Patent Interference No. 105,526. Paper 217. Raz Notice of Filing of a Notice of Appeal (Appeal to the Court of Appeals for the Federal Circuit). Jan. 27, 2009.
- Patent Interference No. 105,526. Paper 219. Raz Notice of Withdrawal of Appeal. May 15, 2009.
- Patent Interference No. 105,674. Paper No. 1. Declaration under 37 C.F.R. §41.203(b) Dec. 1, 2008.
- Patent Interference No. 105,674. Paper No. 6 Raz Notice of Real Party in Interest. Dec. 12, 2008.
- Patent Interference No. 105,674. Paper No. 11 Krieg Designation of Real Party in Interest. Dec. 15, 2008.
- Patent Interference No. 105,674. Paper No. 15. Order—Bd.R. 104(c) Summary of Dec. 23, 2008 Conference Call.
- Patent Interference No. 105,674. Paper No. 19. Order—Bd.R. 104(c). Conference Call. Jan. 16, 2009.
- Patent Interference No. 105,674. Paper No. 21. Raz Observations (regarding evidence to support certain proposed motions. Jan. 27, 2009.
- Patent Interference No. 105,674. Paper No. 23. Raz Miscellaneous Motion 1 (to revive the Raz parent application). Jan. 27, 2009.
- Patent Interference No. 105,674. Paper No. 25. Order—Bd.R. 104(c) (Raz v. Krieg) Summary of Conference Call on Feb. 4, 2009.
- Patent Interference No. 105,674. Paper No. 29. Joint Submission Pursuant to Order Dated Jan. 16, 2009. Mar. 11, 2009.
- Patent Interference No. 105,674. Paper No. 32. Raz Abandonment of Contest. May 15, 2009.
- Patent Interference No. 105,674. Paper No. 33. Judgment—Bd.R. 127. May 20, 2009.
- [No Author Listed] Allergen Nomenclature, List of Allergens (as of Mar. 12, 2004).
- [No Author Listed] Definitions. in *Dorland's Illustrated Medical Dictionary*, Twenty-sixth Edition. W.B. Saunders Company, 1981. Exhibit 2032.
- [No Author Listed] Definitions. in *Immunology*, Roitt et al., Editors. Gower Medical Publishing Company, 1985. Exhibit 2035.
- [No Author Listed] Definitions. in the *Dictionary of Immunology*, Fourth Edition. Herbert et al., editors. Academic Press, 1995. p. 10. Exhibit 1004.
- [No Author Listed] Definitions. in *Webster's New Collegiate Dictionary*. The G&C Merriam Company, 1981. p. 287. Exhibit 2070.
- [No Author Listed] DNA is the primary genetic material. in *Recombinant DNA*, Second Edition. Watson et al., editors. Scientific American Books, New York. p. 24-5, 147. Exhibit 1003.
- [No Author Listed] <http://www.cancer.gov/clinicaltrials/learning/cancervaccines/print>. Exhibit 1043.
- [No Author Listed] <http://www.cancer.gov/newscenter/pressreleases/cancervaccines/print>. Exhibit 1042.
- [No Author Listed] http://www.cancer.org/docroot/ETO/content/ETO_1_4X_Cancer_Vaccines_Active_Specific_Immun_therapies.asp. Exhibit 2072.
- [No Author Listed] http://www.doctorfungus.com/the_fungi/Alternaria.htm. Exhibit 1031.

- [No Author Listed] http://www.doctorfungus.com/the_fungi/Blastomyces.htm. Exhibit 1030.
- [No Author Listed] http://www.doctorfungus.com/the_fungi/Cryptococcus.htm. Exhibit 1029.
- [No Author Listed] <http://www.icare.org/treatment/tcwtv4.htm>. Exhibit 2074.
- [No Author Listed] <http://www.plwc.org/plwc/mainconstructor>. Exhibit 2073.
- [No Author Listed] Macromolecules in Prokaryotic and Eukaryotic Cells. in Molecular Cell Biology, Darnell et al, Editors. W.H. Freeman and Company, 1986. Exhibit 2033.
- [No Author Listed] Recombinant DNA Technology. in Molecular Biology of the Cell. Alberts et al., editors. Garland Publishing, Inc. 1983. Exhibit 2027.
- Abed et al., Interferon-gamma regulation of B lymphocyte differentiation: activation of B cells is a prerequisite for IFN-gamma-mediated inhibition of B cell differentiation. *Cell Immunol.* Feb. 1994;153(2):356-66. Exhibit 2052.
- Adya et al., Expansion of CREB's DNA recognition specificity by Tax results from interaction with Ala-Ala-Arg at positions 282-284 near the conserved DNA-binding domain of CREB. *Proc Natl Acad Sci U S A.* Jun. 7, 1994;91(12):5642-6.
- Agrawal et al., Antisense therapeutics: is it as simple as complementary base recognition? *Mol Med Today.* Feb. 2000;6(2):72-81.
- Agrawal et al., Novel immunomodulatory oligonucleotides prevent development of allergic airway inflammation and airway hyper-responsiveness in asthma. *Int Immunopharmacol.* Jan. 2004;4(1):127-38.
- Agrawal et al., Pharmacokinetics, biodistribution, and stability of oligodeoxynucleotide phosphorothioates in mice. *Proc Natl Acad Sci U S A.* Sep. 1, 1991;88(17):7595-9.
- Alm et al., Early BCG vaccination and development of atopy. *Lancet.* Aug. 9, 1997;350(9075):400-3.
- Angier et al., Microbe DNA seen as alien by immune cells. *New York Times (Science).* Apr. 11, 1995;B5, B9.
- Azad et al., Antiviral activity of a phosphorothioate oligonucleotide complementary to RNA of the human cytomegalovirus major immediate-early region. *Antimicrob Agents Chemother.* Sep. 1993;37(9):1945-54.
- Azuma et al., Biochemical and Immunological Studies on Cellular Components of Tubercle Bacilli. *Kekkaku.* 1992; 69(9):45-65.
- Bayever et al., Systemic administration of a phosphorothioate oligonucleotide with a sequence complementary to p53 for acute myelogenous leukemia and myelodysplastic syndrome: initial results of a phase I trial. *Antisense Res Dev.* 1993 Winter;3(4):383-90.
- Bennett et al., DNA binding to human leukocytes. Evidence for a receptor-mediated association, internalization, and degradation of DNA. *J Clin Invest.* Dec. 1985;76(6):2182-90.
- Berg et al., Interleukin-10 is a central regulator of the response to LPS in murine models of endotoxic shock and the Shwartzman reaction but not endotoxin tolerance. *J Clin Invest.* Nov. 1995;96(5):2339-47.
- Blanchard et al., Interferon-gamma induction by lipopolysaccharide: dependence on interleukin 2 and macrophages. *J Immunol.* Feb. 1, 1986;136(3):963-70.
- Blaxter et al., Genes expressed in *Brugia malayi* infective third stage larvae. *Mol Biochem Parasitol.* Apr. 1996;77(1):77-93.
- Bochner et al., Advances in mechanisms of allergy. *J Allergy Clin Immunol.* May 2004;113(5):868-75.
- Boushey et al., Targets for asthma therapy. *Allerg Immunol (Paris).* Nov. 2000;32(9):336-41 Abstract.
- Branda et al., Amplification of antibody production by phosphorothioate oligodeoxynucleotides. *J Lab Clin Med.* Sep. 1996;128(3):329-38.
- Branda et al., Immune stimulation by an antisense oligomer complementary to the rev gene of HIV-1. *Biochem Pharmacol.* May 25, 1993;45(10):2037-43.
- Briskin et al., Lipopolysaccharide-unresponsive mutant pre-B-cell lines blocked in NF-kappa B activation. *Mol Cell Biol.* Jan. 1990;10(1):422-5.
- Capron et al., Immunologic aspects of schistosomiasis. *Annu Rev Med.* 1992;43:209-18. Review. Exhibit 2068.
- Chace et al., Regulation of differentiation in CD5+ and conventional B cells. Sensitivity to LPS-induced differentiation and interferon-gamma-mediated inhibition of differentiation. *Clin Immunol Immunopathol.* Sep. 1993;68(3):327-32.
- Chang et al., The effect of CpG-oligodeoxynucleotides with different backbone structures and 3' hexameric deoxyriboguanosine run conjugation on the treatment of asthma in mice. *J Allergy Clin Immunol.* 2004;113(2):S323. Abstract 1196.
- Chang et al., The palindromic series I repeats in the simian cytomegalovirus major immediate-early promoter behave as both strong basal enhancers and cyclic AMP response elements. *J Virol.* Jan. 1990;64(1):264-77.
- Chiang et al., Ribavirin or CpG DNA sequence-modulated dendritic cells decrease the IgE level and airway inflammation. *Am J Respir Crit Care Med.* Sep. 1, 2003;168(5):575-80.
- Choudhury et al., In vivo role of p38 mitogen-activated protein kinase in mediating the anti-inflammatory effects of CpG oligodeoxynucleotide in murine asthma. *J Immunol.* Nov. 15, 2002;169(10):5955-61.
- Cockcroft et al., Comparative effects of inhaled salbutamol, sodium cromoglycate, and beclomethasone dipropionate on allergen-induced early asthmatic responses, late asthmatic responses, and increased bronchial responsiveness to histamine. *J Allergy Clin Immunol.* May 1987;79(5):734-40.
- Cooper et al., CPG 7909, an immunostimulatory TLR9 agonist oligodeoxynucleotide, as adjuvant to Engerix-B HBV vaccine in healthy adults: a double-blind phase I/II study. *J Clin Immunol.* Nov. 2004;24(6):693-701.
- Croft et al., Generation of polarized antigen-specific CD8 effector populations: reciprocal action of interleukin (IL)-4 and IL-12 in promoting type 2 versus type 1 cytokine profiles. *J Exp Med.* Nov. 1, 1994;180(5):1715-28.
- Crosby et al., The early response gene NGFI-C encodes a zinc finger transcriptional activator and is a member of the GCGGGGCG (GSG) element-binding protein family. *Mol Cell Biol.* Aug. 1991;11(8):3835-41.
- Crystal et al., Transfer of genes to humans: early lessons and obstacles to success. *Science.* Oct. 20, 1995;270(5235):404-10.
- D'Andrea et al., Interleukin 10 (IL-10) inhibits human lymphocyte interferon gamma-production by suppressing natural killer cell stimulatory factor/IL-12 synthesis in accessory cells. *J Exp Med.* Sep. 1, 1993;178(3):1041-8.
- Dunn et al., The three Es of cancer immunoediting. *Annu Rev Immunol.* 2004;22:329-60.
- Englisch et al., Chemically modified oligonucleotides as probes and inhibitors. *Angew. Chem. Int. Ed. Eng.* Jun. 1991;30(6):613-29.
- Erb et al., Infection of mice with *Mycobacterium bovis*-Bacillus Calmette-Guerin (BCG) suppresses allergen-induced airway eosinophilia. *J Exp Med.* Feb. 16, 1998;187(4):561-9.
- Etlinger et al., Carrier sequence selection—one key to successful vaccines. *Immunol Today.* Feb. 1992;13(2):52-5.
- Feleszko et al., Toll-like receptors—novel targets in allergic airway disease (probiotics, friends and relatives). *Eur J Pharmacol.* Mar. 8, 2006;533(1-3):308-18.
- Fitch et al., The effect of an oral phosphodiesterase (PDE) 4 inhibitor bay 19-8004 in primate asthma models. *American Thoracic Society.* 2000. A52 Poster A94.
- Fox R.I., Mechanism of action of hydroxychloroquine as an antirheumatic drug. *Chemical Abstracts,* 120:15, Abstract No. 182630 (Apr. 29, 1994).
- Francois et al., Examination of the inhibitory and stimulatory effects of IFN-alpha, -beta, and -gamma on human B-cell proliferation induced by various B-cell mitogens. *Clin Immunol Immunopathol.* Sep. 1988;48(3):297-306.
- Friedberg et al., Combination immunotherapy with a CpG oligonucleotide (1018 ISS) and rituximab in patients with non-Hodgkin lymphoma: increased interferon-alpha/beta-inducible gene expression, without significant toxicity. *Blood.* Jan. 15, 2005;105(2):489-95. Epub Sep. 9, 2004.
- Frizzera et al., IFN-gamma-mediated inhibition of antigen receptor-induced B cell proliferation and CREB-1 binding activity requires STAT-1 transcription factor. *Eur J Immunol.* Apr. 2003;33(4):907-12.
- Gibson et al., Cellular requirements for cytokine production in response to the immunomodulators imiquimod and S-27609. *J Interferon Cytokine Res.* Jun. 1995;15(6):537-45.

- Grunig, G. et al., Molecular regulation of Th2 immunity by dendritic cells. *Pharmacol Ther.* Apr. 2005;106(1):75-96. Epub Dec. 20, 2004. 22 pages.
- Gundel et al., Antigen-induced mediator release in primates. *Am Rev Respir Dis.* Jul. 1991;144(1):76-82.
- Gundel et al., Repeated antigen inhalation results in a prolonged airway eosinophilia and airway hyperresponsiveness in primates. *J Appl Physiol.* Feb. 1990;68(2):779-86.
- Gura et al., Antisense has growing pains. *Science.* Oct. 27, 1995;270(5236):575-6.
- Hadden et al., Immunopharmacology. Immunomodulation and immunotherapy. *JAMA.* Nov. 25, 1992;268(20):2964-9.
- Hadden et al., Immunostimulants. *Trends Pharmacol Sci.* May 1993;14(5):169-74.
- Hatzfeld et al., Release of early human hematopoietic progenitors from quiescence by antisense transforming growth factor beta 1 or Rb oligonucleotides. *J Exp Med.* Oct. 1, 1991;174(4):925-9.
- Hemmi et al., A Toll-like receptor recognizes bacterial DNA. *Nature.* Dec. 7, 2000;408(6813):740-5.
- Hemmi et al., The roles of Toll-like receptor 9, MyD88, and DNA-dependent protein kinase catalytic subunit in the effects of two distinct CpG DNAs on dendritic cell subsets. *J Immunol.* Mar. 15, 2003;170(6):3059-64.
- Herz et al., BCG infection suppresses allergic sensitization and development of increased airway reactivity in an animal model. *J Allergy Clin Immunol.* Nov. 1998;102(5):867-74.
- Highfield et al., Sepsis: the more, the murkier. *Biotechnology (NY).* Aug. 1994;12(8):828.
- Hoeffler et al., Identification of multiple nuclear factors that interact with cyclic adenosine 3',5'-monophosphate response element-binding protein and activating transcription factor-2 by protein-protein interactions. *Mol Endocrinol.* Feb. 1991;5(2):256-66.
- Howarth et al., Influence of albuterol, cromolyn sodium and ipratropium bromide on the airway and circulating mediator responses to allergen bronchial provocation in asthma. *Am Rev Respir Dis.* Nov. 1985;132(5):986-92.
- Humlova et al., [Bacteria and their role in allergic diseases] *Cas Lek Cesk.* 2004;143(1):21-5. Review. Czech.
- Iguchi-Arigo et al., CpG methylation of the cAMP-responsive enhancer/promoter sequence TGACGTC A abolishes specific factor binding as well as transcriptional activation. *Genes Dev.* May 1989;3(5):612-9.
- Infante-Duarte et al., Th1/Th2 balance in infection. *Springer Semin Immunopathol.* 1999;21(3):317-38.
- Ishikawa et al., IFN induction and associated changes in splenic leukocyte distribution. *J Immunol.* May 1, 1993;150(9):3713-27.
- Iversen et al., Pharmacokinetics of an antisense phosphorothioate oligodeoxynucleotide against rev from human immunodeficiency virus type 1 in the adult male rat following single injections and continuous infusion. *Antisense Res Dev.* 1994 Spring;4(1):43-52.
- Jakway et al., Growth regulation of the B lymphoma cell line WEHI-231 by anti-immunoglobulin, lipopolysaccharide, and other bacterial products. *J Immunol.* Oct. 1, 1986;137(7):2225-31.
- Jaroszewski et al., Cellular uptake of antisense oligodeoxynucleotides. *Advanced Drug Delivery Reviews.* 1991;6:235-50.
- Jones et al., Pharmacology of montelukast sodium (Singulair), a potent and selective leukotriene D4 receptor antagonist. *Can J Physiol Pharmacol.* Feb. 1995;73(2):191-201.
- Kline et al., T-lymphocyte dysregulation in asthma. *Proc Soc Exp Biol Med.* Dec. 1994;207(3):243-53.
- Kou et al., [Analysis and regulation of interferon-gamma production by peripheral blood lymphocytes patients with bronchial asthma] *Aerugi.* Mar. 1994;43(3):482-91. Japanese.
- Krieg et al., A role for endogenous retroviral sequences in the regulation of lymphocyte activation. *J Immunol.* Oct. 15, 1989;143(8):2448-51.
- Krieg et al., Modification of antisense phosphodiester oligodeoxynucleotides by a 5' cholesteryl moiety increases cellular association and improves efficacy. *Proc Natl Acad Sci U S A.* Feb. 1, 1993;90(3):1048-52.
- Krieg et al., Uptake of oligodeoxyribonucleotides by lymphoid cells is heterogeneous and inducible. *Antisense Res Dev.* 1991; 1(2):161-71.
- Krown et al., Phase I trial with the interferon inducer poly(l/poly-l-lysine) (Poly ICL). *Journal of Interferon Research.* 1983; 3:281-90.
- Kuby et al., Editors, "Chapter 13: Cytokines", *Immunology: Second Edition.* 1994. W.H. Freeman and Company: New York, p. 297-322.
- Lazarus et al., Single-nucleotide polymorphisms in the Toll-like receptor 9 gene (TLR9): frequencies, pairwise linkage disequilibrium, and haplotypes in three U.S. ethnic groups and exploratory case-control disease association studies. *Genomics.* Jan. 2003;81(1):85-91.
- Leigh et al., Effects of montelukast and budesonide on airway responses and airway inflammation in asthma. *Am J Respir Crit Care Med.* Nov. 1, 2002;166(9):1212-7.
- Leonard et al., Conformation of guanine-8-oxoadenine base pairs in the crystal structure of d(CGCGAATT(O8A)GCG). *Biochemistry.* Sep. 15, 1992;31(36):8415-20.
- Litzinger et al., Fate of cationic liposomes and their complex with oligonucleotide in vivo. *Biochim Biophys Acta.* Jun. 11, 1996;1281(2):139-49.
- Liu et al., Hygiene hypothesis: fact or fiction? *J Allergy Clin Immunol.* Mar. 2003;111(3):471-8.
- Lotz et al., Effects of recombinant human interferons on rheumatoid arthritis B lymphocytes activated by Epstein-Barr virus. *J Rheumatol.* Feb. 1987;14(1):42-5. Exhibit 2053.
- Ma et al., DNA-based vaccination against hepatitis C virus (HCV): effect of expressing different forms of HCV E2 protein and use of CpG-optimized vectors in mice. *Vaccine.* Sep. 10, 2002;20(27-28):3263-71.
- Macaubas et al., Respiratory tolerance in the protection against asthma. *Curr Drug Targets Inflamm Allergy.* Jun. 2003;2(2):175-86.
- MacFarlane et al., Antagonism of immunostimulatory CpG-oligodeoxynucleotides by quinacrine, chloroquine, and structurally related compounds. *J Immunol.* Feb. 1, 1998;160(3):1122-31.
- Mardh et al., *Alternaria alternata* as a cause of opportunistic fungal infections in man. *Scand J Infect Dis Suppl.* 1978;(16):36-40.
- Martin et al., Immunotherapy as a disease modifier. *Paediatr Respir Rev.* 2006;7 Suppl 1:S106-7. Epub Jun. 5, 2006.
- Mastrangelo et al., Gene therapy for human cancer: an essay for clinicians. *Semin Oncol.* Feb. 1996;23(1):4-21.
- Matson et al., Nonspecific suppression of [3H]thymidine incorporation by "control" oligonucleotides. *Antisense Res Dev.* 1992 Winter;2(4):325-30.
- McCluskie et al., Intranasal immunization of mice with CpG DNA induces strong systemic and mucosal responses that are influenced by other mucosal adjuvants and antigen distribution. *Mol Med.* Oct. 2000;6(10):867-77.
- McCluskie et al., Route and method of delivery of DNA vaccine influence immune responses in mice and non-human primates. *Mol Med.* May 1999;5(5):287-300.
- McCluskie et al., The potential of CpG oligodeoxynucleotides as mucosal adjuvants. *Crit Rev Immunol.* 2001;21(1-3):103-20.
- McCluskie et al., The potential of oligodeoxynucleotides as mucosal and parenteral adjuvants. *Vaccine.* Mar. 21, 2001;19(17-19):2657-60.
- McIntyre et al., A sense phosphorothioate oligonucleotide directed to the initiation codon of transcription factor NF-kappa B p65 causes sequence-specific immune stimulation. *Antisense Res Dev.* 1993 Winter;3(4):309-22.
- Messina et al., Stimulation of in vitro murine lymphocyte proliferation by bacterial DNA. *J Immunol.* Sep. 15, 1991;147(6):1759-64.
- Metzger et al., Enhancement of humoral immunity by interleukin-12. *Ann NY Acad Sci.* Oct. 31, 1996;795:100-15.
- Milligan et al., Current concepts in antisense drug design. *J Med Chem.* Jul. 9, 1993;36(14):1923-37.
- Mojcik et al., Administration of a phosphorothioate oligonucleotide antisense to murine endogenous retroviral MCF env causes immune effects in vivo in a sequence-specific manner. *Clin Immunol Immunopathol.* May 1993;67(2):130-6.
- Mond et al., Recombinant interferon-gamma inhibits the B cell proliferative response stimulated by soluble but not by Sepharose-bound anti-immunoglobulin antibody. *J Immunol.* Oct. 1985;135(4):2513-7.

- Mottram et al., A novel CDC2-related protein kinase from *Leishmania mexicana*, LmmCRK1, is post-translationally regulated during the life cycle. *J Biol Chem*. 1 Oct. 5, 1993;268(28):21044-52. New England Biolabs 1988-1989 Catalog.
- Pare et al., Lung mechanics following antigen challenge of *Ascaris summ*-sensitive rhesus monkeys. *J Appl Physiol*. Nov. 1976;41(5 Pt. 1):668-76.
- Patterson et al., Inhibition of immunoglobulin E-mediated, antigen-induced monkey asthma and skin reactions by 5, 8, 11, 14-eicosatetraenoic acid. *J Allergy Clin Immunol*. Feb. 1981;67(2):146-52.
- Perez et al., Sequence-independent induction of Sp1 transcription factor activity by phosphorothioate oligodeoxynucleotides. *Proc Natl Acad Sci U S A*. Jun. 21, 1994;91(13):5957-61.
- Pisetsky et al., A role for immunogenic DNA in the pathogenesis of systemic lupus erythematosus. *Arthritis and Rheumatism*. Feb. 1990;33(2):153-159.
- Pisetsky et al., Immune activation by bacterial DNA: a new genetic code. *Immunity*. Oct. 1996;5(4):303-10.
- Pisetsky et al., Immunological properties of bacterial DNA. *Ann NY Acad Sci*. Nov. 27, 1995;772:152-63.
- Pisetsky et al., Stimulation of murine lymphocyte proliferation by a phosphorothioate oligonucleotide with antisense activity for herpes simplex virus. *Life Sci*. 1994;54(2):101-7.
- Raz et al., Intradermal gene immunization: the possible role of DNA uptake in the induction of cellular immunity to viruses. *Proc Natl Acad Sci U S A*. Sep. 27, 1994;91(20):9519-23.
- Redecke et al., Cutting edge: activation of Toll-like receptor 2 induces a Th2 immune response and promotes experimental asthma. *J Immunol*. Mar. 1, 2004;172(5):2739-43.
- Reynolds et al., Inhibition of B lymphocyte activation by interferon-gamma. *J Immunol*. Aug. 1, 1987;139(3):767-73.
- Sander et al., Sequential production of Th1 and Th2 cytokines in response to live bacillus Calmette-Guerin. *Immunology*. Dec. 1995;86(4):512-8 Abstract.
- Schnell et al., Identification and characterization of a *Saccharomyces cerevisiae* gene (PAR1) conferring resistance to iron chelators. *Eur J Biochem*. Sep. 1, 1991;200(2):487-93.
- Schwartz et al., Endotoxin responsiveness and grain dust-induced inflammation in the lower respiratory tract. *Am J Physiol*. Nov. 1994;267(5 Pt 1):L609-17.
- Schwartz et al., The role of endotoxin in grain dust-induced lung disease. *Am J Respir Crit Care Med*. Aug. 1995;152(2):603-8 Abstract.
- Shirakawa et al., The inverse association between tuberculin responses and atopic disorder. *Science*. Jan. 3, 1997;275(5296):77-9.
- Sidman et al., Gamma-interferon is one of several direct B cell-maturing lymphokines. *Nature*. Jun. 28-Jul. 4, 1984;309(5971):801-4.
- Siegrist et al., Co-administration of CpG oligonucleotides enhances the late affinity maturation process of human anti-hepatitis B vaccine response. *Vaccine*. Dec. 16, 2004;23(5):615-22.
- Silverman et al., BCG vaccination and atopy—unfinished business? *Lancet*. Aug. 9, 1997;350(9075):380-1.
- Singh et al., Cationic microparticles are an effective delivery system for immune stimulatory CpG DNA. *Pharm Res*. Oct. 2001;18(10):1476-9.
- Spiegelberg et al., Allergen-immunostimulatory oligodeoxynucleotide conjugate: a novel allergoid for immunotherapy. *Curr Opin Allergy Clin Immunol*. Dec. 2002;2(6):547-51 Abstract.
- Spiegelberg et al., Recognition of T cell epitopes and lymphokine secretion by rye grass allergen *Lolium perenne* I-specific human T cell clones. *J Immunol*. May 1, 1994;152(9):4706-11.
- Stein et al., Antisense oligonucleotides as therapeutic agents—is the bullet really magical? *Science*. Aug. 20, 1993;261(5124):1004-12.
- Stein et al., Oligodeoxynucleotides as inhibitors of gene expression: a review. *Cancer Res*. May 15, 1988;48(10):2659-68.
- Stull et al., Antigene, ribozyme and aptamer nucleic acid drugs: progress and prospects. *Pharm Res*. Apr. 1995;12(4):465-83.
- Subramanian et al., Theoretical considerations on the “spine of hydration” in the minor groove of d(CGCGAATTCGCG). d(GCGCTTAAGCGC): Monte Carlo computer simulation. *Proc Natl Acad Sci U S A*. Mar. 1988;85(6):1836-40.
- Tanaka et al., An antisense oligonucleotide complementary to a sequence in I gamma 2b increases gamma 2b germline transcripts, stimulates B cell DnA synthesis, and inhibits immunoglobulin secretion. *J Exp Med*. Feb. 1, 1992;175(2):597-607.
- Thorne et al., Experimental grain dust atmospheres generated by wet and dry aerosolization techniques. *Am J Ind Med*. Jan. 1994;25(1):109-12 Abstract.
- Tomai et al., Immunomodulating and antiviral activities of the imidazoquinoline S-28463. *Antiviral Res*. Nov. 1995;28(3):253-64.
- Turner et al., Characterization of a primate model of asthma using anti-allergy/anti-asthma agents. *Inflamm Res*. May 1996;45(5):239-45.
- Turner et al., Effects of rolipram on responses to acute and chronic antigen exposure in monkeys. *Am J Respir Crit Care Med*. May 1994;149(5):1153-9.
- Turner et al., In vitro and in vivo effects of leukotriene B4 antagonism in a primate model of asthma. *J Clin Invest*. Jan. 15, 1996;97(2):381-7.
- Turner et al., Leukotriene D4 receptor antagonism reduces airway hyperresponsiveness in monkeys. *Pulm Pharmacol*. Feb. 1994;7(1):49-58.
- Uhlmann et al., Antisense oligonucleotides: a new therapeutic principle. *Chem Rev*. Jun. 1990;90(4):544-84.
- Vandenbulcke et al., The innate immune system and its role in allergic disorders. *Int Arch Allergy Immunol*. 2006;139(2):159-65. Abstract.
- Verthelyi et al., Immunoregulatory activity of CpG oligonucleotides in humans and nonhuman primates. *Clin Immunol*. Oct. 2003;109(1):64-71.
- Vlassov et al., Oligonucleotides in cells and in organisms: pharmacological considerations. Prospects for Antisense Nucleic Acid Therapy of Cancer and Aids. 1991:243-66.
- Wagner et al., Gene inhibition using antisense oligodeoxynucleotides. *Nature*. Nov. 24, 1994;372(6504):333-5 Abstract.
- Walker et al., Activated T cells and cytokines in bronchoalveolar lavages from patients with various lung diseases associated with eosinophilia. *Am J Respir Crit Care Med*. Oct. 1994;150(4):1038-48 Abstract.
- Wallace et al., Oligonucleotide probes for the screening of recombinant DNA libraries. *Methods Enzymol*. 1987;152:432-42.
- Wallner et al., Immunotherapy with T-cell-reactive peptides derived from allergens. *Allergy*. May 1994;49(5):302-8.
- Weeratna et al., CPG ODN allows lower dose of antigen against hepatitis B surface antigen in BALB/c mice. *Immunol Cell Biol*. Feb. 2003;81(1):59-62.
- Weeratna et al., Optimization strategies for DnA vaccines. *Intervirology*. 2000;43(4-6):218-26.
- Weiss et al., DNA vaccines for allergy treatment. *Methods Mol Med*. 2006;127:253-67. Review.
- Weiss et al., Upping the antisense ante—scientists bet on profits from reverse genetics. *Science News*. Feb. 16, 1991;139:108-9.
- Whalen et al., DNA vaccines for emerging infectious diseases: what if? *Emerg Infect Dis*. Jul.-Sep. 1996;2(3):168-75.
- Whalen et al., DNA-mediated immunization to the hepatitis B surface antigen. Activation and entrainment of the immune response. *Ann NY Acad Sci*. Nov. 27, 1995;772:64-76.
- Witt et al., Phase I trial of an oral immunomodulator and interferon inducer in cancer patients. *Cancer Res*. Nov. 1, 1993;53(21):5176-80.
- Wong et al., Formoterol compared with beclomethasone and placebo on allergen-induced asthmatic responses. *Am Rev Respir Dis*. Nov. 1992;146(5 Pt 1):1156-60.
- Wu et al., Receptor-mediated gene delivery and expression in vivo. *J Biol Chem*. Oct. 15, 1988;263(29):14621-4.
- Wu-Pong et al., Oligonucleotides: opportunities for drug therapy and research. *Pharm Technol*. Oct. 1994;18:102-14.
- Wyatt et al., Combinatorially selected guanosine-quartet structure is a potent inhibitor of human immunodeficiency virus envelope-mediated cell fusion. *Proc Natl Acad Sci U S A*. Feb. 15, 1994;91(4):1356-60.

Yamamoto et al., DNA from bacteria, but not from vertebrates, induces interferons, activates natural killer cells and inhibits tumor growth. *Microbiol Immunol.* 1992;36(9):983-97.

Yamamoto et al., In vitro augmentation of natural killer cell activity and production of interferon-alpha/beta and—gamma with deoxyribonucleic acid fraction from *Mycobacterium bovis* BCG. *Jpn J Cancer Res.* Jul. 1988;79(7):866-73.

Yamamoto et al., Mode of action of oligonucleotide fraction extracted from *Mycobacterium bovis* BCG. *Kekkaku.* 1994; 69(9):29-32.

Zhang et al., Antigen- and isotype-specific immune responses to a recombinant antigen-allergen chimeric (RAAC) protein. *J Immunol.* Jul. 15, 1993;151(2):791-9.

Zhao et al., Comparison of cellular binding and uptake of antisense phosphodiester, phosphorothioate, and mixed phosphorothioate and

methylphosphonate oligonucleotides. *Antisense Res Dev.* 1993 Spring;3(1):53-66.

Zhao et al., Stage-specific oligonucleotide uptake in murine bone marrow B-cell precursors. *Blood.* Dec. 1, 1994;84(11):3660-6.

Patent Interference No. 105,171. Miscellaneous Motion 1 (Unopposed Additional Request for File Histories). Apr. 15, 2004.

Patent Interference No. 105,171. Miscellaneous Motion 2 (Unopposed Request for Addition to Iowa's file History). Apr. 15, 2004.

Patent Interference No. 105,171. Regents of the University of California Notice of Filing Appeal. Mar. 17, 2005.

Patent Interference No. 105,526. Krieg Substantive Motion 1 (for unpatentability based on interference estoppel). (Electronically filed, unsigned). Jun. 18, 2007.

* cited by examiner




FIG. 1A




FIG. 1B




FIG. 1C




FIG. 2




FIG. 3




FIG. 4A




FIG. 4B




FIG. 5




FIG. 6

Treatment:	0	EC	CT	LPS
		DNA	DNA	
min:	0	15 30	15 30	15 30




FIG. 7

FIG. 8A





FIG. 8B






FIG. 9




FIG. 10




FIG. II




FIG. 12




FIG.13




FIG. 14




FIG. 15

**METHODS OF USING
IMMUNOSTIMULATORY NUCLEIC ACID
MOLECULES TO TREAT ALLERGIC
CONDITIONS**

RELATED APPLICATIONS

This application is a continuation of U.S. patent application Ser. No. 10/894,657, filed Jul. 16, 2004, now abandoned, which is a continuation of U.S. patent application Ser. No. 09/818,918, filed Mar. 27, 2001, now abandoned, which is a divisional of U.S. patent application Ser. No. 08/738,652, filed Oct. 30, 1996, now issued as U.S. Pat. No. 6,207,646 B1, which is a continuation-in-part of U.S. patent application Ser. No. 08/386,063, filed Feb. 7, 1995, now issued as U.S. Pat. No. 6,194,388 B1, which is a continuation-in-part of U.S. patent application Ser. No. 08/276,358, filed Jul. 15, 1994, now abandoned, each of which is incorporated herein by reference.

GOVERNMENT SUPPORT

The work resulting in this invention was supported in part by National Institute of Health Grant No. R29-AR42556-01. The U.S. Government has certain rights in the invention.

BACKGROUND OF THE INVENTION

DNA Binds to Cell Membranes and is Internalized

In the 1970's, several investigators reported the binding of high molecular weight DNA to cell membranes (Lerner, R. A., W. Meinke, and D. A. Goldstein. 1971. "Membrane-associated DNA in the cytoplasm of diploid human lymphocytes". *Proc. Natl. Acad. Sci. USA* 68:1212; Agrawal, S. K., R. W. Wagner, P. K. McAllister, and B. Rosenberg. 1975. "Cell-surface-associated nucleic acid in tumorigenic cells made visible with platinum-pyrimidine complexes by electron microscopy". *Proc. Natl. Acad. Sci. USA* 72:928). In 1985, Bennett et al. presented the first evidence that DNA binding to lymphocytes is similar to a ligand receptor interaction: binding is saturable, competitive, and leads to DNA endocytosis and degradation into oligonucleotides (Bennett, R. M., G. T. Gabor, and M. M. Merritt. 1985. "DNA binding to human leukocytes. Evidence for a receptor-mediated association, internalization, and degradation of DNA". *J. Clin. Invest.* 76:2182). Like DNA, oligodeoxyribonucleotides (ODNs) are able to enter cells in a saturable, sequence independent, and temperature and energy dependent fashion (reviewed in Jaroszewski, J. W., and J. S. Cohen. 1991. "Cellular uptake of antisense oligodeoxynucleotides". *Advanced Drug Delivery Reviews* 6:235; Akhtar, S., Y. Shoji, and R. L. Juliano. 1992. "Pharmaceutical aspects of the biological stability and membrane transport characteristics of antisense oligonucleotides". In: *Gene Regulation: Biology of Antisense RNA and DNA*. R. P. Erickson, and J. G. Izant, eds. Raven Press, Ltd. New York, pp. 133; and Zhao, Q., T. Waldschmidt, E. Fisher, C. J. Herrera, and A. M. Krieg., 1994. "Stage specific oligonucleotide uptake in murine bone marrow B cell precursors". *Blood*, 84:3660). No receptor for DNA or ODN uptake has yet been cloned, and it is not yet clear whether ODN binding and cell uptake occurs through the same or a different mechanism from that of high molecular weight DNA.

Lymphocyte ODN uptake has been shown to be regulated by cell activation. Spleen cells stimulated with the B cell mitogen LPS had dramatically enhanced ODN uptake in the

B cell population, while spleen cells treated with the T cell mitogen Con A showed enhanced ODN uptake by T but not B cells (Krieg, A. M., F. Gmelig-Meyling, M. F. Gourley, W. J. Kisch, L. A. Chrisey, and A. D. Steinberg. 1991. "Uptake of oligodeoxyribonucleotides by lymphoid cells is heterogeneous and inducible". *Antisense Research and Development* 1:161).

Immune Effects of Nucleic Acids

Several polynucleotides have been extensively evaluated as biological response modifiers. Perhaps the best example is poly (I,C) which is a potent inducer of IFN production as well as a macrophage activator and inducer of NK activity (Talmadge, J. E., J. Adams, H. Phillips, M. Collins, B. Lenz, M. Schneider, E. Schlick, R. Ruffmann, R. H. Wilttrout, and M. A. Chirigos. 1985. "Immunomodulatory effects in mice of polyinosinic-polycytidylic acid complexed with poly-L-lysine and carboxymethylcellulose". *Cancer Res.* 45:1058; Wilttrout, R. H., R. R. Salup, T. A. Twilley, and J. E. Talmadge. 1985. "Immunomodulation of natural killer activity by polyribonucleotides". *J. Biol. Resp. Mod.* 4:5.12; Krown, S. E. 1986. "Interferons and interferon inducers in cancer treatment". *Sem. Oncol.* 13:207; and Ewel, C. H., S. J. Urba, W. C. Kopp, J. W. Smith II, R. G. Steis, J. L. Rossio, D. L. Longo, M. J. Jones, W. G. Alvord, C. M. Pinsky, J. M. Beveridge, K. L. McNitt, and S. P. Creekmore. 1992. "Polyinosinic-polycytidylic acid complexed with poly-L-lysine and carboxymethylcellulose in combination with interleukin-2 in patients with cancer: clinical and immunological effects". *Canc. Res.* 52:3005). It appears that this murine NK activation may be due solely to induction of IFN- β secretion (Ichikawa, R., and C. A. Biron. 1993. "IFN induction and associated changes in splenic leukocyte distribution". *J. Immunol.* 150:3713). This activation was specific for the ribose sugar since deoxyribose was ineffective. Its potent in vitro antitumor activity led to several clinical trials using poly (I,C) complexed with poly-L-lysine and carboxymethylcellulose (to reduce degradation by RNase) (Talmadge, J. E., et al., 1985. cited supra; Wilttrout, R. H., et al., 1985. cited supra; Krown, S. E., 1986. cited supra; and Ewel, C. H., et al., 1992. cited supra). Unfortunately, toxic side effects have thus far prevented poly (I,C) from becoming a useful therapeutic agent.

Guanine ribonucleotides substituted at the C8 position with either a bromine or a thiol group are B cell mitogens and may replace "B cell differentiation factors" (Feldbush, T. L., and Z. K. Ballas. 1985. "Lymphokine-like activity of 8-mercaptoguanosine: induction of T and B cell differentiation". *J. Immunol.* 134:3204; and Goodman, M. G. 1986. "Mechanism of synergy between T cell signals and C8-substituted guanine nucleosides in humoral immunity: B lymphotropic cytokines induce responsiveness to 8-mercaptoguanosine". *J. Immunol.* 136:3335). 8-mercaptoguanosine and 8-bromoguanosine also can substitute for the cytokine requirement for the generation of MHC restricted CTL (Feldbush, T. L., 1985. cited supra), augment murine NK activity (Koo, G. C., M. E. Jewell, C. L. Manyak, N. H. Sigal, and L. S. Wicker. 1988. "Activation of murine natural killer cells and macrophages by 8-bromoguanosine". *J. Immunol.* 140:3249), and synergize with IL-2 in inducing murine LAK generation (Thompson, R. A., and Z. K. Ballas. 1990. "Lymphokine-activated killer (LAK) cells. V. 8-Mercaptoguanosine as an IL-2-sparing agent in LAK generation". *J. Immunol.* 145:3524). The NK and LAK augmenting activities of these C8-substituted guanines appear to be due to their induction of IFN (Thompson, R. A., et al. 1990. cited supra). Recently, a 5' triphosphorylated thymidine produced by a mycobacterium was found to be mitogenic for a subset of human $\gamma\delta$ T cells (Constant, P., F. Davodeau, M.-A. Peyrat, Y. Poquet, G. Puzo, M. Bonne-

ille, and J.-J. Fournie. 1994. "Stimulation of human $\gamma\delta$ T cells by nonpeptidic mycobacterial ligands" *Science* 264:267). This report indicated the possibility that the immune system may have evolved ways to preferentially respond to microbial nucleic acids.

Several observations suggest that certain DNA structures may also have the potential to activate lymphocytes. For example, Bell et al. reported that nucleosomal protein-DNA complexes (but not naked DNA) in spleen cell supernatants caused B cell proliferation and immunoglobulin secretion (Bell, D. A., B. Morrison, and P. VandenBygaart. 1990. "Immunogenic DNA-related factors". *J. Clin. Invest.* 85:1487). In other cases, naked DNA has been reported to have immune effects. For example, Messina et al. have recently reported that 260 to 800 bp fragments of poly (dG) \cdot (dC) and poly (dG \cdot dC) were mitogenic for B cells (Messina, J. P., G. S. Gilkeson, and D. S. Pisetsky. 1993. "The influence of DNA structure on the in vitro stimulation of murine lymphocytes by natural and synthetic polynucleotide antigens". *Cell. Immunol.* 147:148). Tokunaga, et al. have reported that dG \cdot dC induces IFN- γ and NK activity (Tokunaga, S. Yamamoto, and K. Namba. 1988. "A synthetic single-stranded DNA, poly(dG,dC), induces interferon- α/β and - γ , augments natural killer activity, and suppresses tumor growth" *Jpn. J. Cancer Res.* 79:682). Aside from such artificial homopolymer sequences, Pisetsky et al. reported that pure mammalian DNA has no detectable immune effects, but that DNA from certain bacteria induces B cell activation and immunoglobulin secretion (Messina, J. P., G. S. Gilkeson, and D. S. Pisetsky. 1991. "Stimulation of in vitro murine lymphocyte proliferation by bacterial DNA". *J. Immunol.* 147:1759). Assuming that these data did not result from some unusual contaminant, these studies suggested that a particular structure or other characteristic of bacterial DNA renders it capable of triggering B cell activation. Investigations of mycobacterial DNA sequences have demonstrated that ODN which contain certain palindrome sequences can activate NK cells (Yamamoto, S., T. Yamamoto, T. Kataoka, E. Kuramoto, O. Yano, and T. Tokunaga. 1992. "Unique palindromic sequences in synthetic oligonucleotides are required to induce INF and augment INF-mediated natural killer activity". *J. Immunol.* 148:4072; Kuramoto, E., O. Yano, Y. Kimura, M. Baba, T. Makino, S. Yamamoto, T. Yamamoto, T. Kataoka, and T. Tokunaga. 1992. "Oligonucleotide sequences required for natural killer cell activation". *Jpn. J. Cancer Res.* 83:1128).

Several phosphorothioate modified ODN have been reported to induce in vitro or in vivo B cell stimulation (Tanaka, T., C. C. Chu, and W. E. Paul. 1992. "An antisense oligonucleotide complementary to a sequence in Iy2b increases γ 2b germline transcripts, stimulates B cell DNA synthesis, and inhibits immunoglobulin secretion". *J. Exp. Med.* 175:597; Branda, R. F., A. L. Moore, L. Mathews, J. J. McCormack, and G. Zon. 1993. "Immune stimulation by an antisense oligomer complementary to the rev gene of HIV-1". *Biochem. Pharmacol.* 45:2037; McIntyre, K. W., K. Lombard-Gillooly, J. R. Perez, C. Kunsch, U. M. Sarmiento, J. D. Larigan, K. T. Landreth, and R. Narayanan. 1993. "A sense phosphorothioate oligonucleotide directed to the initiation codon of transcription factor NF κ B T65 causes sequence-specific immune stimulation". *Antisense Res. Develop.* 3:309; and Pisetsky, D. S., and C. F. Reich. 1993. "Stimulation of murine lymphocyte proliferation by a phosphorothioate oligonucleotide with antisense activity for herpes simplex virus". *Life Sciences* 54:101). These reports do not suggest a common structural motif or sequence element in these ODN that might explain their effects.

The CREB/ATF Family of Transcription Factors and their Role in Replication

The cAMP response element binding protein (CREB) and activating transcription factor (ATF) or CREB/ATF family of transcription factors is a ubiquitously expressed class of transcription factors of which 11 members have so far been cloned (reviewed in de Groot, R. P., and P. Sassone-Corsi: "Hormonal control of gene expression: Multiplicity and versatility of cyclic adenosine 3',5'-monophosphate-responsive nuclear regulators". *Mol. Endocrin.* 7:145, 1993; Lee, K. A. W., and N. Masson: "Transcriptional regulation by CREB and its relatives". *Biochim. Biophys. Acta* 1174:221, 1993.). They all belong to the basic region/leucine zipper (bZip) class of proteins. All cells appear to express one or more CREB/ATF proteins, but the members expressed and the regulation of mRNA splicing appear to be tissue-specific. Differential splicing of activation domains can determine whether a particular CREB/ATF protein will be a transcriptional inhibitor or activator. Many CREB/ATF proteins activate viral transcription, but some splicing variants which lack the activation domain are inhibitory. CREB/ATF proteins can bind DNA as homo- or hetero-dimers through the cAMP response element, the CRE, the consensus form of which is the unmethylated sequence TGACGTC (binding is abolished if the CpG is methylated) (Iguchi-Ariga, S. M. M., and W. Schaffner: "CpG methylation of the cAMP-responsive enhancer/promoter sequence TGACGTCA abolishes specific factor binding as well as transcriptional activation". *Genes & Develop.* 3:612, 1989.).

The transcriptional activity of the CRE is increased during B cell activation (Xie, H. T. C. Chiles, and T. L. Rothstein: "Induction of CREB activity via the surface Ig receptor of B cells". *J. Immunol.* 151:880, 1993.). CREB/ATF proteins appear to regulate the expression of multiple genes through the CRE including immunologically important genes such as fos, jun B, Rb-1, IL-6, IL-1 (Tsukada, J., K. Saito, W. R. Waterman, A. C. Webb, and P. E. Auron: "Transcription factors NF-IL6 and CREB recognize a common essential site in the human prointerleukin 1 β gene". *Mol. Cell. Biol.* 14:7285, 1994; Gray, G. D., O. M. Hernandez, D. Hebel, M. Root, J. M. Pow-Sang, and E. Wickstrom: "Antisense DNA inhibition of tumor growth induced by c-Ha-ras oncogene in nude mice". *Cancer Res.* 53:577, 1993), IFN- β (Du, W., and T. Maniatis: "An ATF/CREB binding site protein is required for virus induction of the human interferon B gene". *Proc. Natl. Acad. Sci. USA* 89:2150, 1992), TGF- β 1 (Asiedu, C. K., L. Scott, R. K. Assoian, M. Ehrlich: "Binding of AP-1/CREB proteins and of MDPB to contiguous sites downstream of the human TGF-B1 gene". *Biochim. Biophys. Acta* 1219:55, 1994.), TGF- β 2, class II MHC (Cox, P. M., and C. R. Goding: "An ATF/CREB binding motif is required for aberrant constitutive expression of the MHC class II DR α promoter and activation by SV40 T-antigen". *Nucl. Acids Res.* 20:4881, 1992.), E-selectin, GM-CSF, CD-8 α , the germline Iga constant region gene, the TCR V β gene, and the proliferating cell nuclear antigen (Huang, D., P. M. Shipman-Appasamy, D. J. Orten, S. H. Hinrichs, and M. B. Prystowsky: "Promoter activity of the proliferating-cell nuclear antigen gene is associated with inducible CRE-binding proteins in interleukin 2-stimulated T lymphocytes". *Mol. Cell. Biol.* 14:4233, 1994.). In addition to activation through the cAMP pathway, CREB can also mediate transcriptional responses to changes in intracellular Ca⁺⁺ concentration (Sheng, M., G. McFadden, and M. E. Greenberg: "Membrane depolarization and calcium induce c-fos transcription via phosphorylation of transcription factor CREB". *Neuron* 4:571, 1990).

The role of protein-protein interactions in transcriptional activation by CREB/ATF proteins appears to be extremely important. There are several published studies reporting direct or indirect interactions between NFκB proteins and CREB/ATF proteins (Whitley, et. al., (1994) *Mol. & Cell. Biol.* 14:6464; Cogswell, et al., (1994) *J. Immun.* 153:712; Hines, et al., (1993) *Oncogene* 8:3189; and Du, et al., (1993) *Cell* 74:887. Activation of CREB through the cyclic AMP pathway requires protein kinase A (PKA), which phosphorylates CREB³⁴¹ on ser¹³³ and allows it to bind to a recently cloned protein, CBP (Kwok, R. P. S., J. R. Lundblad, J. C. Chrvia, J. P. Richards, H. P. Bachinger, R. G. Brennan, S. G. E. Roberts, M. R. Green, and R. H. Goodman: "Nuclear protein CBP is a coactivator for the transcription factor CREB". *Nature* 370:223, 1994; Arias, J., A. S. Alberts, P. Brindle, F. X. Claret, T. Smea, M. Karin, J. Feramisco, and M. Montminy: "Activation of cAMP and mitogen responsive genes relies on a common nuclear factor". *Nature* 370:226, 1994.). CBP in turn interacts with the basal transcription factor TFIIB causing increased transcription. CREB also has been reported to interact with dTAFII 110, a TATA binding protein-associated factor whose binding may regulate transcription (Ferreri, K., G. Gill, and M. Montminy: "The cAMP-regulated transcription factor CREB interacts with a component of the TFIID complex". *Proc. Natl. Acad. Sci. USA* 91:1210, 1994.). In addition to these interactions, CREB/ATF proteins can specifically bind multiple other nuclear factors (Hoeffler, J. P., J. W. Lustbader, and C.-Y. Chen: "Identification of multiple nuclear factors that interact with cyclic adenosine 3',5'-monophosphate response element-binding protein and activating transcription factor-2 by protein-protein interactions". *Mol. Endocrinol.* 5:256, 1991) but the biologic significance of most of these interactions is unknown. CREB is normally thought to bind DNA either as a homodimer or as a heterodimer with several other proteins. Surprisingly, CREB monomers constitutively activate transcription (Krajewski, W., and K. A. W. Lee: "A monomeric derivative of the cellular transcription factor CREB functions as a constitutive activator". *Mol. Cell. Biol.* 14:7204, 1994.).


Aside from their critical role in regulating cellular transcription, it has recently been shown that CREB/ATF proteins are subverted by some infectious viruses and retroviruses, which require them for viral replication. For example, the cytomegalovirus immediate early promoter, one of the strongest known mammalian promoters, contains eleven copies of the CRE which are essential for promoter function (Chang, Y.-N., S. Crawford, J. Stall, D. R. Rawlins, K.-T. Jeang, and G. S. Hayward: "The palindromic series I repeats in the simian cytomegalovirus major immediate-early promoter behave as both strong basal enhancers and cyclic AMP response elements". *J. Virol.* 64:264, 1990). At least some of the transcriptional activating effects of the adenovirus E1A protein, which induces many promoters, are due to its binding to the DNA binding domain of the CREB/ATF protein, ATF-2, which mediates E1A inducible transcription activation (Liu, F., and M. R. Green: "Promoter targeting by adenovirus E1a through interaction with different cellular DNA-binding domains". *Nature* 368:520, 1994). It has also been suggested that E1A binds to the CREB-binding protein, CBP (Arany, Z., W. R. Sellers, D. M. Livingston, and R. Eckner: "E1A-associated p300 and CREB-associated CBP belong to a conserved family of coactivators". *Cell* 77:799, 1994). Human T lymphotropic virus-I (HTLV-1), the retrovirus which causes human T cell leukemia and tropical spastic paresis, also requires CREB/ATF proteins for replication. In this case, the retrovirus produces a protein, Tax, which binds to CREB/ATF proteins and redirects them from their normal cellular binding

sites to different DNA sequences (flanked by G- and C-rich sequences) present within the HTLV transcriptional enhancer (Paca-Uccaralertkun, S., L.-J. Zhao, N. Adya, J. V. Cross, B. R. Cullen, I. M. Boros, and C. Z. Giam: "In vitro selection of DNA elements highly responsive to the human T-cell lymphotropic virus type I transcriptional activator, Tax". *Mol. Cell. Biol.* 14:456, 1994; Adya, N., L.-J. Zhao, W. Huang, I. Boros, and C.-Z. Giam: "Expansion of CREB's DNA recognition specificity by Tax results from interaction with Ala-Ala-Arg at positions 282-284 near the conserved DNA-binding domain of CREB". *Proc. Natl. Acad. Sci. USA* 91:5642, 1994).

SUMMARY OF THE INVENTION


The instant invention is based on the finding that certain nucleic acids containing unmethylated cytosine-guanine (CpG) dinucleotides activate lymphocytes in a subject and redirect a subject's immune response from a Th2 to a Th1 (e.g. by inducing monocytic cells and other cells to produce Th1 cytokines, including IL-12, IFN-γ and GM-CSF). Based on this finding, the invention features, in one aspect, novel immunostimulatory nucleic acid compositions.

In a preferred embodiment, the immunostimulatory nucleic acid contains a consensus mitogenic CpG motif represented by the formula:



wherein X₁ is selected from the group consisting of A, G and T; and X₂ is C or T.

In a particularly preferred embodiment an immunostimulatory nucleic acid molecule contains a consensus mitogenic CpG motif represented by the formula:



wherein C and G are unmethylated; and X₁, X₂, X₃ and X₄ are nucleotides.

Enhanced immunostimulatory activity of human cells occurs where X₁X₂ is selected from the group consisting of GpT, GpG, GpA and ApA and/or X₃X₄ is selected from the group consisting of TpT, CpT and GpT (Table 5). For facilitating uptake into cells, CpG containing immunostimulatory nucleic acid molecules are preferably in the range of 8 to 40 base pairs in size. However, nucleic acids of any size (even many kb long) are immunostimulatory if sufficient immunostimulatory motifs are present, since such larger nucleic acids are degraded into oligonucleotides inside of cells. Preferred synthetic oligonucleotides do not include a GCG trinucleotide sequence at or near the 5' and/or 3' terminals and/or the consensus mitogenic CpG motif is not a palindrome. Prolonged immunostimulation can be obtained using stabilized oligonucleotides, particularly phosphorothioate stabilized oligonucleotides.

In a second aspect, the invention features useful therapies, which are based on the immunostimulatory activity of the nucleic acid molecules. For example, the immunostimulatory nucleic acid molecules can be used to treat, prevent or ameliorate an immune system deficiency (e.g., a tumor or cancer or a viral, fungal, bacterial or parasitic infection in a subject). In addition, immunostimulatory nucleic acid molecules can be administered to stimulate a subject's response to a vaccine.

Further, by redirecting a subject's immune response from Th2 to Th1, the instant claimed nucleic acid molecules can be administered to treat or prevent the symptoms of asthma. In

addition, the instant claimed nucleic acid molecules can be administered in conjunction with a particular allergen to a subject as a type of desensitization therapy to treat or prevent the occurrence of an allergic reaction.

Further, the ability of immunostimulatory nucleic acid molecules to induce leukemic cells to enter the cell cycle supports the use of immunostimulatory nucleic acid molecules in treating leukemia by increasing the sensitivity of chronic leukemia cells and then administering conventional ablative chemotherapy, or combining the immunostimulatory nucleic acid molecules with another immunotherapy.

Other features and advantages of the invention will become more apparent from the following detailed description and claims.

BRIEF DESCRIPTION OF THE FIGURES

FIG. 1A-C are graphs plotting dose-dependent IL-6 production in response to various DNA sequences in T cell depleted spleen cell cultures. A. *E. coli* DNA (●) and calf thymus DNA (■) sequences and LPS (at 10× the concentration of *E. coli* and calf thymus DNA) (◆). B. Control phosphodiester oligodeoxynucleotide (ODN) 5' ATGGAAGTCCAGTGTTC 3' (SEQ ID NO:1) (■) and two phosphodiester CpG ODN 5' ATCGACCTACGTGCGT-TCTC 3' (SEQ ID NO:2) (◆) and 5' TCCATAACGTTCTGATGCT 3' (SEQ ID NO:3) (●). C. Control phosphorothioate ODN 5' GCTAGATGTTAGCGT 3' (SEQ ID NO:4) (■) and two phosphorothioate CpG ODN 5' GAGAACGTC-GACCTTCGAT 3' (SEQ ID NO:5) (◆) and 5' GCATGACGTTGAGCT 3' (SEQ ID NO:6) (●). Data present the mean±standard deviation of triplicates.

FIG. 2 is a graph plotting IL-6 production induced by CpG DNA in vivo as determined 1-8 hrs after injection. Data represent the mean from duplicate analyses of sera from two mice. BALB/c mice (two mice/group) were injected iv. with 100 μl of PBS (□) or 200 μg of CpG phosphorothioate ODN 5' TCCATGACGTTCTGATGCT 3' (SEQ ID NO: 7) (■) or non-CpG phosphorothioate ODN 5' TCCATGAGCTTCCTGAGTCT 3' (SEQ ID NO: 8) (◆).

FIG. 3 is an autoradiograph showing IL-6 mRNA expression as determined by reverse transcription polymerase chain reaction in liver, spleen, and thymus at various time periods after in vivo stimulation of BALB/c mice (two mice/group) injected iv with 100 μl A of PBS, 200 μg of CpG phosphorothioate ODN 5' TCCATGACGTTCTGATGCT 3' (SEQ ID NO: 7) or non-CpG phosphorothioate ODN 5' TCCATGAGCTTCCTGAGTCT 3' (SEQ ID NO: 8).

FIG. 4A is a graph plotting dose-dependent inhibition of CpG-induced IgM production by anti-IL-6. Splenic B-cells from DBA/2 mice were stimulated with CpG ODN 5' TCCAAAGACGTTCTGATGCT 3' (SEQ ID NO: 9) in the presence of the indicated concentrations of neutralizing anti-IL-6 (◆) or isotype control Ab (●) and IgM levels in culture supernatants determined by ELISA. In the absence of CpG ODN, the anti-IL-6 Ab had no effect on IgM secretion (■).

FIG. 4B is a graph plotting the stimulation index of CpG-induced splenic B cells cultured with anti-IL-6 and CpG S-ODN 5' TCCATGACGTTCTGATGCT 3' (SEQ ID NO: 7) (◆) or anti-IL-6 antibody only (■). Data present the mean±standard deviation of triplicates.

FIG. 5 is a bar graph plotting chloramphenicol acetyltransferase (CAT) activity in WEHI-231 cells transfected with a promoter-less CAT construct (pCAT), positive control plasmid (RSV), or IL-6 promoter-CAT construct alone or cultured with CpG 5' TCCATGACGTTCTGATGCT 3' (SEQ ID NO: 7) or non-CpG 5' TCCATGAGCTTCCTGAGTCT 3'

(SEQ ID NO: 8) phosphorothioate ODN at the indicated concentrations. Data present the mean of triplicates.

FIG. 6 is a schematic overview of the immune effects of the immunostimulatory unmethylated CpG containing nucleic acids, which can directly activate both B cells and monocytic cells (including macrophages and dendritic cells) as shown. The immunostimulatory oligonucleotides do not directly activate purified NK cells, but render them competent to respond to IL-12 with a marked increase in their IFN-γ production. By inducing IL-12 production and the subsequent increased IFN-γ secretion by NK cells, the immunostimulatory nucleic acids promote a Th1 type immune response. No direct activation of proliferation of cytokine secretion by highly purified T cells has been found. However, the induction of Th1 cytokine secretion by the immunostimulatory oligonucleotides promotes the development of a cytotoxic lymphocyte response.

FIG. 7 is an autoradiograph showing NFκB mRNA induction in monocytes treated with *E. coli* (EC) DNA (containing unmethylated CpG motifs), control (CT) DNA (containing no unmethylated CpG motifs) and lipopolysaccharide (LPS) at various measured times, 15 and 30 minutes after contact.

FIG. 8A shows the results from a flow cytometry study using mouse B cells with the dihydrorhodamine 123 dye to determine levels of reactive oxygen species. The dye only sample in Panel A of the figure shows the background level of cells positive for the dye at 28.6%. This level of reactive oxygen species was greatly increased to 80% in the cells treated for 20 minutes with PMA and ionomycin, a positive control (Panel B). The cells treated with the CpG oligo (TC-CATGACGTTCTGACGTT SEQ ID NO: 10) also showed an increase in the level of reactive oxygen species such that more than 50% of the cells became positive (Panel D). However, cells treated with an oligonucleotide with the identical sequence except that the CpGs were switched (TCCATGAGCTTCCTGAGTCT SEQ ID NO: 11) did not show this significant increase in the level of reactive oxygen species (Panel E).

FIG. 8B shows the results from a flow cytometry study using mouse B cells in the presence of chloroquine with the dihydrorhodamine 123 dye to determine levels of reactive oxygen species. Chloroquine slightly lowers the background level of reactive oxygen species in the cells such that the untreated cells in Panel A have only 4.3% that are positive. Chloroquine completely abolishes the induction of reactive oxygen species in the cells treated with CpG DNA (Panel B) but does not reduce the level of reactive oxygen species in the cells treated with PMA and ionomycin (Panel E).

FIG. 9 is a graph plotting lung lavage cell count over time. The graph shows that when the mice are initially injected with *Schistosoma mansoni* eggs "egg", which induces a Th2 immune response, and subsequently inhale *Schistosoma mansoni* egg antigen "SEA" (open circle), many inflammatory cells are present in the lungs. However, when the mice are initially given CpG oligo (SEQ ID NO: 10) along with egg, the inflammatory cells in the lung are not increased by subsequent inhalation of SEA (open triangles).

FIG. 10 is a graph plotting lung lavage eosinophil count over time. Again, the graph shows that when the mice are initially injected with egg and subsequently inhale SEA (open circle), many eosinophils are present in the lungs. However, when the mice are initially given CpG oligo (SEQ ID NO: 10) along with egg, the inflammatory cells in the lung are not increased by subsequent inhalation of the SEA (open triangles).

FIG. 11 is a bar graph plotting the effect on the percentage of macrophage, lymphocyte, neutrophil and eosinophil cells

induced by exposure to saline alone; egg, then SEA; egg and SEQ ID NO: 11, then SEA; and egg and control oligo (SEQ ID NO: 11), then SEA. When the mice are treated with the control oligo at the time of the initial exposure to the egg, there is little effect on the subsequent influx of eosinophils into the lungs after inhalation of SEA. Thus, when mice inhale the eggs on days 14 or 21, they develop an acute inflammatory response in the lungs. However, giving a CpG oligo along with the eggs at the time of initial antigen exposure on days 0 and 7 almost completely abolishes the increase in eosinophils when the mice inhale the egg antigen on day 14.

FIG. 12 is a bar graph plotting eosinophil count in response to injection of various amounts of the protective oligo SEQ ID NO: 10.

FIG. 13 is a graph plotting interleukin 4 (IL-4) production (pg/ml) in mice over time in response to injection of egg, then SEA (open diamond); egg and SEQ ID NO: 10, then SEA (open circle); or saline, then saline (open square). The graph shows that the resultant inflammatory response correlates with the levels of the Th2 cytokine IL-4 in the lung.

FIG. 14 is a bar graph plotting interleukin 12 (IL-12) production (pg/ml) in mice over time in response to injection of saline; egg, then SEA; or SEQ ID NO: 10 and egg, then SEA. The graph shows that administration of an oligonucleotide containing an unmethylated CpG motif can actually redirect the cytokine response of the lung to production of IL-12, indicating a Th1 type of immune response.

FIG. 15 is a bar graph plotting interferon gamma (IFN- γ) production (pg/ml) in mice over time in response to injection of saline; egg, then saline; or SEQ ID NO: 10 and egg, then SEA. The graph shows that administration of an oligonucleotide containing an unmethylated CpG motif can also redirect the cytokine response of the lung to production of IFN- γ , indicating a Th1 type of immune response.

DETAILED DESCRIPTION OF THE INVENTION

Definitions

As used herein, the following terms and phrases shall have the meanings set forth below:

An “allergen” refers to a substance that can induce an allergic or asthmatic response in a susceptible subject. The list of allergens is enormous and can include pollens, insect venoms, animal dander, dust, fungal spores and drugs (e.g. penicillin). Examples of natural, animal and plant allergens include proteins specific to the following genera: Canine (*Canis familiaris*); *Dermatophagoides* (e.g. *Dermatophagoides farinae*); *Felis* (*Felis domesticus*); *Ambrosia* (*Ambrosia artemisiifolia*); *Lolium* (e.g. *Lolium perenne* or *Lolium multiflorum*); *Cryptomeria* (*Cryptomeria japonica*); *Alternaria* (*Alternaria alternata*); *Alder*; *Alnus* (*Alnus glutinosa*); *Betula* (*Betula verrucosa*); *Quercus* (*Quercus alba*); *Olea* (*Olea europa*); *Artemisia* (*Artemisia vulgaris*); *Plantago* (e.g. *Plantago lanceolata*); *Parietaria* (e.g. *Parietaria officinalis* or *Parietaria judaica*); *Blattella* (e.g. *Blattella germanica*); *Apis* (e.g. *Apis multiflorum*); *Cupressus* (e.g. *Cupressus sempervirens*, *Cupressus arizonica* and *Cupressus macrocarpa*); *Juniperus* (e.g. *Juniperus sabinooides*, *Juniperus virginiana*, *Juniperus communis* and *Juniperus ashei*); *Thuya* (e.g. *Thuya orientalis*); *Chamaecyparis* (e.g. *Chamaecyparis obtusa*); *Periplaneta* (e.g. *Periplaneta americana*); *Agropyron* (e.g. *Agropyron repens*); *Secale* (e.g. *Secale cereale*); *Triticum* (e.g. *Triticum aestivum*); *Dactylis* (e.g. *Dactylis glomerata*); *Festuca* (e.g. *Festuca elatior*); *Poa* (e.g. *Poa pratensis* or *Poa compressa*); *Avena* (e.g. *Avena sativa*); *Hol-*

cus (e.g. *Holcus lanatus*); *Anthoxanthum* (e.g. *Anthoxanthum odoratum*); *Arrhenatherum* (e.g. *Arrhenatherum elatius*); *Agrostis* (e.g. *Agrostis alba*); *Phleum* (e.g. *Phleum pratense*); *Phalaris* (e.g. *Phalaris arundinacea*); *Paspalum* (e.g. *Paspalum notatum*); *Sorghum* (e.g. *Sorghum halepensis*); and *Bromus* (e.g. *Bromus inermis*).

An “allergy” refers to acquired hypersensitivity to a substance (allergen). Allergic conditions include eczema, allergic rhinitis or coryza, hay fever, bronchial asthma, urticaria (hives) and food allergies, and other atopic conditions.

“Asthma”—refers to a disorder of the respiratory system characterized by inflammation, narrowing of the airways and increased reactivity of the airways to inhaled agents. Asthma is frequently, although not exclusively associated with atopic or allergic symptoms.

An “immune system deficiency” shall mean a disease or disorder in which the subject’s immune system is not functioning in normal capacity or in which it would be useful to boost a subject’s immune response for example to eliminate a tumor or cancer (e.g. tumors of the brain, lung (e.g. small cell and non-small cell), ovary, breast, prostate, colon, as well as other carcinomas and sarcomas) or an infection in a subject.

Examples of infectious virus include: Retroviridae (e.g., human immunodeficiency viruses, such as HIV-1 (also referred to as HTLV-III, LAV or HTLV-III/LAV, or HIV-III; and other isolates, such as HIV-LP; Picornaviridae (e.g., polio viruses, hepatitis A virus; enteroviruses, human coxsackie viruses, rhinoviruses, echoviruses); Calciviridae (e.g., strains that cause gastroenteritis); Togaviridae (e.g., equine encephalitis viruses, rubella viruses); Flaviridae (e.g., dengue viruses, encephalitis viruses, yellow fever viruses); Coronaviridae (e.g., coronaviruses); Rhabdoviridae (e.g., vesicular stomatitis viruses, rabies viruses); Filoviridae (e.g., ebola viruses); Paramyxoviridae (e.g., parainfluenza viruses, mumps virus, measles virus, respiratory syncytial virus); Orthomyxoviridae (e.g., influenza viruses); Bunyaviridae (e.g., Hantaan viruses, bunya viruses, phleboviruses and Nairo viruses); Arenaviridae (hemorrhagic fever viruses); Reoviridae (e.g., reoviruses, orbiviruses and rotaviruses); Birnaviridae; Hepadnaviridae (Hepatitis B virus); Parvoviridae (parvoviruses); Papovaviridae (papilloma viruses, polyoma viruses); Adenoviridae (most adenoviruses); Herpesviridae (herpes simplex virus (HSV) 1 and 2, varicella zoster virus, cytomegalovirus (CMV), herpes viruses); Poxviridae (variola viruses, vaccinia viruses, pox viruses); and Iridoviridae (e.g., African swine fever virus); and unclassified viruses (e.g., the etiological agents of Spongiform encephalopathies, the agent of delta hepatitis (thought to be a defective satellite of hepatitis B virus), the agents of non-A, non-B hepatitis (class 1=internally transmitted; class 2=parenterally transmitted (i.e., Hepatitis C); Norwalk and related viruses, and astroviruses).

Examples of infectious bacteria include: *Helicobacter pylori*, *Borelia burgdorferi*, *Legionella pneumophila*, *Mycobacteria* spp. (e.g., *M. tuberculosis*, *M. avium*, *M. intracellulare*, *M. kansasii*, *M. gordonae*), *Staphylococcus aureus*, *Neisseria gonorrhoeae*, *Neisseria meningitidis*, *Listeria monocytogenes*, *Streptococcus pyogenes* (Group A *Streptococcus*), *Streptococcus agalactiae* (Group B *Streptococcus*), *Streptococcus* (viridans group), *Streptococcus faecalis*, *Streptococcus bovis*, *Streptococcus* (anaerobic spp.), *Streptococcus pneumoniae*, pathogenic *Campylobacter* sp., *Enterococcus* sp., *Haemophilus influenzae*, *Bacillus anthracis*, *Corynebacterium diphtheriae*, *Corynebacterium* sp., *Erysipelothrix rhusiopathiae*, *Clostridium perfringens*, *Clostridium tetani*, *Enterobacter aerogenes*, *Klebsiella pneumoniae*, *Pasturella multocida*, *Bacteroides* sp., *Fusobacte-*

11

rium nucleatum, *Streptobacillus moniliformis*, *Treponema pallidum*, *Treponema pertenuis*, *Leptospira*, and *Actinomyces israelii*.

Examples of infectious fungi include: *Cryptococcus neoformans*, *Histoplasma capsulatum*, *Coccidioides immitis*, *Blastomyces dermatitidis*, *Chlamydia trachomatis*, *Candida albicans*. Other infectious organisms (i.e., protists) include: *Plasmodium falciparum* and *Toxoplasma gondii*.


An "immunostimulatory nucleic acid molecule" refers to a nucleic acid molecule, which contains an unmethylated cytosine, guanine dinucleotide sequence (i.e. "CpG DNA" or DNA containing a cytosine followed by guanosine and linked by a phosphate bond) and stimulates (e.g. has a mitogenic effect on, or induces or increases cytokine expression by) a vertebrate lymphocyte. An immunostimulatory nucleic acid molecule can be double-stranded or single-stranded. Generally, double-stranded molecules are more stable in vivo, while single-stranded molecules have increased immune activity.

In a preferred embodiment, the immunostimulatory nucleic acid contains a consensus mitogenic CpG motif represented by the formula:



wherein X_1 is selected from the group consisting of A, G and T; and X_2 is C or T.

In a particularly preferred embodiment, immunostimulatory nucleic acid molecules are between 2 to 100 base pairs in size and contain a consensus mitogenic CpG motif represented by the formula:



wherein C and G are unmethylated, X_1 , X_2 , X_3 and X_4 are nucleotides.

For economic reasons, preferably the immunostimulatory CpG DNA is in the range of between 8 to 40 base pairs in size if it is synthesized as an oligonucleotide. Alternatively, CpG dinucleotides can be produced on a large scale in plasmids, which after being administered to a subject are degraded into oligonucleotides. Preferred immunostimulatory nucleic acid molecules (e.g. for use in increasing the effectiveness of a vaccine or to treat an immune system deficiency by stimulating an antibody [humoral] response in a subject) have a relatively high stimulation index with regard to B cell, monocyte and/or natural killer cell responses (e.g. cytokine, proliferative, lytic or other responses).

The stimulation index of a particular immunostimulatory CpG DNA can be tested in various immune cell assays. Preferably, the stimulation index of the immunostimulatory CpG DNA with regard to B-cell proliferation is at least about 5, preferably at least about 10, more preferably at least about 15 and most preferably at least about 20 as determined by incorporation of 3H uridine in a murine B cell culture, which has been contacted with a 20 μM of ODN for 20 h at 37° C. and has been pulsed with 1 μCi of 3H uridine; and harvested and counted 4 h later as described in detail in Example 1. For use in vivo, for example to treat an immune system deficiency by stimulating a cell-mediated (local) immune response in a subject, it is important that the immunostimulatory CpG DNA be capable of effectively inducing cytokine secretion by monocytic cells and/or Natural Killer (NK) cell lytic activity.

Preferred immunostimulatory CpG nucleic acids should effect at least about 500 pg/ml of TNF- α , 15 pg/ml IFN- γ , 70 pg/ml of GM-CSF 275 pg/ml of IL-6, 200 pg/ml IL-12, depending on the therapeutic indication, as determined by the

12

assays described in Example 12. Other preferred immunostimulatory CpG DNAs should effect at least about 10%, more preferably at least about 15% and most preferably at least about 20% YAC-1 cell specific lysis or at least about 30, more preferably at least about 35 and most preferably at least about 40% 2C11 cell specific lysis as determined by the assay described in detail in Example 4.

A "nucleic acid" or "DNA" shall mean multiple nucleotides (i.e. molecules comprising a sugar (e.g. ribose or deoxyribose) linked to a phosphate group and to an exchangeable organic base, which is either a substituted pyrimidine (e.g. cytosine (C), thymine (T) or uracil (U)) or a substituted purine (e.g. adenine (A) or guanine (G)). As used herein, the term refers to ribonucleotides as well as oligodeoxyribonucleotides. The term shall also include polynucleotides (i.e. a polynucleotide minus the phosphate) and any other organic base containing polymer. Nucleic acid molecules can be obtained from existing nucleic acid sources (e.g. genomic or cDNA), but are preferably synthetic (e.g. produced by oligonucleotide synthesis).

A "nucleic acid delivery complex" shall mean a nucleic acid molecule associated with (e.g. ionically or covalently bound to; or encapsulated within) a targeting means (e.g. a molecule that results in higher affinity binding to target cell (e.g. B-cell and natural killer (NK) cell) surfaces and/or increased cellular uptake by target cells). Examples of nucleic acid delivery complexes include nucleic acids associated with: a sterol (e.g. cholesterol), a lipid (e.g. a cationic lipid, virosome or liposome), or a target cell specific binding agent (e.g. a ligand recognized by target cell specific receptor). Preferred complexes must be sufficiently stable in vivo to prevent significant uncoupling prior to internalization by the target cell. However, the complex should be cleavable under appropriate conditions within the cell so that the nucleic acid is released in a functional form.

"Palindromic sequence" shall mean an inverted repeat (i.e. a sequence such as ABCDEE'D'C'B'A' in which A and A' are bases capable of forming the usual Watson-Crick base pairs. In vivo, such sequences may form double stranded structures.

A "stabilized nucleic acid molecule" shall mean a nucleic acid molecule that is relatively resistant to in vivo degradation (e.g. via an exo- or endo-nuclease). Stabilization can be a function of length or secondary structure. Unmethylated CpG containing nucleic acid molecules that are tens to hundreds of kbs long are relatively resistant to in vivo degradation. For shorter immunostimulatory nucleic acid molecules, secondary structure can stabilize and increase their effect. For example, if the 3' end of a nucleic acid molecule has self-complementarity to an upstream region, so that it can fold back and form a sort of stem loop structure, then the nucleic acid molecule becomes stabilized and therefore exhibits more activity.

Preferred stabilized nucleic acid molecules of the instant invention have a modified backbone. For use in immune stimulation, especially preferred stabilized nucleic acid molecules are phosphorothioate modified nucleic acid molecules (i.e. at least one of the phosphate oxygens of the nucleic acid molecule is replaced by sulfur). Preferably the phosphate modification occurs at or near the 5' and/or 3' end of the nucleic acid molecule. In addition to stabilizing nucleic acid molecules, as reported further herein, phosphorothioate-modified nucleic acid molecules (including phosphorodithioate-modified) can increase the extent of immune stimulation of the nucleic acid molecule, which contains an unmethylated CpG dinucleotide as shown herein. International Patent Application Publication Number: WO 95/26204 entitled "Immune Stimulation By Phosphorothioate Oligonucleotide

Analogous” also reports on the non-sequence specific immunostimulatory effect of phosphorothioate modified oligonucleotides. As reported herein, unmethylated CpG containing nucleic acid molecules having a phosphorothioate backbone have been found to preferentially activate B-cell activity, while unmethylated CpG containing nucleic acid molecules having a phosphodiester backbone have been found to preferentially activate monocytic (macrophages, dendritic cells and monocytes) and NK cells. Phosphorothioate CpG oligonucleotides with preferred human motifs are also strong activators of monocytic and NK cells.

Other stabilized nucleic acid molecules include: nonionic DNA analogs, such as alkyl- and aryl-phosphonates (in which the charged phosphonate oxygen is replaced by an alkyl or aryl group), phosphodiester and alkylphosphotriesters, in which the charged oxygen moiety is alkylated. Nucleic acid molecules which contain a diol, such as tetraethyleneglycol or hexaethyleneglycol, at either or both termini have also been shown to be substantially resistant to nuclease degradation.

A “subject” shall mean a human or vertebrate animal including a dog, cat, horse, cow, pig, sheep, goat, chicken, monkey, rat, mouse, etc.

As used herein, the term “vector” refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. Preferred vectors are those capable of autonomous replication and expression of nucleic acids to which they are linked (e.g., an episome). Vectors capable of directing the expression of genes to which they are operatively linked are referred to herein as “expression vectors.” In general, expression vectors of utility in recombinant DNA techniques are often in the form of “plasmids” which refer generally to circular double stranded DNA loops which, in their vector form, are not bound to the chromosome. In the present specification, “plasmid” and “vector” are used interchangeably as the plasmid is the most commonly used form of vector. However, the invention is intended to include such other forms of expression vectors which serve equivalent functions and which become known in the art subsequently hereto.

Certain Unmethylated CpG Containing Nucleic Acids have B Cell Stimulatory Activity as Shown In Vitro and In Vivo

In the course of investigating the lymphocyte stimulatory effects of two antisense oligonucleotides specific for endogenous retroviral sequences, using protocols described in the attached Examples 1 and 2, it was surprisingly found that two out of twenty-four “controls” (including various scrambled, sense, and mismatch controls for a panel of “antisense” ODN) also mediated B cell activation and IgM secretion, while the other “controls” had no effect.

Two observations suggested that the mechanism of this B cell activation by the “control” ODN may not involve antisense effects 1) comparison of vertebrate DNA sequences listed in GenBank showed no greater homology than that seen with non-stimulatory ODN and 2) the two controls showed no hybridization to Northern blots with 10 µg of spleen poly A+ RNA. Resynthesis of these ODN on a different synthesizer or extensive purification by polyacrylamide gel electrophoresis or high pressure liquid chromatography gave identical stimulation, eliminating the possibility of an impurity. Similar stimulation was seen using B cells from C3H/HeJ mice, eliminating the possibility that lipopolysaccharide (LPS) contamination could account for the results.

The fact that two “control” ODN caused B cell activation similar to that of the two “antisense” ODN raised the possibility that all four ODN were stimulating B cells through

some non-antisense mechanism involving a sequence motif that was absent in all of the other nonstimulatory control ODN. In comparing these sequences, it was discovered that all of the four stimulatory ODN contained CpG dinucleotides that were in a different sequence context from the nonstimulatory control.

To determine whether the CpG motif present in the stimulatory ODN was responsible for the observed stimulation, over 300 ODN ranging in length from 5 to 42 bases that contained methylated, unmethylated, or no CpG dinucleotides in various sequence contexts were synthesized. These ODNs, including the two original “controls” (ODN 1 and 2) and two originally synthesized as “antisense” (ODN 3D and 3M; Krieg, A. M. *J. Immunol.* 143:2448 (1989)), were then examined for in vitro effects on spleen cells (representative sequences are listed in Table 1). Several ODN that contained CpG dinucleotides induced B cell activation and IgM secretion; the magnitude of this stimulation typically could be increased by adding more CpG dinucleotides (Table 1; compare ODN 2 to 2a or 3D to 3Da and 3Db). Stimulation did not appear to result from an antisense mechanism or impurity. ODN caused no detectable proliferation of γδ or other T cell populations.

Mitogenic ODN sequences uniformly became nonstimulatory if the CpG dinucleotide was mutated (Table 1; compare ODN 1 to 1a; 3D to 3Dc; 3M to 3Ma; and 4 to 4a) or if the cytosine of the CpG dinucleotide was replaced by 5-methylcytosine (Table 1; ODN 1b, 2b, 3Dd, and 3Mb). Partial methylation of CpG motifs caused a partial loss of stimulatory effect (compare 2a to 2c, Table 1). In contrast, methylation of other cytosines did not reduce ODN activity (ODN 1c, 2d, 3De and 3Mc). These data confirmed that a CpG motif is the essential element present in ODN that activate B cells.

In the course of these studies, it became clear that the bases flanking the CpG dinucleotide played an important role in determining the murine B cell activation induced by an ODN. The optimal stimulatory motif was determined to consist of a CpG flanked by two 5' purines (preferably a GpA dinucleotide) and two 3' pyrimidines (preferably a TpT or TpC dinucleotide). Mutations of ODN to bring the CpG motif closer to this ideal improved stimulation (e.g. Table 1, compare ODN 2 to 2e; 3M to 3Md) while mutations that disturbed the motif reduced stimulation (e.g. Table 1, compare ODN 3D to 3Df; 4 to 4b, 4c and 4d). On the other hand, mutations outside the CpG motif did not reduce stimulation (e.g. Table 1, compare ODN 1 to 1d; 3D to 3Dg; 3M to 3Me). For activation of human cells, the best flanking bases are slightly different (See Table 5).

Of those tested, ODNs shorter than 8 bases were non-stimulatory (e.g. Table 1, ODN 4e). Among the forty-eight 8 base ODN tested, the most stimulatory sequence identified was TCAACGTT (ODN 4) which contains the self complementary “palindrome” AACGTT. In further optimizing this motif, it was found that ODN containing Gs at both ends showed increased stimulation, particularly if the ODN were rendered nuclease resistant by phosphorothioate modification of the terminal internucleotide linkages. ODN 1585 (5' GGGGTCAACGTTGAGGGGGG 3' (SEQ ID NO:12), in which the first two and last five internucleotide linkages are phosphorothioate modified caused an average 25.4 fold increase in mouse spleen cell proliferation compared to an average 3.2 fold increase in proliferation induced by ODN 1638, which has the same sequence as ODN 1585 except that the 10 Gs at the two ends are replaced by 10 As. The effect of the G-rich ends is cis; addition of an ODN with poly G ends

but no CpG motif to cells along with 1638 gave no increased proliferation. For nucleic acid molecules longer than 8 base pairs, non-palindromic motifs containing an unmethylated CpG were found to be more immunostimulatory.

Other octamer ODN containing a 6 base palindrome with a TpC dinucleotide at the 5' end were also active (e.g. Table 1,

ODN 4b, 4c). Other dinucleotides at the 5' end gave reduced stimulation (e.g. ODN 4f; all sixteen possible dinucleotides were tested). The presence of a 3' dinucleotide was insufficient to compensate for the lack of a 5' dinucleotide (e.g. Table 1, ODN 4g). Disruption of the palindrome eliminated stimulation in octamer ODN (e.g. Table 1, ODN 4h), but palindromes were not required in longer ODN.

TABLE 1

Oligonucleotide Stimulation of Mouse B Cells			
ODN	Sequence (5' to 3')†	Stimulation Index'	
		³ H Uridine	IgM Production
1 (SEQ ID NO: 13)	GCTAGAC <u>CG</u> TTAGCGT	6.1 ± 0.8	17.9 ± 3.6
1a (SEQ ID NO: 4)T.....	1.2 ± 0.2	1.7 ± 0.5
1b (SEQ ID NO: 14)Z.....	1.2 ± 0.1	1.8 ± 0.0
1c (SEQ ID NO: 15)Z..	10.3 ± 4.4	9.5 ± 1.8
1d (SEQ ID NO: 16)	..AT.....GAGC.	13.0 ± 2.3	18.3 ± 7.5
2 (SEQ ID NO: 17)	ATGGAAGGTCCAG <u>CG</u> TTCTC	2.9 ± 0.2	13.6 ± 2.0
2a (SEQ ID NO: 18)	..C..CTC..G.....	7.7 ± 0.8	24.2 ± 3.2
2b (SEQ ID NO: 19)	..Z..CTC.ZG..Z.....	1.6 ± 0.5	2.8 ± 2.2
2c (SEQ ID NO: 20)	..Z..CTC..G.....	3.1 ± 0.6	7.3 ± 1.4
2d (SEQ ID NO: 21)	..C..CTC..G.....Z..	7.4 ± 1.4	27.7 ± 5.4
2e (SEQ ID NO: 22)A.....	5.6 ± 2.0	ND
3D (SEQ ID NO: 23)	GAGA <u>AC</u> GCTGGACCTCCAT	4.9 ± 0.5	19.9 ± 3.6
3Da (SEQ ID NO: 24)C.....	6.6 ± 1.5	33.9 ± 6.8
3Db (SEQ ID NO: 25)C.....G..	10.1 ± 2.8	25.4 ± 0.8
3Dc (SEQ ID NO: 26)	...C.A.....	1.0 ± 0.1	1.2 ± 0.5
3Dd (SEQ ID NO: 27)Z.....	1.2 ± 0.2	1.0 ± 0.4
3De (SEQ ID NO: 28)Z.....	4.4 ± 1.2	18.8 ± 4.4
3Df (SEQ ID NO: 29)A.....	1.6 ± 0.1	7.7 ± 0.4
3Dg (SEQ ID NO: 30)CC.G.ACTG..	6.1 ± 1.5	18.6 ± 1.5
3M (SEQ ID NO: 31)	TCCATGT <u>CG</u> GCCTGATGCT	4.1 ± 0.2	23.2 ± 4.9
3Ma (SEQ ID NO: 32)CT.....	0.9 ± 0.1	1.8 ± 0.5
3Mb (SEQ ID NO: 33)Z.....	1.3 ± 0.3	1.5 ± 0.6
3Mc (SEQ ID NO: 34)Z.....	5.4 ± 1.5	8.5 ± 2.6
3Md (SEQ ID NO: 35)A..T.....	17.2 ± 9.4	ND
3Me (SEQ ID NO: 36)C..A..	3.6 ± 0.2	14.2 ± 5.2
4	TCAACGTT	6.1 ± 1.4	19.2 ± 5.2
4aGC..	1.1 ± 0.2	1.5 ± 1.1
4b	...G <u>CG</u> C.	4.5 ± 0.2	9.6 ± 3.4
4c	...TCGA.	2.7 ± 1.0	ND
4d	..TT..AA	1.3 ± 0.2	ND
4e	1.3 ± 0.2	1.1 ± 0.5
4f	C.....	3.9 ± 1.4	ND
4g	---...CT	1.4 ± 0.3	ND
4hC	1.2 ± 0.2	ND
LPS		7.8 ± 2.5	4.8 ± 1.0

'Stimulation indexes are the means and std. dev. derived from at least 3 separate experiments, and are compared to wells cultured with no added ODN.

ND = not done.

CpG dinucleotides are underlined.

Dots indicate identity; dashes indicate deletions.

Z indicates 5 methyl cytosine.

TABLE 2

Identification of the optimal CpG motif for Murine IL-6 production and B cell activation.					
ODN	SEQUENCE (5'-3')	IL-6 (pgl/l) ^a			IgM (ng/ml) ^c
		CHI2.LX	SPLENIC B CELL	SI ^b	
512 (SEQ ID NO: 37)	TCCATGTCGCTCCTGATGCT	1300 ± 106	627 ± 43	5.8 ± 0.3	7315 ± 1324
1637 (SEQ ID NO: 38)C.....	136 ± 27	46 ± 6	1.7 ± 0.2	770 ± 72
1615 (SEQ ID NO: 39)G.....	1201 ± 155	850 ± 202	3.7 ± 0.3	3212 ± 617
1614 (SEQ ID NO: 40)A.....	1533 ± 321	1812 ± 103	10.8 ± 0.6	7558 ± 414
1636 (SEQ ID NO: 41)A.....	1181 ± 76	947 ± 132	5.4 ± 0.4	3983 ± 485
1634 (SEQ ID NO: 42)C.....	1049 ± 223	1671 ± 175	9.2 ± 0.9	6256 ± 261
1619 (SEQ ID NO: 43)T.....	1555 ± 304	2908 ± 129	12.5 ± 1.0	8243 ± 698
1618 (SEQ ID NO: 44)A..T.....	2109 ± 291	2596 ± 166	12.9 ± 0.7	10425 ± 674
1639 (SEQ ID NO: 45)AA..T.....	1827 ± 83	2012 ± 132	11.5 ± 0.4	9489 ± 103
1707 (SEQ ID NO: 46)A..TC.....	ND	1147 ± 175	4.0 ± 0.2	3534 ± 217
1708 (SEQ ID NO: 47)CA..TG.....	ND	59 ± 3	1.5 ± 0.1	466 ± 109

Dots indicate identity; CpG dinucleotides are underlined; ND = not done

^aThe experiment was done at least three times with similar results. The level of IL-6 of unstimulated control cultures of both CHI2.LX and splenic B cells was ≤ 10 pg/ml. The IgM level of unstimulated culture was 547 ± 82 ng/ml. CpG dinucleotides are underlined and dots indicate identity.

^b[³H] Uridine uptake was indicated as a fold increase (SI: stimulation index) from unstimulated control (2322.67 ± 213.68 cpm). Cells were stimulated with 20 μ M of various CpG O-ODN. Data present the mean \pm SD of triplicates

^cMeasured by ELISA.

The kinetics of lymphocyte activation were investigated using mouse spleen cells. When the cells were pulsed at the same time as ODN addition and harvested just four hours later, there was already a two-fold increase in ³H uridine incorporation. Stimulation peaked at 12-48 hours and then decreased. After 24 hours, no intact ODN were detected, perhaps accounting for the subsequent fall in stimulation when purified B cells with or without anti-IgM (at a submitogenic dose) were cultured with CpG ODN, proliferation was found to synergistically increase about 10-fold by the two mitogens in combination after 48 hours. The magnitude of stimulation was concentration dependent and consistently exceeded that of LPS under optimal conditions for both. Oligonucleotides containing a nuclease resistant phosphorothioate backbone were approximately two hundred times more potent than unmodified oligonucleotides.

Cell cycle analysis was used to determine the proportion of B cells activated by CpG-ODN. CpG-ODN induced cycling in more than 95% of B cells. Splenic B lymphocytes sorted by flow cytometry into CD23- (marginal zone) and CD23+ (follicular) subpopulations were equally responsive to ODN-induced stimulation, as were both resting and activated populations of B cells isolated by fractionation over Percoll gradients. These studies demonstrated that CpG-ODN induce essentially all B cells to enter the cell cycle.

Immunostimulatory Nucleic Acid Molecules Block Murine B Cell Apoptosis

Certain B cell lines such as WEHI-231 are induced to undergo growth arrest and/or apoptosis in response to crosslinking of their antigen receptor by anti-IgM (Jakway, J. P. et al., "Growth regulation of the B lymphoma cell line WEHI-231 by anti-immunoglobulin, lipopolysaccharide and other bacterial products" *J. Immunol.* 137: 2225 (1986); Tsubata, T., J. Wu and T. Honjo: B-cell apoptosis induced by

antigen receptor crosslinking is blocked by a T-cell signal through CD40." *Nature* 364: 645 (1993)). WEHI-231 cells are rescued from this growth arrest by certain stimuli such as LPS and by the CD40 ligand. ODN containing the CpG motif were also found to protect WEHI-231 from anti-IgM induced growth arrest, indicating that accessory cell populations are not required for the effect. Subsequent work indicates that CpG ODN induce Bcl-x and myc expression, which may account for the protection from apoptosis. Also, CpG nucleic acids have been found to block apoptosis in human cells. This inhibition of apoptosis is important, since it should enhance and prolong immune activation by CpG DNA.

Induction of Murine Cytokine Secretion by CpG Motifs in Bacterial DNA or Oligonucleotides.

As described in Example 9, the amount of IL-6 secreted by spleen cells after CpG DNA stimulation was measured by ELISA. T cell depleted spleen cell cultures rather than whole spleen cells were used for in vitro studies following preliminary studies showing that T cells contribute little or nothing to the IL-6 produced by CpG DNA-stimulated spleen cells. As shown in Table 3, IL-6 production was markedly increased in cells cultured with *E. coli* DNA but not in cells cultured with calf thymus DNA. To confirm that the increased IL-6 production observed with *E. coli* DNA was not due to contamination by other bacterial products, the DNA was digested with DNase prior to analysis. DNase pretreatment abolished IL-6 production induced by *E. coli* DNA (Table 3). In addition, spleen cells from LPS-nonresponsive C3H/HeJ mouse produced similar levels of IL-6 in response to bacterial DNA. To analyze whether the IL-6 secretion induced by *E. coli* DNA was mediated by the unmethylated CpG dinucleotides in bacterial DNA, methylated *E. coli* DNA and a panel of synthetic ODN were examined. As shown in Table 3, CpG ODN significantly induced IL-6 secretion (ODN 5a, 5b, 5c) while

CpG methylated *E. coli* DNA, or ODN containing methylated CpG (ODN 50 or no CpG (ODN 5d) did not. Changes at sites other than CpG dinucleotides (ODN 5b) or methylation of other cytosines (ODN 5g) did not reduce the effect of CpG ODN. Methylation of a single CpG in an ODN with three CpGs resulted in a partial reduction in the stimulation (compare ODN 5c to 5e; Table 3).

IL-6 production plateaued at approximately 50 µg/ml of bacterial DNA or 40 µM of CpG O-ODN. The maximum levels of IL-6 induced by bacterial DNA and CpG ODN were 1-1.5 ng/ml and 2-4 ng/ml respectively. These levels were significantly greater than those seen after stimulation by LPS (0.35 ng/ml) (FIG. 1A). To evaluate whether CpG ODN with a nuclease-resistant DNA backbone would also induce IL-6

TABLE 3

Induction of Murine IL-6 secretion by CpG motifs in bacterial DNA or oligonucleotides.	
Treatment	IL-6 (pg/ml)
calf thymus DNA	≤10
calf thymus DNA + DNase	≤10
<i>E. coli</i> DNA	1169.5 ± 94.1
<i>E. coli</i> DNA + DNase	≤10
CpG methylated <i>E. coli</i> DNA	≤10
LPS	280.1 ± 17.1
Media (no DNA)	≤10
ODN 5a SEQ ID NO: 59 ATGGACTCTCCAG <u>C</u> GTTC	1096.4 ± 372.0
5b SEQ ID NO: 22AGG.A.	1124.5 ± 126.2
5c SEQ ID NO: 18 . . <u>C</u> <u>G</u>	1783.0 ± 189.5
5d SEQ ID NO: 1AGG. . C. . T.	≤10
5e SEQ ID NO: 60 . . <u>C</u>G.Z.	851.1 ± 114.4
5f SEQ ID NO: 19 . . Z.ZG. . Z.	≤10
5g SEQ ID NO: 21 . . <u>C</u> <u>G</u>Z.	1862.3 ± 87.26

T cell depleted spleen cells from DBA/2 mice were stimulated with phosphodiester modified oligonucleotides (O-ODN) (20 µM), calf thymus DNA (50 µg/ml) or *E. coli* DNA (50 µg/ml) with or without enzyme treatment, or LPS (10 µg/ml) for 24 hr. Data represent the mean (pg/ml) ± SD of triplicates. CpG dinucleotides are underlined and dots indicate identity. Z indicates 5-methylcytosine.

Identification of the Optimal CpG Motif for Induction of Murine IL-6 and IgM Secretion and B Cell Proliferation.

To evaluate whether the optimal B cell stimulatory CpG motif was identical with the optimal CpG motif for IL-6 secretion, a panel of ODN in which the bases flanking the CpG dinucleotide were progressively substituted was studied. This ODN panel was analyzed for effects on B cell proliferation, Ig production, and IL-6 secretion, using both splenic B cells and CH12.LX cells. As shown in Table 2, the optimal stimulatory motif is composed of an unmethylated CpG flanked by two 5' purines and two 3' pyrimidines. Generally a mutation of either 5' purine to pyrimidine or 3' pyrimidine to purine significantly reduced its effects. Changes in 5' purines to C were especially deleterious, but changes in 5' purines to T or 3' pyrimidines to purines had less marked effects. Based on analyses of these and scores of other ODN, it was determined that the optimal CpG motif for induction of IL-6 secretion is TGACGTT, which is identical with the optimal mitogenic and IgM-inducing CpG motif (Table 2). This motif was more stimulatory than any of the palindrome containing sequences studied (1639, 1707 and 1708).

Titration of Induction of Murine IL-6 Secretion by CpG Motifs.

Bacterial DNA and CpG ODN induced IL-6 production in T cell depleted murine spleen cells in a dose-dependent manner, but vertebrate DNA and non-CpG ODN did not (FIG. 1).

production, S-ODN were added to T cell depleted murine spleen cells. CpG S-ODN also induced IL-6 production in a dose-dependent manner to approximately the same level as CpG O-ODN while non-CpG S-ODN failed to induce IL-6 (FIG. 1C). CpG S-ODN at a concentration of 0.05 µM could induce maximal IL-6 production in these cells. This result indicated that the nuclease-resistant DNA backbone modification retains the sequence specific ability of CpG DNA to induce IL-6 secretion and that CpG S-ODN are more than 80-fold more potent than CpG O-ODN in this assay system.

Induction of Murine IL-6 Secretion by CpG DNA In Vivo.

To evaluate the ability of bacterial DNA and CpG S-ODN to induce IL-6 secretion in vivo, BALB/c mice were injected iv. with 100 µg of *E. coli* DNA, calf thymus DNA, or CpG or non-stimulatory S-ODN and bled 2 hr after stimulation. The level of IL-6 in the sera from the *E. coli* DNA injected group was approximately 13 ng/ml while IL-6 was not detected in the sera from calf thymus DNA or PBS injected groups (Table 4). CpG S-ODN also induced IL-6 secretion in vivo. The IL-6 level in the sera from CpG S-ODN injected groups was approximately 20 ng/ml. In contrast, IL-6 was not detected in the sera from non-stimulatory S-ODN stimulated group (Table 4).

TABLE 4

Secretion of Murine IL-6 induced by CpG DNA stimulation <i>in vivo</i> .	
Stimulant	IL-6 (pg/ml)
PBS	<50
<i>E. coli</i> DNA	13858 ± 3143
Calf Thymus DNA	<50
CpG S-ODN	20715 ± 606
non-CpG S-ODN	<50

Mice (2 mice/group) were i.v. injected with 100 µl of PBS, 200 µg of *E. coli* DNA or calf thymus DNA, or 500 µg of CpG S-ODN or non-CpG control S-ODN. Mice were bled 2 hr after injection and 1:10 dilution of each serum was analyzed by IL-6 ELISA. Sensitivity limit of IL-6 ELISA was 5 pg/ml. Sequences of the CpG S-ODN is 5'GCATGACGTTGAGCT3' (SEQ ID NO: 48) and of the non-stimulatory S-ODN is 5'GCTAGATGTTAGCGT3' (SEQ ID NO: 49). Note that although there is a CpG in sequence 48, it is too close to the 3' end to effect stimulation, as explained herein. Data represent mean ± SD of duplicates. The experiment was done at least twice with similar results.

Kinetics of Murine IL-6 Secretion after Stimulation by CpG Motifs *In Vivo*.

To evaluate the kinetics of induction of IL-6 secretion by CpG DNA *in vivo*, BALB/c mice were injected iv. with CpG or control non-CpG S-ODN. Serum IL-6 levels were significantly increased within 1 hr and peaked at 2 hr to a level of approximately 9 ng/ml in the CpG S-ODN injected group (FIG. 2). IL-6 protein in sera rapidly decreased after 4 hr and returned to basal level by 12 hr after stimulation. In contrast to CpG DNA stimulated groups, no significant increase of IL-6 was observed in the sera from the non-stimulatory S-ODN or PBS injected groups (FIG. 2).

Tissue Distribution and Kinetics of IL-6 mRNA Expression Induced by CpG Motifs *In Vivo*.

As shown in FIG. 2, the level of serum IL-6 increased rapidly after CpG DNA stimulation. To investigate the possible tissue origin of this serum IL-6, and the kinetics of IL-6 gene expression *in vivo* after CpG DNA stimulation, BALB/c mice were injected iv with CpG or non-CpG S-ODN and RNA was extracted from liver, spleen, thymus, and bone marrow at various time points after stimulation. As shown in FIG. 3A, the level of IL-6 mRNA in liver, spleen, and thymus was increased within 30 min. after injection of CpG S-ODN. The liver IL-6 mRNA peaked at 2 hr post-injection and rapidly decreased and reached basal level 8 hr after stimulation (FIG. 3A). Splenic IL-6 mRNA peaked at 2 hr after stimulation and then gradually decreased (FIG. 3A). Thymus IL-6 mRNA peaked at 1 hr post-injection and then gradually decreased (FIG. 3A). IL-6 mRNA was significantly increased in bone marrow within 1 hr after CpG S-ODN injection but then returned to basal level. In response to CpG S-ODN, liver, spleen and thymus showed more substantial increases in IL-6 mRNA expression than the bone marrow.

Patterns of Murine Cytokine Expression Induced by CpG DNA

In vivo or in whole spleen cells, no significant increase in the protein levels of the following interleukins: IL-2, IL-3, IL-4, IL-5, or IL-10 was detected within the first six hours (Klinman, D. M. et al., (1996) *Proc. Natl. Acad. Sci. USA* 93:2879-2883). However, the level of TNF-α is increased within 30 minutes and the level of IL-6 increased strikingly within 2 hours in the serum of mice injected with CpG ODN. Increased expression of IL-12 and interferon gamma (IFN-γ) mRNA by spleen cells was also detected within the first two hours.

TABLE 5

Induction of human PBMC cytokine secretion by CpG oligos						
ODN	Sequence (5'-3')	IL-6 ¹	TNF-α ¹	IFN-γ ¹	GM-CSF	IL-12
512 SEQ ID NO: 37	TCCATGTC <u>CG</u> TCCTGATGCT	500	140	15.6	70	250
1637 SEQ ID NO: 38C.....	550	16	7.8	15.6	35
1615 SEQ ID NO: 39G.....	600	145	7.8	45	250
1614 SEQ ID NO: 40A.....	550	31	0	50	250
1636 SEQ ID NO: 41A.....	325	250	35	40	0
1634 SEQ ID NO: 42C.....	300	400	40	85	200
1619 SEQ ID NO: 43T.....	275	450	200	80	>500
1618 SEQ ID NO: 44A..T.....	300	60	15.6	15.6	62
1639 SEQ ID NO: 45AA..T.....	625	220	15.6	40	60
1707 SEQ ID NO: 46A..TC.....	300	70	17	0	0

TABLE 5-continued

Induction of human PBMC cytokine secretion by CpG oligos						
ODN	Sequence (5'-3')	IL-6 ¹	TNF- α ¹	IFN- γ ¹	GM-CSF	IL-12
1708CA..TG.....	270	10	17	0	0

dots indicate identity; CpG dinucleotides are underlined

¹measured by ELISA using Quantikine kits from R&D Systems (pg/ml) Cells were cultured in 10% autologous serum with the indicated oligodeoxynucleotides (12 μ g/ml) for 4 hr in the case of TNF- α or 24 hr for the other cytokines before supernatant harvest and assay. Data are presented as the level of cytokine above that in wells with no added oligodeoxynucleotide.

¹measured by ELISA using Quantikine kits from R&D SYSTEMS® (pg/ml) Cells were cultured in 10% autologous serum with the indicated oligodeoxynucleotides (12 μ g/ml) for 4 hr in the case of TNF- α or 24 hr for the other cytokines before supernatant harvest and assay. Data are presented as the level of cytokine above that in wells with no added oligodeoxynucleotide.

CpG DNA Induces Cytokine Secretion by Human PBMC, Specifically Monocytes

The same panels of ODN used for studying mouse cytokine expression were used to determine whether human cells also are induced by CpG motifs to express cytokine (or proliferate), and to identify the CpG motif(s) responsible. Oligonucleotide 1619 (GTCGTT) was the best inducer of TNF- α and IFN- γ secretion, and was closely followed by a nearly identical motif in oligonucleotide 1634 (GTCGCT) (Table 5). The motifs in oligodeoxynucleotides 1637 and 1614 (GC-CGGT and GACGGT) led to strong IL-6 secretion with relatively little induction of other cytokines. Thus, it appears that human lymphocytes, like murine lymphocytes, secrete cytokines differentially in response to CpG dinucleotides, depending on the surrounding bases. Moreover, the motifs that stimulate murine cells best differ from those that are most effective with human cells. Certain CpG oligodeoxynucleotides are poor at activating human cells (oligodeoxynucleotides 1707, 1708, which contain the palindrome forming sequences GACGTC and CACGTG respectively).

The cells responding to the DNA appear to be monocytes, since the cytokine secretion is abolished by treatment of the cells with L-leucyl-L-leucine methyl ester (L-LME), which is selectively toxic to monocytes (but also to cytotoxic T lymphocytes and NK cells), and does not affect B cell Ig secretion (Table 6, and data not shown). The cells surviving L-LME treatment had >95% viability by trypan blue exclusion, indicating that the lack of a cytokine response among these cells did not simply reflect a nonspecific death all cell types. Cytokine secretion in response to *E. coli* (EC) DNA requires unmethylated CpG motifs, since it is abolished by methylation of the EC DNA (next to the bottom row, Table 6). LPS contamination of the DNA cannot explain the results since the level of contamination was identical in the native and methylated DNA, and since addition of twice the highest amount of contaminating LPS had no effect (not shown).

TABLE 6

CpG DNA induces cytokine secretion by human PBMC				
DNA	TNF- α (pg/ml) ¹	IL-6 (pg/ml)	IFN- γ (pg/ml)	RANTES (pg/ml)
EC DNA (50 μ g/ml)	900	12,000	700	1560
EC DNA (5 μ g/ml)	850	11,000	400	750
EC DNA (0.5 μ g/ml)	500	ND	200	0

TABLE 6-continued

CpG DNA induces cytokine secretion by human PBMC				
DNA	TNF- α (pg/ml) ¹	IL-6 (pg/ml)	IFN- γ (pg/ml)	RANTES (pg/ml)
EC DNA (0.05 μ g/ml)	62.5	10,000	15.6	0
EC DNA (50 μ g/ml) + L-LME ²	0	ND	ND	ND
EC DNA (10 μ g/ml) Methyl ³	0	5	ND	ND
CT DNA (50 μ g/ml)	0	600	0	0

¹Levels of all cytokines were determined by ELISA using Quantikine kits from R&D Systems as described in the previous table. Results are representative using PBMC from different donors.

²Cells were pretreated for 15 min. with L-leucyl-L-leucine methyl ester (L-LME) to determine whether the cytokine production under these conditions was from monocytes (or other L-LME-sensitive cells).

³EC DNA was methylated using 2 U/ μ g DNA of CpG methylase (New England Biolabs) according to the manufacturer's directions, and methylation confirmed by digestion with Hpa-II and Msp-I. As a negative control, samples were included containing twice the maximal amount of LPS contained in the highest concentration of EC DNA which failed to induce detectable cytokine production under these experimental conditions. ND = not done

¹Levels of all cytokines were determined by ELISA using Quantikine kits from R&D SYSTEMS® as described in the previous table. Results are representative using PBMC from different donors.

³EC DNA was methylated using 2 U/ μ g DNA of CpG methylase (NEW ENGLAND BIOLABS® Inc.) according to the manufacturer's directions, and methylation confirmed by digestion with Hpa-II and Msp-I. As a negative control, samples were included containing twice the maximal amount of LPS contained in the highest concentration of EC DNA which failed to induce detectable cytokine production under these experimental conditions. ND=not done

The loss of cytokine production in the PBMC treated with L-LME suggested that monocytes may be responsible for cytokine production in response to CpG DNA. To test this hypothesis more directly, the effects of CpG DNA on highly purified human monocytes and macrophages was tested. As hypothesized, CpG DNA directly activated production of the cytokines IL-6, GM-CSF, and TNF- α by human macrophages, whereas non-CpG DNA did not (Table 7).

TABLE 7

CpG DNA induces cytokine expression in purified human macrophages			
	IL-6 (pg/ml)	GM-CSF (pg/ml)	TNF- α (pg/ml)
Cells alone	0	0	0
CT DNA (50 μ g/ml)	0	0	0
EC DNA (50 μ g/ml)	2000	15.6	1000

Biological Role of IL-6 in Inducing Murine IgM Production in Response to CpG Motifs.

The kinetic studies described above revealed that induction of IL-6 secretion, which occurs within 1 hr post CpG stimulation, precedes IgM secretion. Since the optimal CpG motif for ODN inducing secretion of IL-6 is the same as that for IgM (Table 2), whether the CpG motifs independently induce IgM and IL-6 production or whether the IgM production is dependent on prior IL-6 secretion was examined. The addition of neutralizing anti-IL-6 antibodies inhibited *in vitro* IgM production mediated by CpG ODN in a dose-dependent manner but a control antibody did not (FIG. 4A). In contrast, anti-IL-6 addition did not affect either the basal level or the CpG-induced B cell proliferation (FIG. 4B).

Increased Transcriptional Activity of the IL-6 Promoter in Response to CpG DNA.

The increased level of IL-6 mRNA and protein after CpG DNA stimulation could result from transcriptional or post-transcriptional regulation. To determine if the transcriptional activity of the IL-6 promoter was upregulated in B cells cultured with CpG ODN, a murine B cell line, WEHI-231, which produces IL-6 in response to CpG DNA, was transfected with an IL-6 promoter-CAT construct (pIL-6/CAT) (Pottratz, S. T. et al., 17 β -estradiol) inhibits expression of human interleukin-6-promoter-reporter constructs by a receptor-dependent mechanism. *J. Clin. Invest.* 93:944). CAT assays were performed after stimulation with various concentrations of CpG or non-CpG ODN. As shown in FIG. 5, CpG ODN induced increased CAT activity in dose-dependent manner while non-CpG ODN failed to induce CAT activity. This confirms that CpG induces the transcriptional activity of the IL-6 promoter.

Dependence of B Cell Activation by CpG ODN on the Number of 5' and 3' Phosphorothioate Internucleotide Linkages.

To determine whether partial sulfur modification of the ODN backbone would be sufficient to enhance B cell activation, the effects of a series of ODN with the same sequence, but with differing numbers of S internucleotide linkages at the 5' and 3' ends were tested. Based on previous studies of nuclease degradation of ODN, it was determined that at least two phosphorothioate linkages at the 5' end of ODN were required to provide optimal protection of the ODN from degradation by intracellular exo- and endo-nucleases. Only chimeric ODN containing two 5' phosphorothioate-modified linkages, and a variable number of 3' modified linkages were therefore examined.

The lymphocyte stimulating effects of these ODN were tested at three concentrations (3.3, 10, and 30 μ M) by measuring the total levels of RNA synthesis (by 3 H uridine incorporation) or DNA synthesis (by 3 H thymidine incorporation) in treated spleen cell cultures (Example 10). O-ODN (0/0 phosphorothioate modifications) bearing a CpG motif caused no spleen cell stimulation unless added to the cultures at concentrations of at least 10 μ M (Example 10). However, when this sequence was modified with two S linkages at the 5' end and at least three S linkages at the 3' end, significant stimulation was seen at a dose of 3.3 μ M. At this low dose, the level of stimulation showed a progressive increase as the number of 3' modified bases was increased, until this reached or exceeded six, at which point the stimulation index began to decline. In general, the optimal number of 3' S linkages for spleen cell stimulation was five. At all three concentrations tested in these experiments, the S-ODN was less stimulatory than the optimal chimeric compounds.

Dependence of CpG-Mediated Lymphocyte Activation on the Type of Backbone Modification.

Phosphorothioate modified ODN (S-ODN) are far more nuclease resistant than phosphodiester modified ODN (O-ODN). Thus, the increased immune stimulation caused by S-ODN and S-O-ODN (i.e. chimeric phosphorothioate ODN in which the central linkages are phosphodiester, but the two 5' and five 3' linkages are phosphorothioate modified) compared to O-ODN may result from the nuclease resistance of the former. To determine the role of ODN nuclease resistance in immune stimulation by CpG ODN, the stimulatory effects of chimeric ODN in which the 5' and 3' ends were rendered nuclease resistant with either methylphosphonate (MP-), methylphosphorothioate (MPS-), phosphorothioate (S-), or phosphorodithioate (S₂-) internucleotide linkages were tested (Example 10). These studies showed that despite their nuclease resistance, MP-O-ODN were actually less immune stimulatory than O-ODN. However, combining the MP and S modifications by replacing both nonbridging O molecules with 5' and 3' MPS internucleotide linkages restored immune stimulation to a slightly higher level than that triggered by O-ODN.

S-O-ODN were far more stimulatory than O-ODN, and were even more stimulatory than S-ODN, at least at concentrations above 3.3 μ M. At concentrations below 3 μ M, the S-ODN with the 3M sequence was more potent than the corresponding S-O-ODN, while the S-ODN with the 3D sequence was less potent than the corresponding S-O-ODN (Example 10). In comparing the stimulatory CpG motifs of these two sequences, it was noted that the 3D sequence is a perfect match for the stimulatory motif in that the CpG is flanked by two 5' purines and two 3' pyrimidines. However, the bases immediately flanking the CpG in ODN 3D are not optimal; it has a 5' pyrimidine and a 3' purine. Based on further testing, it was found that the sequence requirement for immune stimulation is more stringent for S-ODN than for S-O- or O-ODN. S-ODN with poor matches to the optimal CpG motif cause little or no lymphocyte activation (e.g. Sequence 3D). However, S-ODN with good matches to the motif, most critically at the positions immediately flanking the CpG, are more potent than the corresponding S-O-ODN (e.g. Sequence 3M, Sequences 4 and 6), even though at higher concentrations (greater than 3 μ M) the peak effect from the S-O-ODN is greater (Example 10).

S₂-O-ODN were remarkably stimulatory, and caused substantially greater lymphocyte activation than the corresponding S-ODN or S-O-ODN at every tested concentration.

The increased B cell stimulation seen with CpG ODN bearing S or S₂ substitutions could result from any or all of the following effects: nuclease resistance, increased cellular uptake, increased protein binding, and altered intracellular localization. However, nuclease resistance can not be the only explanation, since the MP-O-ODN were actually less stimulatory than the O-ODN with CpG motifs. Prior studies have shown that ODN uptake by lymphocytes is markedly affected by the backbone chemistry (Zhao et al., (1993) Comparison of cellular binding and uptake of antisense phosphodiester, phosphorothioate, and mixed phosphorothioate and methylphosphonate oligonucleotides. (Antisense Research and Development 3, 53-66; Zhao et al., (1994) Stage specific oligonucleotide uptake in murine bone marrow B cell precursors. *Blood* 84, 3660-3666.) The highest cell membrane binding and uptake was seen with S-ODN, followed by S-O-ODN, O-ODN, and MP-ODN. This differential uptake correlates well with the degree of immune stimulation.

Unmethylated CpG Containing Oligos Have NK Cell Stimulatory Activity

Experiments were conducted to determine whether CpG containing oligonucleotides stimulated the activity of natural killer (NK) cells in addition to B cells. As shown in Table 8, a marked induction of NK activity among spleen cells cultured

dinucleotide and two 3' pyrimidines. Kinetic experiments demonstrated that NK activity peaked around 18 hrs. after addition of the ODN. The data indicates that the murine NK response is dependent on the prior activation of monocytes by CpG DNA, leading to the production of IL-12, TNF- α , and IFN- α/β (Example 11).

TABLE 9

Induction of NK Activity by DNA Containing CpG Motifs but not by Non-CpG DNA			
DNA or Cytokine Added		LU/10 ⁶	
		Mouse Cells	Human Cells
Expt. 1	None	0.00	0.00
	IL-2	16.68	15.82
	<i>E. coli</i> DNA	7.23	5.05
	Calf thymus DNA	0.00	0.00
Expt. 2	None	0.00	3.28
	1585 <u>gggGTCAACGTTG</u> agggggG (SEQ ID NO: 12)	7.38	17.98
	1629gtc..... (SEQ ID NO: 50)	0.00	4.4
Expt. 3	None	0.00	
	1613 <u>GCTAGACGTTAGTGT</u> (SEQ ID NO: 51)	5.22	
	1769z..... (SEQ ID NO: 52)	0.02	ND
	1619 <u>TCCATGTCGTTCCCTGATGCT</u> (SEQ ID NO: 43)	3.35	
	1765z..... (SEQ ID NO: 53)	0.11	

CpG dinucleotides in ODN sequences are indicated by underlying; Z indicates methylcytosine. Lower case letters indicate nuclease resistant phosphorothioate modified internucleotide linkages which, in titration experiments, were more than 20 times as potent as non-modified ODN, depending on the flanking bases. Poly G ends (g) were used in some ODN, because they significantly increase the level of ODN uptake.

with CpG ODN 1 and 3Dd was observed. In contrast, there was relatively no induction in effectors that had been treated with non-CpG control ODN.

TABLE 8

Induction Of NK Activity By CpG Oligodeoxynucleotides (ODN)				
ODN	% YAC-1 Specific Lysis*		% 2C11 Specific Lysis	
	50:1 Effector:Target	100:1	50:1 Effector:Target	100:1
None	-1.1	-1.4	15.3	16.6
1	16.1	24.5	38.7	47.2
3Dd	17.1	27.0	37.0	40.0
non-CpG ODN	-1.6	-1.7	14.8	15.4

Induction of NK Activity by DNA Containing CpG Motifs, but Not by Non-CpG DNA.

Bacterial DNA cultured for 18 hrs. at 37° C. and then assayed for killing of K562 (human) or Yac-1 (mouse) target cells induced NK lytic activity in both mouse spleen cells depleted of B cells and human PBMC, but vertebrate DNA did not (Table 9). To determine whether the stimulatory activity of bacterial DNA may be a consequence of its increased level of unmethylated CpG dinucleotides, the activating properties of more than 50 synthetic ODN containing unmethylated, methylated, or no CpG dinucleotides was tested. The results, summarized in Table 9, demonstrate that synthetic ODN can stimulate significant NK activity, as long as they contain at least one unmethylated CpG dinucleotide. No difference was observed in the stimulatory effects of ODN in which the CpG was within a palindrome (such as ODN 1585, which contains the palindrome AACGTT) from those ODN without palindromes (such as 1613 or 1619), with the caveat that optimal stimulation was generally seen with ODN in which the CpG was flanked by two 5' purines or a 5' GpT

35

From all of these studies, a more complete understanding of the immune effects of CpG DNA has been developed, which is summarized in FIG. 6.

Identification of B Cell and Monocyte/NK Cell-Specific Oligonucleotides

As shown in FIG. 6, CpG DNA can directly activate highly purified B cells and monocytic cells. There are many similarities in the mechanism through which CpG DNA activates these cell types. For example, both require NF κ B activation as explained further below.

In further studies of different immune effects of CpG DNA, it was found that there is more than one type of CpG motif. Specifically, oligo 1668, with the best mouse B cell motif, is a strong inducer of both B cell and natural killer (NK) cell activation, while oligo 1758 is a weak B cell activator, but still induces excellent NK responses (Table 10).

TABLE 10

Different CpG motifs stimulate optimal murine B cell and NK activation		
ODN Sequence	B cell activation ¹	NK activation ²
1668 <u>TCCATGACGTTCCCTGATGCT</u> (SEQ ID NO: 54)	42,849	2.52
1758 <u>TCTCCCAGCGTGC</u> CCCAT (SEQ ID NO: 55)	1,747	6.66
NONE	367	0.00

CpG dinucleotides are underlined; oligonucleotides were synthesized with phosphorothioate modified backbones to improve their nuclease resistance.

¹Measured by ³H thymidine incorporation after 48 hr culture with oligodeoxynucleotides at a 200 nM concentration as described in Example 1.

²Measured in lytic units.

65

Teleological Basis of Immunostimulatory, Nucleic Acids

Vertebrate DNA is highly methylated and CpG dinucleotides are underrepresented. However, the stimulatory CpG motif is common in microbial genomic DNA, but quite rare in vertebrate DNA. In addition, bacterial DNA has been reported to induce B cell proliferation and immunoglobulin (Ig) production, while mammalian DNA does not (Messina, J. P. et al., *J. Immunol.* 147:1759 (1991)). Experiments further described in Example 3, in which methylation of bacterial DNA with CpG methylase was found to abolish mitogenicity, demonstrates that the difference in CpG status is the cause of B cell stimulation by bacterial DNA. This data supports the following conclusion: that unmethylated CpG dinucleotides present within bacterial DNA are responsible for the stimulatory effects of bacterial DNA.

Teleologically, it appears likely that lymphocyte activation by the CpG motif represents an immune defense mechanism that can thereby distinguish bacterial from host DNA. Host DNA, which would commonly be present in many anatomic regions and areas of inflammation due to apoptosis (cell death), would generally induce little or no lymphocyte activation due to CpG suppression and methylation. However, the presence of bacterial DNA containing unmethylated CpG motifs can cause lymphocyte activation precisely in infected anatomic regions, where it is beneficial. This novel activation pathway provides a rapid alternative to T cell dependent antigen specific B cell activation. Since the CpG pathway synergizes with B cell activation through the antigen receptor, B cells bearing antigen receptor specific for bacterial antigens would receive one activation signal through cell membrane Ig and a second signal from bacterial DNA, and would therefore tend to be preferentially activated. The interrelationship of this pathway with other pathways of B cell activation provide a physiologic mechanism employing a polyclonal antigen to induce antigen-specific responses.

However, it is likely that B cell activation would not be totally nonspecific. B cells bearing antigen receptors specific for bacterial products could receive one activation signal through cell membrane Ig, and a second from bacterial DNA, thereby more vigorously triggering antigen specific immune responses. As with other immune defense mechanisms, the response to bacterial DNA could have undesirable consequences in some settings. For example, autoimmune responses to self antigens would also tend to be preferentially triggered by bacterial infections, since autoantigens could also provide a second activation signal to autoreactive B cells triggered by bacterial DNA. Indeed the induction of autoimmunity by bacterial infections is a common clinical observation. For example, the autoimmune disease systemic lupus erythematosus, which is: i) characterized by the production of anti-DNA antibodies; ii) induced by drugs which inhibit DNA methyltransferase (Cornacchia, E. J. et al., *J. Clin. Invest.* 92:38 (1993)); and iii) associated with reduced DNA methylation (Richardson, B. L. et al., *Arth. Rheum* 35:647 (1992)), is likely triggered at least in part by activation of DNA-specific B cells through stimulatory signals provided by CpG motifs, as well as by binding of bacterial DNA to antigen receptors.

Further, sepsis, which is characterized by high morbidity and mortality due to massive and nonspecific activation of the immune system may be initiated by bacterial DNA and other products released from dying bacteria that reach concentrations sufficient to directly activate many lymphocytes. Further evidence of the role of CpG DNA in the sepsis syndrome is described in Cowdery, J., et al., (1996) *The Journal of Immunology* 156:4570-4575.

Proposed Mechanisms of Action

Unlike antigens that trigger B cells through their surface Ig receptor, CpG-ODN did not induce any detectable Ca^{2+} flux, changes in protein tyrosine phosphorylation, or IP 3 generation. Flow cytometry with FITC-conjugated ODN with or without a CpG motif was performed as described in Zhao, Q et al., (*Antisense Research and Development* 3:53-66 (1993)), and showed equivalent membrane binding, cellular uptake, efflux, and intracellular localization. This suggests that there may not be cell membrane proteins specific for CpG ODN. Rather than acting through the cell membrane, that data suggests that unmethylated CpG containing oligonucleotides require cell uptake for activity: ODN covalently linked to a solid Teflon support were nonstimulatory, as were biotinylated ODN immobilized on either avidin beads or avidin coated petri dishes. CpG ODN conjugated to either FITC or biotin retained full mitogenic properties, indicating no steric hindrance.

Recent data indicate the involvement of the transcription factor NF κ B as a direct or indirect mediator of the CpG effect. For example, within 15 minutes of treating B cells or monocytes with CpG DNA, the level of NF κ B binding activity is increased (FIG. 7). However, it is not increased by DNA that does not contain CpG motifs. In addition, it was found that two different inhibitors of NF κ B activation, PDTC and gliotoxin, completely block the lymphocyte stimulation by CpG DNA as measured by B cell proliferation or monocytic cell cytokine secretion, suggesting that NF κ B activation is required for both cell types.

There are several possible mechanisms through which NF κ B can be activated. These include through activation of various protein kinases, or through the generation of reactive oxygen species. No evidence for protein kinase activation induced immediately after CpG DNA treatment of B cells or monocytic cells have been found, and inhibitors of protein kinase A, protein kinase C, and protein tyrosine kinases had no effects on the CpG induced activation. However, CpG DNA causes a rapid induction of the production of reactive oxygen species in both B cells and monocytic cells, as detected by the sensitive fluorescent dye dihydrorhodamine 123 as described in Royall, J. A., and Ischiropoulos, H. (*Archives of Biochemistry and Biophysics* 302:348-355 (1993)). Moreover, inhibitors of the generation of these reactive oxygen species completely block the induction of NF κ B and the later induction of cell proliferation and cytokine secretion by CpG DNA.

Working backwards, the next question was how CpG DNA leads to the generation of reactive oxygen species so quickly. Previous studies by the inventors demonstrated that oligonucleotides and plasmid or bacterial DNA are taken up by cells into endosomes. These endosomes rapidly become acidified inside the cell. To determine whether this acidification step may be important in the mechanism through which CpG DNA activates reactive oxygen species, the acidification step was blocked with specific inhibitors of endosome acidification including chloroquine, monensin, and bafilomycin, which work through different mechanisms. FIG. 8A shows the results from a flow cytometry study using mouse B cells with the dihydrorhodamine 123 dye to determine levels of reactive oxygen species. The dye only sample in Panel A of the figure shows the background level of cells positive for the dye at 28.6%. As expected, this level of reactive oxygen species was greatly increased to 80% in the cells treated for 20 minutes with PMA and ionomycin, a positive control (Panel B). The cells treated with the CpG oligo also showed an increase in the level of reactive oxygen species such that more than 50% of the cells became positive (Panel D). However,

cells treated with an oligonucleotide with the identical sequence except that the CpG was switched did not show this significant increase in the level of reactive oxygen species (Panel E).

In the presence of chloroquine, the results are very different (FIG. 8B). Chloroquine slightly lowers the background level of reactive oxygen species in the cells such that the untreated cells in Panel A have only 4.3% that are positive. Chloroquine completely abolishes the induction of reactive oxygen species in the cells treated with CpG DNA (Panel B) but does not reduce the level of reactive oxygen species in the cells treated with PMA and ionomycin (Panel E). This demonstrates that unlike the PMA plus ionomycin, the generation of reactive oxygen species following treatment of B cells with CpG DNA requires that the DNA undergo an acidification step in the endosomes. This is a completely novel mechanism of leukocyte activation. Chloroquine, monensin, and bafilomycin also appear to block the activation of NFκB by CpG DNA as well as the subsequent proliferation and induction of cytokine secretion.

Presumably, there is a protein in or near the endosomes that specifically recognizes DNA containing CpG motifs and leads to the generation of reactive oxygen species. To detect any protein in the cell cytoplasm that may specifically bind CpG DNA, we used electrophoretic mobility shift assays (EMSA) with 5' radioactively labeled oligonucleotides with or without CpG motifs. A band was found that appears to represent a protein binding specifically to single stranded oligonucleotides that have CpG motifs, but not to oligonucleotides that lack CpG motifs or to oligonucleotides in which the CpG motif has been methylated. This binding activity is blocked if excess of oligonucleotides that contain the NFκB binding site was added. This suggests that an NFκB or related protein is a component of a protein or protein complex that binds the stimulatory CpG oligonucleotides.

No activation of CREB/ATF proteins was found at time points where NFκB was strongly activated. These data therefore do not provide proof that NFκB proteins actually bind to the CpG nucleic acids, but rather that the proteins are required in some way for the CpG activity. It is possible that a CREB/ATF or related protein may interact in some way with NFκB proteins or other proteins thus explaining the remarkable similarity in the binding motifs for CREB proteins and the optimal CpG motif. It remains possible that the oligos bind to a CREB/ATF or related protein, and that this leads to NFκB activation.

Alternatively, it is very possible that the CpG nucleic acids may bind to one of the TRAF proteins that bind to the cytoplasmic region of CD40 and mediate NFκB activation when CD40 is cross-linked. Examples of such TRAF proteins include TRAF-2 and TRAF-5.

Method for Making Immunostimulatory Nucleic Acids

For use in the instant invention, nucleic acids can be synthesized de novo using any of a number of procedures well known in the art. For example, the β-cyanoethyl phosphoramidite method (S. L. Beaucage and M. H. Caruthers, (1981) *Tet. Let.* 22:1859); nucleoside H-phosphonate method (Garegg et al., (1986) *Tet. Let.* 27: 4051-4054; Froehler et al., (1986) *Nucl. Acid. Res.* 14: 5399-5407; Garegg et al., (1986) *Tet. Let.* 27: 4055-4058, Gaffney et al., (1988) *Tet. Let.* 29:2619-2622). These chemistries can be performed by a variety of automated oligonucleotide synthesizers available in the market. Alternatively, oligonucleotides can be prepared from existing nucleic acid sequences (e.g. genomic or cDNA) using known techniques, such as those employing restriction enzymes, exonucleases or endonucleases.

For use in vivo, nucleic acids are preferably relatively resistant to degradation (e.g. via endo- and exo-nucleases). Secondary structures, such as stem loops, can stabilize nucleic acids against degradation. Alternatively, nucleic acid stabilization can be accomplished via phosphate backbone modifications. A preferred stabilized nucleic acid has at least a partial phosphorothioate modified backbone. Phosphorothioates may be synthesized using automated techniques employing either phosphoramidate or H-phosphonate chemistries. Aryl- and alkyl-phosphonates can be made e.g. as described in U.S. Pat. No. 4,469,863; and alkylphosphotriesters (in which the charged oxygen moiety is alkylated as described in U.S. Pat. No. 5,023,243 and European Patent No. 092,574) can be prepared by automated solid phase synthesis using commercially available reagents. Methods for making other DNA backbone modifications and substitutions have been described (Uhlmann, E. and Peyman, A. (1990) *Chem. Rev.* 90:544; Goodchild, J. (1990) *Bioconjugate Chem.* 1:165). 2'-O-methyl nucleic acids with CpG motifs also cause immune activation, as do ethoxy-modified CpG nucleic acids. In fact, no backbone modifications have been found that completely abolish the CpG effect, although it is greatly reduced by replacing the C with a 5-methyl C.

For administration in vivo, nucleic acids may be associated with a molecule that results in higher affinity binding to target cell (e.g. B-cell, monocytic cell and natural killer (NK) cell) surfaces and/or increased cellular uptake by target cells to form a "nucleic acid delivery complex". Nucleic acids can be ionically, or covalently associated with appropriate molecules using techniques which are well known in the art. A variety of coupling or crosslinking agents can be used e.g. protein A, carbodiimide, and N-succinimidyl-3-(2-pyridyldithio) propionate (SPDP). Nucleic acids can alternatively be encapsulated in liposomes or virosomes using well-known techniques.

Therapeutic Uses of Immunostimulatory Nucleic Acid Molecules

Based on their immunostimulatory properties, nucleic acid molecules containing at least one unmethylated CpG dinucleotide can be administered to a subject in vivo to treat an "immune system deficiency". Alternatively, nucleic acid molecules containing at least one unmethylated CpG dinucleotide can be contacted with lymphocytes (e.g. B cells, monocytic cells or NK cells) obtained from a subject having an immune system deficiency ex vivo and activated lymphocytes can then be reimplanted in the subject.

As reported herein, in response to unmethylated CpG containing nucleic acid molecules, an increased number of spleen cells secrete IL-6, IL-12, IFN-α, IFN-β, IL-1, IL-3, IL-10, TNF-α, TNF-β, GM-CSF, RANTES, and probably others. The increased IL-6 expression was found to occur in B cells, CD4⁺ T cells and monocytic cells.

Immunostimulatory nucleic acid molecules can also be administered to a subject in conjunction with a vaccine to boost a subject's immune system and thereby effect a better response from the vaccine. Preferably the immunostimulatory nucleic acid molecule is administered slightly before or at the same time as the vaccine. A conventional adjuvant may optionally be administered in conjunction with the vaccine, which is minimally comprised of an antigen, as the conventional adjuvant may further improve the vaccination by enhancing antigen absorption.

When the vaccine is a DNA vaccine at least two components determine its efficacy. First, the antigen encoded by the vaccine determines the specificity of the immune response. Second, if the backbone of the plasmid contains CpG motifs, it functions as an adjuvant for the vaccine. Thus, CpG DNA

acts as an effective “danger signal” and causes the immune system to respond vigorously to new antigens in the area. This mode of action presumably results primarily from the stimulatory local effects of CpG DNA on dendritic cells and other “professional” antigen presenting cells, as well as from the costimulatory effects on B cells.

Immunostimulatory oligonucleotides and unmethylated CpG containing vaccines, which directly activate lymphocytes and co-stimulate an antigen-specific response, are fundamentally different from conventional adjuvants (e.g. aluminum precipitates), which are inert when injected alone and are thought to work through absorbing the antigen and thereby presenting it more effectively to immune cells. Further, conventional adjuvants only work for certain antigens, only induce an antibody (humoral) immune response (Th2), and are very poor at inducing cellular immune responses (Th1). For many pathogens, the humoral response contributes little to protection, and can even be detrimental.

In addition, an immunostimulatory oligonucleotide can be administered prior to, along with or after administration of a chemotherapy or immunotherapy to increase the responsiveness of the malignant cells to subsequent chemotherapy or immunotherapy or to speed the recovery of the bone marrow through induction of restorative cytokines such as GM-CSF. CpG nucleic acids also increase natural killer cell lytic activity and antibody dependent cellular cytotoxicity (ADCC). Induction of NK activity and ADCC may likewise be beneficial in cancer immunotherapy, alone or in conjunction with other treatments.

Another use of the described immunostimulatory nucleic acid molecules is in desensitization therapy for allergies, which are generally caused by IgE antibody generation against harmless allergens. The cytokines that are induced by unmethylated CpG nucleic acids are predominantly of a class called “Th1” which is most marked by a cellular immune response and is associated with IL-12 and IFN- γ . The other major type of immune response is termed a Th2 immune response, which is associated with more of an antibody immune response and with the production of IL-4, IL-5 and IL-10. In general, it appears that allergic diseases are mediated by Th2 type immune responses and autoimmune diseases by Th1 immune response. Based on the ability of the immunostimulatory nucleic acid molecules to shift the immune response in a subject from a Th2 (which is associated with production of IgE antibodies and allergy) to a Th1 response (which is protective against allergic reactions), an effective dose of an immunostimulatory nucleic acid (or a vector containing a nucleic acid) alone or in conjunction with an allergen can be administered to a subject to treat or prevent an allergy.

Nucleic acids containing unmethylated CpG motifs may also have significant therapeutic utility in the treatment of asthma. Th2 cytokines, especially IL-4 and IL-5 are elevated in the airways of asthmatic subjects. These cytokines promote important aspects of the asthmatic inflammatory response, including IgE isotype switching, eosinophil chemotaxis and activation and mast cell growth. Th1 cytokines, especially IFN- γ and IL-12, can suppress the formation of Th2 clones and production of Th2 cytokines.

As described in detail in the following Example 12, oligonucleotides containing an unmethylated CpG motif (i.e. TCCATGACGTTCTGACGTT; SEQ ID NO: 10), but not a control oligonucleotide (TCCATGAGCTTCTGAGTCT; SEQ ID NO: 11) prevented the development of an inflammatory cellular infiltrate and eosinophilia in a murine model of asthma. Furthermore, the suppression of eosinophilic inflam-

mation was associated with a suppression of a Th2 response and induction of a Th1 response.

For use in therapy, an effective amount of an appropriate immunostimulatory nucleic acid molecule alone or formulated as a delivery complex can be administered to a subject by any mode allowing the oligonucleotide to be taken up by the appropriate target cells (e.g., B-cells and monocytic cells). Preferred routes of administration include oral and transdermal (e.g., via a patch). Examples of other routes of administration include injection (subcutaneous, intravenous, parenteral, intraperitoneal, intrathecal, etc.). The injection can be in a bolus or a continuous infusion.

A nucleic acid alone or as a nucleic acid delivery complex can be administered in conjunction with a pharmaceutically acceptable carrier. As used herein, the phrase “pharmaceutically acceptable carrier” is intended to include substances that can be coadministered with a nucleic acid or a nucleic acid delivery complex and allows the nucleic acid to perform its indicated function. Examples of such carriers include solutions, solvents, dispersion media, delay agents, emulsions and the like. The use of such media for pharmaceutically active substances are well known in the art. Any other conventional carrier suitable for use with the nucleic acids falls within the scope of the instant invention.

The language “effective amount” of a nucleic acid molecule refers to the amount necessary or sufficient to realize a desired biologic effect. For example, an effective amount of a nucleic acid containing at least one unmethylated CpG for treating an immune system deficiency could be that amount necessary to eliminate a tumor, cancer, or bacterial, viral or fungal infection. An effective amount for use as a vaccine adjuvant could be that amount useful for boosting a subjects immune response to a vaccine. An “effective amount” for treating asthma can be that amount useful for redirecting a Th2 type of immune response that is associated with asthma to a Th1 type of response. The effective amount for any particular application can vary depending on such factors as the disease or condition being treated, the particular nucleic acid being administered (e.g. the number of unmethylated CpG motifs or their location in the nucleic acid), the size of the subject, or the severity of the disease or condition. One of ordinary skill in the art can empirically determine the effective amount of a particular oligonucleotide without necessitating undue experimentation.

The present invention is further illustrated by the following Examples which in no way should be construed as further limiting. The entire contents of all of the references (including literature references, issued patents, published patent applications, and co-pending patent applications) cited throughout this application are hereby expressly incorporated by reference.

EXAMPLES

Example 1

Effects of ODNs on B Cell Total RNA Synthesis and Cell Cycle

B cells were purified from spleens obtained from 6-12 wk old specific pathogen free DBA/2 or BXSJ mice (bred in the University of Iowa animal care facility; no substantial strain differences were noted) that were depleted of T cells with anti-Thy-1.2 and complement and centrifugation over lympholyte M (CEDARLANE® Laboratories Ltd., Hornby, Ontario, Canada) (“B cells”). B cells contained fewer than 1% CD4⁺ or CD8⁺ cells. 8×10⁴ B cells were dispensed in tripli-

35

cate into 96 well microtiter plates in 100 μ l RPMI containing 10% FBS (heat inactivated to 65° C. for 30 min.), 50 μ M 2-mercaptoethanol, 100 U/ml penicillin, 100 μ g/ml streptomycin, and 2 mM L-glutamate. 20 μ M ODN were added at the start of culture for 20 h at 37° C., cells pulsed with 1 μ Ci of 3 H uridine, and harvested and counted 4 hr later. Ig secreting B cells were enumerated using the ELISA spot assay after culture of whole spleen cells with ODN at 20 μ M for 48 hr. Data, reported in Table 1, represent the stimulation index compared to cells cultured without ODN. 3 H thymidine incorporation assays showed similar results, but with some nonspecific inhibition by thymidine released from degraded ODN (Matsun. S and A. M. Krieg (1992) Nonspecific suppression of 3 H-thymidine incorporation by control oligonucleotides. *Antisense Research and Development* 2:325).

Example 2

Effects of ODN on Production of IgM from B Cells

Single cell suspensions from the spleens of freshly killed mice were treated with anti-Thy1, anti-CD4, and anti-CD8 and complement by the method of Leibson et al., *J. Exp. Med.* 154:1681 (1981). Resting B cells (<02% T cell contamination) were isolated from the 63-70% band of a discontinuous Percoll gradient by the procedure of DeFranco et al., *J. Exp. Med.* 155:1523 (1982). These were cultured as described above in 30 μ M ODN or 20 μ g/ml LPS for 48 hr. The number of B cells actively secreting IgM was maximal at this time point, as determined by ELISpot assay (Klinman, D. M. et al. *J. Immunol.* 144:506 (1990)). In that assay, B cells were incubated for 6 hrs on anti-Ig coated microtiter plates. The Ig they produced (>99% IgM) was detected using phosphatase-labelled anti-Ig (SouthernBiotech, Birmingham, Ala.). The antibodies produced by individual B cells were visualized by addition of BCIP (SIGMA-ALDRICH™, St. Louis Mo.) which forms an insoluble blue precipitate in the presence of phosphatase. The dilution of cells producing 20-40 spots/well was used to determine the total number of antibody-secreting B cells/sample. All assays were performed in triplicate (data reported in Table 1). In some experiments, culture supernatants were assayed for IgM by ELISA, and showed similar increases in response to CpG-ODN.

Example 3

B Cell Stimulation by Bacterial DNA

DBA/2 B cells were cultured with no DNA or 50 μ g/ml of a) *Micrococcus lysodeikticus*; b) NZB/N mouse spleen; and c) NFS/N mouse spleen genomic DNAs for 48 hours, then pulsed with 3 H thymidine for 4 hours prior to cell harvest. Duplicate DNA samples were digested with DNase I for 30 minutes at 37 C prior to addition to cell cultures. *E. coli* DNA also induced an 8.8 fold increase in the number of IgM secreting B cells by 48 hours using the ELISA-spot assay.

DBA/2 B cells were cultured with either no additive, 50 μ g/ml LPS or the ODN 1; 1a; 4; or 4a at 20 μ M. Cells were cultured and harvested at 4, 8, 24 and 48 hours. BXSb cells were cultured as in Example 1 with 5, 10, 20, 40 or 80 μ M of ODN 1; 1a; 4; or 4a or LPS. In this experiment, wells with no ODN had 3833 cpm. Each experiment was performed at least three times with similar results. Standard deviations of the triplicate wells were <5%.

36

Example 4

Effects of ODN on Natural Killer (NK) Activity

10 \times 10⁶ C57BL/6 spleen cells were cultured in two ml RPMI (supplemented as described for Example 1) with or without 40 μ M CpG or non-CpG ODN for forty-eight hours. Cells were washed, and then used as effector cells in a short term 51 Cr release assay with YAC-1 and 2C11, two NK sensitive target cell lines (Ballas, Z. K. et al. (1993) *J. Immunol.* 150:17). Effector cells were added at various concentrations to 10⁴ 51 Cr-labeled target cells in V-bottom microtiter plates in 0.2 ml, and incubated in 5% CO₂ for 4 hr. at 37° C. Plates were then centrifuged, and an aliquot of the supernatant counted for radioactivity. Percent specific lysis was determined by calculating the ratio of the 51 Cr released in the presence of effector cells minus the 51 Cr released when the target cells are cultured alone, over the total counts released after cell lysis in 2% acetic acid minus the 51 Cr cpm released when the cells are cultured alone.

Example 5

In Vivo Studies with CpG Phosphorothioate ODN

Mice were weighed and injected IP with 0.25 ml of sterile PBS or the indicated phosphorothioate ODN dissolved in PBS. Twenty four hours later, spleen cells were harvested, washed, and stained for flow cytometry using phycoerythrin conjugated 6B2 to gate on B cells in conjunction with biotin conjugated anti Ly-6A/E or anti-Ia^d (Pharmingen, San Diego, Calif.) or anti-Bla-1 (Hardy, R. R. et al., *J. Exp. Med.* 159: 1169 (1984). Two mice were studied for each condition and analyzed individually.

Example 6

Titration of Phosphorothioate ODN for B Cell Stimulation

B cells were cultured with phosphorothioate ODN with the sequence of control ODN 1a or the CpG ODN 1d and 3Db and then either pulsed after 20 hr with 3 H uridine or after 44 hr with 3 H thymidine before harvesting and determining cpm.

Example 7

Rescue of B Cells from Apoptosis

WEHI-231 cells (5 \times 10⁴/well) were cultured for 1 hr. at 37 C. in the presence or absence of LPS or the control ODN 1a or the CpG ODN 1d and 3Db before addition of anti-IgM (1 μ /ml). Cells were cultured for a further 20 hr. before a 4 hr. pulse with 2 μ Ci/well 3 H thymidine. In this experiment, cells with no ODN or anti-IgM gave 90.4 \times 10³ cpm of 3 H thymidine incorporation by addition of anti-IgM. The phosphodiester ODN shown in Table 1 gave similar protection, though with some nonspecific suppression due to ODN degradation. Each experiment was repeated at least 3 times with similar results.

Example 8

In Vivo Induction of Murine IL-6

DBA/2 female mice (2 mos. old) were injected IP with 500 μ g CpG or control phosphorothioate ODN. At various

time points after injection, the mice were bled. Two mice were studied for each time point. IL-6 was measured by Elisa, and IL-6 concentration was calculated by comparison to a standard curve generated using recombinant IL-6. The sensitivity of the assay was 10 pg/ml. Levels were undetectable after 8 hr.

Example 9

Systemic Induction of Murine IL-6 Transcription

Mice and cell lines. DBA/2, BALB/c, and C3H/HeJ mice at 5-10 wk of age were used as a source of lymphocytes. All mice were obtained from The Jackson Laboratory (Bar Harbor, Me.), and bred and maintained under specific pathogen-free conditions in the University of Iowa Animal Care Unit. The mouse B cell line CH12.LX was kindly provided by Dr. G. Bishop (University of Iowa, Iowa City).

Cell preparation. Mice were killed by cervical dislocation. Single cell suspensions were prepared aseptically from the spleens from mice. T cell depleted mouse splenocytes were prepared by using anti-Thy-1.2 and complement and centrifugation over lympholyte M (CEDARLANE® Laboratories Ltd., Hornby, Ontario, Canada) as described (Krieg, A. M. et al., (1989) A role for endogenous retroviral sequences in the regulation of lymphocyte activation. *J. Immunol.* 143: 2448).

ODN and DNA. Phosphodiester oligonucleotides (O-ODN) and the backbone modified phosphorothioate oligonucleotides (S-ODN) were obtained from the DNA Core facility at the University of Iowa or from OPERON® Technologies (Alameda, Calif.). *E. coli* DNA (Strain B) and calf thymus DNA were purchased from SIGMA-ALDRICH™ (St. Louis, Mo.). All DNA and ODN were purified by extraction with phenol:chloroform:isoamyl alcohol (25:24:1) and/or ethanol precipitation. *E. coli* and calf thymus DNA were single stranded prior to use by boiling for 10 min. followed by cooling on ice for 5 min. For some experiments, *E. coli* and calf thymus DNA were digested with DNase I (2 U/μg of DNA) at 37° C. for 2 hr in 1×SSC with 5 mM MgCl₂. To methylate the cytosine in CpG dinucleotides in *E. coli* DNA, *E. coli* DNA was treated with CpG methylase (M. SssI; 2 U/μg of DNA) in NEBuffer 2 supplemented with 160 μM S-adenosyl methionine and incubated overnight at 37° C. Methylated DNA was purified as above. Efficiency of methylation was confirmed by Hpa II digestion followed by analysis by gel electrophoresis. All enzymes were purchased from NEW ENGLAND BIOLABS® Inc. (Beverly, Mass.). LPS level in ODN was less than 12.5 ng/mg and *E. coli* and calf thymus DNA contained less than 2.5 ng of LPS/mg of DNA by *Limulus* assay.

Cell Culture. All cells were cultured at 37° C. in a 5% CO₂ humidified incubator maintained in RPMI-1640 supplemented with 10% (v/v) heat inactivated fetal calf serum (FCS), 1.5 mM L-glutamine, 50 μg/ml, CpG or non-CpG phosphodiester ODN (O-ODN) (20 μM), phosphorothioate ODN (S-ODN) (0.5 μM), or *E. coli* or calf thymus DNA (50 μg/ml) at 37° C. for 24 hr. (for IL-6 production) or 5 days (for IgM production). Concentrations of stimulants were chosen based on preliminary studies with titrations. In some cases, cells were treated with CpG O-ODN along with various concentrations (1-10 μg/ml) of neutralizing rat IgG1 antibody against murine IL-6 (hybridoma MP5-20F3) or control rat IgG1 mAb to *E. coli* β-galactosidase (hybridoma GL113; ATCC, Rockville, Md.) (20) for 5 days. At the end of incubation, culture supernatant fractions were analyzed by ELISA as below.

In vivo induction of IL-6 and IgM. BALB/c mice were injected intravenously (iv) with PBS, calf thymus DNA (200 μg/100 μl PBS/mouse), *E. coli* DNA (200 μg/100 μl PBS/mouse), or CpG or non-CpG S-ODN (200 μg/100 μl PBS/mouse). Mice (two/group) were bled by retroorbital puncture and sacrificed by cervical dislocation at various time points. Liver, spleen, thymus, and bone marrow were removed and RNA was prepared from those organs using RNeasy RBB (Tel-Test, Friendswood, Tex.) according to the manufacturers protocol.

ELISA. Flat-bottomed Immun 1 plates (Dynatech Laboratories, Inc., Chantilly, Va.) were coated with 100 μl/well of anti-mouse IL-6 mAb (MP5-20F3) (2 μg/ml) or anti-mouse IgM μ-chain specific (5 μg/ml; SIGMA-ALDRICH™, St. Louis, Mo.) in carbonate-bicarbonate, pH 9.6 buffer (15 nM Na₂CO₃, 35 mM NaHCO₃) overnight at 4° C. The plates were then washed with TPBS (0.5 mM MgCl₂·6H₂O, 2.68 mM KCl, 1.47 mM KH₂PO₄, 0.14 M NaCl, 6.6 mM K₂HPO₄, 0.5% Tween 20) and blocked with 10% FCS in TPBS for 2 hr at room temperature and then washed again. Culture supernatants, mouse sera, recombinant mouse IL-6 (PHARMINGEN®, San Diego, Calif.) or purified mouse IgM (CALBIOCHEM®, San Diego, Calif.) were appropriately diluted in 10% FCS and incubated in triplicate wells for 6 hr at room temperature. The plates were washed and 100 μl/well of biotinylated rat anti-mouse IL-6 monoclonal antibodies (MP5-32C11, PHARMINGEN®, San Diego, Calif.) (1 μg/ml in 10% FCS) or biotinylated anti-mouse Ig (SIGMA-ALDRICH™, St. Louis, Mo.) were added and incubated for 45 min. at room temperature following washes with TPBS. Horseradish peroxidase (HRP) conjugated avidin (BIORAD® Laboratories, Hercules, Calif.) at 1:4000 dilution in 10% FCS (100 μl/well) was added and incubated at room temperature for 30 min. The plates were washed and developed with o-phenyldiamine dihydrochloride (OPD; SIGMA-ALDRICH™, St. Louis Mo.) 0.05 M phosphate-citrate buffer, pH 5.0, for 30 min. The reaction was stopped with 0.67 N H₂SO₄ and plates were read on a microplate reader (Cambridge Technology, Inc., Watertown, Mass.) at 490-600 nm. The results are shown in FIGS. 1 and 2.

RT-PCR. A sense primer, an antisense primer, and an internal oligonucleotide probe for IL-6 were synthesized using published sequences (Montgomery, R. A. and M. S. Dallman (1991), Analysis of cytokine gene expression during fetal thymic ontogeny using the polymerase chain reaction (*J. Immunol.*) 147:554). cDNA synthesis and IL-6 PCR was done essentially as described by Montgomery and Dallman (Montgomery, R. A. and M. S. Dallman (1991), Analysis of cytokine gene expression during fetal thymic ontogeny using the polymerase chain reaction (*J. Immunol.*) 147:554) using RT-PCR reagents from PERKINELMER® (Hayward, Calif.). Samples were analyzed after 30 cycles of amplification by gel electrophoresis followed by unblot analysis (Stoye, J. P. et al., (1991) DNA hybridization in dried gels with fragmented probes: an improvement over blotting techniques, *Techniques* 3:123). Briefly, the gel was hybridized at room temperature for 30 min. in denaturation buffer (0.05 M NaOH, 1.5M NaCl) followed by incubation for 30 min. in renaturation buffer (1.5 M NaCl, 1 M Tris, pH 8) and a 30 min. wash in double distilled water. The gel was dried and prehybridized at 47° C. for 2 hr. hybridization buffer (5×SSPE, 0.1% SDS) containing 10 μg/ml denatured salmon sperm DNA. The gel was hybridized with 2×10⁶ cpm/ml γ-³²P]ATP end-labeled internal oligonucleotide probe for IL-6 (5'CATTTCCACGATTTCCA3') SEQ ID NO:56) over-

night at 47° C., washed 4 times (2×SSC, 0.2% SDS) at room temperature and autoradiographed. The results are shown in FIG. 3.

Cell Proliferation assay. DBA/2 mice spleen B cells (5×10⁴ cells/100 μl/well) were treated with media, CpG or non-CpG S-ODN (0.5 μM) or O-ODN (20 μM) for 24 hr at 37° C. Cells were pulsed for the last four hr. with either [³H] Thymidine or [³H] Uridine (1 μCi/well). Amounts of [³H] incorporated were measured using Liquid Scintillation Analyzer (Packard Instrument Co., Downers Grove, Ill.).

Transfections and CAT assays. WEHI-231 cells (10⁷ cells) were electroporated with 20 μg of control or human IL-6 promoter-CAT construct (kindly provided by S. Manolagas, Univ. of Arkansas) (Pottratz, S. T. et al., (1994) 17B-estradiol inhibits expression of human interleukin-6 promoter-reporter constructs by a receptor-dependent mechanism. *J. Clin. Invest.* 93:944) at 250 mV and 960 μF. Cells were stimulated with various concentrations of CpG or non-CpG ODN after electroporation. Chloramphenicol acetyltransferase (CAT) activity was measured by a solution assay (Seed, B. and J. Y. Sheen (1988) A single phase-extraction assay for chloramphenicol acetyl transferase activity. *Gene* 76:271) 16 hr. after transfection. The results are presented in FIG. 5.

Example 10

Oligodeoxynucleotide Modifications Determine the Magnitude of B Cell Stimulation by CpG Motifs

ODN were synthesized on an APPLIED BIOSYSTEMS® Inc. (Foster City, Calif.) model 380A, 380B, or 394 DNA synthesizer using standard procedures (Beacage and Caruthers (1981) Deoxynucleoside phosphoramidites—A new class of key intermediates for deoxypolynucleotide synthesis. *Tetrahedron Letters* 22, 1859-1862.). Phosphodiester ODN were synthesized using standard beta-cyanoethyl phosphoramidite chemistry. Phosphorothioate linkages were introduced by oxidizing the phosphite linkage with elemental sulfur instead of the standard iodine oxidation. The four common nucleoside phosphoramidites were purchased from Applied Biosystems. All phosphodiester and thioate containing ODN were deprotected by treatment with concentrated ammonia at 55° C. for 12 hours. The ODN were purified by gel exclusion chromatography and lyophilized to dryness prior to use. Phosphorodithioate linkages were introduced by using deoxynucleoside S-(b-benzoylmercaptoethyl)pyrrolidino thiophosphoramidites (Wiesler, W. T. et al., (1993) In *Methods in Molecular Biology: Protocols for Oligonucleotides and Analogs—Synthesis and Properties*, Agrawal, S. (ed.), Humana Press, 191-206.). Dithioate containing ODN were deprotected by treatment with concentrated ammonia at 55° C. for 12 hours followed by reverse phase HPLC purification.

In order to synthesize oligomers containing methylphosphonothioates or methylphosphonates as well as phosphodiester at any desired internucleotide linkage, two different synthetic cycles were used. The major synthetic differences in the two cycles are the coupling reagent where dialkylaminomethylnucleoside phosphines are used and the oxidation reagents in the case of methylphosphonothioates. In order to synthesize either derivative, the condensation time has been increased for the dialkylaminomethylnucleoside phosphines due to the slower kinetics of coupling (Jager and Engels, (1984) Synthesis of deoxynucleoside methylphosphonates via a phosphoramidite approach. *Tetrahedron Letters* 24, 1437-1440). After the coupling step has been completed, the methylphosphinodiester is treated with the sulfurizing reagent (5% elemental sulfur, 100 millimolar

N,N-dimethylaminopyridine in carbon disulfide/pyridine/triethylamine), four consecutive times for 450 seconds each to produce methylphosphonothioates. To produce methylphosphonate linkages, the methylphosphinodiester is treated with standard oxidizing reagent (0.1 M iodine in tetrahydrofuran/2,6-lutidine/water).

The silica gel bound oligomer was treated with distilled pyridine/concentrated ammonia, 1:1, (v/v) for four days at 4 degrees centigrade. The supernatant was dried in vacuo, dissolved in water and chromatographed on a G50/50 Sephadex column.

As used herein, O-ODN refers to ODN which are phosphodiester; S-ODN are completely phosphorothioate modified; S-O-ODN are chimeric ODN in which the central linkages are phosphodiester, but the two 5' and five 3' linkages are phosphorothioate modified; S₂-O-ODN are chimeric ODN in which the central linkages are phosphodiester, but the two 5' and five 3' linkages are phosphorodithioate modified; and MP-O-ODN are chimeric ODN in which the central linkages are phosphodiester, but the two 5' and five 3' linkages are methylphosphonate modified. The ODN sequences studied (with CpG dinucleotides indicated by underlining) include:

3D (5' GAGAACGCTGGACCTTCCAT) ; ; (SEQ ID NO: 23)
 3M (5' TCCATGTCCGGTCTGATGCT) ; ; (SEQ ID NO: 31)
 5 (5' GGCGTTATTCTGACTCGCC) ; ; (SEQ ID NO: 57)
 and
 6 (5' CCTACCGTTGTATGCGCCCAGCT) ; ; (SEQ ID NO: 58)

These sequences are representative of literally hundreds of CpG and non-CpG ODN that have been tested in the course of these studies.

Mice. DBA/2, or BXSB mice obtained from The Jackson Laboratory (Bar Harbor, Me.), and maintained under specific pathogen-free conditions were used as a source of lymphocytes at 5-10 wk of age with essentially identical results.

Cell proliferation assay. For cell proliferation assays, mouse spleen cells (5×10⁴ cells/100 μl/well) were cultured at 37° C. in a 5% CO₂ humidified incubator in RPMI-1640 supplemented with 10% (v/v) heat inactivated fetal calf serum (heated to 65° C. for experiments with O-ODN, or 56° C. for experiments using only modified ODN), 1.5 μM L-glutamine, 50 μM 2-mercaptoethanol, 100 U/ml penicillin and 100 μg/ml streptomycin for 24 hr or 48 hr as indicated. 1 μCi of [³H] uridine or thymidine (as indicated) was added to each well, and the cells harvested after an additional 4 hours of culture. Filters were counted by scintillation counting. Standard deviations of the triplicate wells were <5%. The results are presented in FIGS. 6-8.

Example 11

Induction of NK Activity

Phosphodiester ODN were purchased from OPERON® Technologies (Alameda, Calif.). Phosphorothioate ODN were purchased from the DNA core facility, University of Iowa, or from The Midland Certified Reagent Company (Midland Tex.). *E. coli* (strain B) DNA and calf thymus DNA were purchased from Sigma, SIGMA-ALDRICH™, (St. Louis, Mo.). All DNA and ODN were purified by extraction with phenol:chloroform:isoamyl alcohol (25:24:1) and/or ethanol precipitation. The LPS level in ODN was less than 12.5 ng/mg and *E. coli* and calf thymus DNA contained less than 2.5 ng of LPS/mg of DNA by *Limulus* assay.

Virus-free, 4-6 week old, DBA/2, C57BL/6 (B6) and congenitally athymic BALB/C mice were obtained on contract through the Veterans Affairs from the National Cancer Institute (Bethesda, Md.). C57BL/6 SCID mice were bred in the SPF barrier facility at the University of Iowa Animal Care Unit.

Human peripheral mononuclear blood leukocytes (PBMC) were obtained as previously described (Ballas, Z. K. et al., (1990) *J. Allergy Clin. Immunol.* 85:453; Ballas, Z. K. and W. Rasmussen (1990) *J. Immunol.* 145:1039; Ballas, Z. K. and W. Rasmussen (1993) *J. Immunol.* 150: 17). Human or murine cells were cultured at 5×10^6 /well, at 37° C. in a 5% CO₂ humidified atmosphere in 24-well plates (Ballas, Z. K. et al., (1990) *J. Allergy Clin. Immunol.* 85:453; Ballas, Z. K. and W. Rasmussen (1990) *J. Immunol.* 145:1039; and Ballas, Z. K. and W. Rasmussen (1993) *J. Immunol.* 150:17), with medium alone or with CpG or non-CpG ODN at the indicated concentrations, or with *E. coli* or calf thymus (50 µg/ml) at 37° C. for 24 hr. All cultures were harvested at 18 hr. and the cells were used as effectors in a standard 4 hr. ⁵¹Cr-release assay against K562 (human) or YAC-1 (mouse) target cells as previously described. For calculation of lytic units (LU), 1 LU was defined as the number of cells needed to effect 30% specific lysis. Where indicated, neutralizing antibodies against IFN-β (Lee Biomolecular, San Diego, Calif.) or IL-12 (C15.1, C15.6, C17.8, and C17.15; provided by Dr. Giorgio Trinchieri, The Wistar Institute, Philadelphia, Pa.) or their isotype controls were added at the initiation of cultures to a concentration of 10 µg/ml. For anti-IL-12 addition, 10 µg of each of the 4 MAB (or isotype controls) were added simultaneously. Recombinant human IL-2 was used at a concentration of 100 U/ml.

Example 12

Prevention of the Development of an Inflammatory Cellular Infiltrate and Eosinophilia in a Murine Model of Asthma

6-8 week old C56BL/6 mice (from The Jackson Laboratory, Bar Harbor, Me.) were immunized with 5,000 *Schistosoma mansoni* eggs by intraperitoneal (i.p.) injection on days 0 and 7. *Schistosoma mansoni* eggs contain an antigen (*Schistosoma mansoni* egg antigen (SEA)) that induces a Th2 immune response (e.g. production of IgE antibody). IgE antibody production is known to be an important cause of asthma.

The immunized mice were then treated with oligonucleotides (30 µg in 200 µl saline by i.p. injection), which either contained an unmethylated CpG motif (i.e. TCCATGACGTTCCCTGACGTT; SEQ ID NO. 10) or did not (i.e. control, TCCATGAGCTTCCTGAGTCT; SEQ ID NO. 11). Soluble SEA (10 µg in 25 µl of saline) was administered by intranasal instillation on days 14 and 21. Saline was used as a control.

Mice were sacrificed at various times after airway challenge. Whole lung lavage was performed to harvest airway and alveolar inflammatory cells. Cytokine levels were measured from lavage fluid by ELISA. RNA was isolated from whole lung for Northern analysis and RT-PCR studies using CsCl gradients. Lungs were inflated and perfused with 4% paraformaldehyde for histologic examination.

FIG. 9 shows that when the mice are initially injected with the eggs i.p., and then inhale the egg antigen (open circle),

many inflammatory cells are present in the lungs. However, when the mice are initially given a nucleic acid containing an unmethylated CpG motif along with the eggs, the inflammatory cells in the lung are not increased by subsequent inhalation of the egg antigen (open triangles).

FIG. 10 shows that the same results are obtained when only eosinophils present in the lung lavage are measured. Eosinophils are the type of inflammatory cell most closely associated with asthma.

FIG. 11 shows that when the mice are treated with a control oligo at the time of the initial exposure to the egg, there is little effect on the subsequent influx of eosinophils into the lungs after inhalation of SEA. Thus, when mice inhale the eggs on days 14 or 21, they develop an acute inflammatory response in the lungs. However, giving a CpG oligo along with the eggs at the time of initial antigen exposure on days 0 and 7 almost completely abolishes the increase in eosinophils when the mice inhale the egg antigen on day 14.

FIG. 12 shows that very low doses of oligonucleotide (<10 µg) can give this protection.

FIG. 13 shows that the resultant inflammatory response correlates with the levels of the Th2 cytokine IL-4 in the lung.

FIG. 14 shows that administration of an oligonucleotide containing an unmethylated CpG motif can actually redirect the cytokine response of the lung to production of IL-12, indicating a Th1 type of immune response.

FIG. 15 shows that administration of an oligonucleotide containing an unmethylated CpG motif can also redirect the cytokine response of the lung to production of IFN-γ, indicating a Th1 type of immune response.

Example 13

CpG Oligonucleotides Induce Human PBMC to Secrete Cytokines

Human PBMC were prepared from whole blood by standard centrifugation over ficoll hypaque. Cells (5×10^5 /ml) were cultured in 10% autologous serum in 96 well microtiter plates with CpG or control oligodeoxynucleotides (24 µg/ml for phosphodiester oligonucleotides; 6 µg/ml for nucleaseresistant phosphorothioate oligonucleotides) for 4 hr in the case of TNF-α or 24 hr. for the other cytokines before supernatant harvest and assay, measured by ELISA using Quantikine kits or reagents from R&D SYSTEMS® (pg/ml) or cytokine ELISA kits from BIOSOURCE™ (for IL-12 assay). Assays were performed as per the manufacturer's instructions. Data are presented in Table 6 as the level of cytokine above that in wells with no added oligodeoxynucleotide.

EQUIVALENTS

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents of the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 60

<210> SEQ ID NO 1
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 1

atggaaggct cagtgttctc 20

<210> SEQ ID NO 2
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 2

atcgacctac gtgcgttctc 20

<210> SEQ ID NO 3
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 3

tccataacgt tctgtgct 20

<210> SEQ ID NO 4
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 4

gctagatggt agcgt 15

<210> SEQ ID NO 5
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 5

gagaacgtcg accttcgat 19

<210> SEQ ID NO 6
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 6

gcatgacggt gagct 15

<210> SEQ ID NO 7
<211> LENGTH: 20

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 7

tccatgacgt tcttgatgct 20

<210> SEQ ID NO 8
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 8

tccatgagct tcttgagtct 20

<210> SEQ ID NO 9
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 9

tccaagacgt tcttgatgct 20

<210> SEQ ID NO 10
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 10

tccatgacgt tcttgacgct 20

<210> SEQ ID NO 11
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 11

tccatgagct tcttgagtct 20

<210> SEQ ID NO 12
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 12

ggggtcaacg ttgagggggg 20

<210> SEQ ID NO 13
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 13

-continued

gctagacggtt agcgt	15
<pre> <210> SEQ ID NO 14 <211> LENGTH: 15 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic oligonucleotide <221> NAME/KEY: modified_base <222> LOCATION: (7)...(7) <223> OTHER INFORMATION: m5c <400> SEQUENCE: 14 </pre>	
gctagacggtt agcgt	15
<pre> <210> SEQ ID NO 15 <211> LENGTH: 15 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic oligonucleotide <221> NAME/KEY: modified_base <222> LOCATION: (13)...(13) <223> OTHER INFORMATION: m5c <400> SEQUENCE: 15 </pre>	
gctagacggtt agcgt	15
<pre> <210> SEQ ID NO 16 <211> LENGTH: 15 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic oligonucleotide <400> SEQUENCE: 16 </pre>	
gcatgacggtt gagct	15
<pre> <210> SEQ ID NO 17 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic oligonucleotide <400> SEQUENCE: 17 </pre>	
atggaaggtc cagcgttctc	20
<pre> <210> SEQ ID NO 18 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic oligonucleotide <400> SEQUENCE: 18 </pre>	
atcgactctc gagcgttctc	20
<pre> <210> SEQ ID NO 19 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic oligonucleotide <221> NAME/KEY: modified_base <222> LOCATION: (3)...(3) <223> OTHER INFORMATION: m5c <221> NAME/KEY: modified_base </pre>	

-continued

<222> LOCATION: (10)...(10)
<223> OTHER INFORMATION: m5c
<221> NAME/KEY: modified_base
<222> LOCATION: (14)...(14)
<223> OTHER INFORMATION: m5c

<400> SEQUENCE: 19

atcgactctc gagcgttctc 20

<210> SEQ ID NO 20
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide
<221> NAME/KEY: modified_base
<222> LOCATION: (3)...(3)
<223> OTHER INFORMATION: m5c

<400> SEQUENCE: 20

atcgactctc gagcgttctc 20

<210> SEQ ID NO 21
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide
<221> NAME/KEY: modified_base
<222> LOCATION: (18)...(18)
<223> OTHER INFORMATION: m5c

<400> SEQUENCE: 21

atcgactctc gagcgttctc 20

<210> SEQ ID NO 22
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 22

atggaaggtc caacgttctc 20

<210> SEQ ID NO 23
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 23

gagaacgctg gaccttccat 20

<210> SEQ ID NO 24
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 24

gagaacgctc gaccttccat 20

<210> SEQ ID NO 25
<211> LENGTH: 20

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 25

gagaacgctc gaccttcgat 20

<210> SEQ ID NO 26
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 26

gagcaagctg gaccttccat 20

<210> SEQ ID NO 27
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide
<221> NAME/KEY: modified_base
<222> LOCATION: (6)...(6)
<223> OTHER INFORMATION: m5c

<400> SEQUENCE: 27

gagaacgctg gaccttccat 20

<210> SEQ ID NO 28
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide
<221> NAME/KEY: modified_base
<222> LOCATION: (14)...(14)
<223> OTHER INFORMATION: m5c

<400> SEQUENCE: 28

gagaacgctg gaccttccat 20

<210> SEQ ID NO 29
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 29

gagaacgatg gaccttccat 20

<210> SEQ ID NO 30
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 30

gagaacgctc cagcactgat 20

<210> SEQ ID NO 31
<211> LENGTH: 20
<212> TYPE: DNA

-continued

```

<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 31

tccatgtcgg tctctgatgct                20

<210> SEQ ID NO 32
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 32

tccatgctgg tctctgatgct                20

<210> SEQ ID NO 33
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide
<221> NAME/KEY: modified_base
<222> LOCATION: (8)...(8)
<223> OTHER INFORMATION: m5c

<400> SEQUENCE: 33

tccatgtcgg tctctgatgct                20

<210> SEQ ID NO 34
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide
<221> NAME/KEY: modified_base
<222> LOCATION: (12)...(12)
<223> OTHER INFORMATION: m5c

<400> SEQUENCE: 34

tccatgtcgg tctctgatgct                20

<210> SEQ ID NO 35
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 35

tccatgacgt tctctgatgct                20

<210> SEQ ID NO 36
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 36

tccatgtcgg tctctgctgat                20

<210> SEQ ID NO 37
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence

```

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 37

tccatgctcg tctctgatgct 20

<210> SEQ ID NO 38
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 38

tccatgccgg tctctgatgct 20

<210> SEQ ID NO 39
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 39

tccatggcgg tctctgatgct 20

<210> SEQ ID NO 40
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 40

tccatgacgg tctctgatgct 20

<210> SEQ ID NO 41
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 41

tccatgtcga tctctgatgct 20

<210> SEQ ID NO 42
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 42

tccatgtcgc tctctgatgct 20

<210> SEQ ID NO 43
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 43

tccatgtcgt tctctgatgct 20

-continued

<210> SEQ ID NO 44
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 44

tccatgacgt tectgatgct 20

<210> SEQ ID NO 45
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 45

tccataacgt tectgatgct 20

<210> SEQ ID NO 46
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 46

tccatgacgt cectgatgct 20

<210> SEQ ID NO 47
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 47

tccatcacgt gectgatgct 20

<210> SEQ ID NO 48
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 48

gcatgacggt gagct 15

<210> SEQ ID NO 49
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 49

gctagatggt agcgt 15

<210> SEQ ID NO 50
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

-continued

```

<400> SEQUENCE: 50
ggggtcagtc ttgagggggg          20

<210> SEQ ID NO 51
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 51
gctagacggt agtgt              15

<210> SEQ ID NO 52
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide
<221> NAME/KEY: modified_base
<222> LOCATION: (8)...(8)
<223> OTHER INFORMATION: m5c

<400> SEQUENCE: 52
gctagacctt agtgt              15

<210> SEQ ID NO 53
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide
<221> NAME/KEY: modified_base
<222> LOCATION: (8)...(8)
<223> OTHER INFORMATION: m5c

<400> SEQUENCE: 53
tccatgtcgt tctctgatgct       20

<210> SEQ ID NO 54
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 54
tccatgacgt tctctgatgct       20

<210> SEQ ID NO 55
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 55
tctcccagcg tgcgccat          18

<210> SEQ ID NO 56
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

```


-continued

```

<400> SEQUENCE: 56
catttccacg atttccca                18

<210> SEQ ID NO 57
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 57
ggcgttattc ctgactcgcc            20

<210> SEQ ID NO 58
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 58
cctacgttgt atgcgcccag ct         22

<210> SEQ ID NO 59
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 59
atggactctc cagcgttctc            20


<210> SEQ ID NO 60
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide
<221> NAME/KEY: modified_base
<222> LOCATION: (14)...(14)
<223> OTHER INFORMATION: m5c

<400> SEQUENCE: 60
atcgactctc gagcgttctc            20

```

The invention claimed is:

1. A method for treating an allergic condition, the method comprising administering to a subject in need of treatment of an allergic condition an immunostimulatory oligonucleotide 8 to 100 nucleotides long, comprising:



wherein the C is unmethylated and wherein X_1 , X_2 , X_3 , and X_4 are nucleotides, wherein the oligonucleotide is administered without an allergen, in an effective amount to treat the allergic condition.

2. The method of claim 1, wherein 5' $X_1X_2CGX_3X_4$ 3' is not a palindrome.

3. The method of claim 1, wherein the allergic condition is selected from the group consisting of eczema, allergic rhinitis, hay fever, urticaria, and food allergies.

4. The method of claim 1, wherein the immunostimulatory oligonucleotide is 8 to 40 nucleotides long.

5. The method of claim 1, wherein the immunostimulatory oligonucleotide is a stabilized oligonucleotide.

6. The method of claim 5, wherein the immunostimulatory oligonucleotide comprises a phosphate backbone modification which is a phosphorothioate or phosphorodithioate modification.

7. The method of claim 1, wherein the immunostimulatory nucleic acid is administered to the subject orally.

8. The method of claim 1, wherein the immunostimulatory nucleic acid is administered to the subject transdermally.

9. The method of claim 1, wherein the immunostimulatory nucleic acid is administered to the subject by injection.

* * * * *