

Extended Abstract

Impact of vasopressin receptors on regulation of immune response in preeclampsia

Kathryn Huber-Keener, MD, PhD,¹ Sabrina M. Scroggins, PhD,¹ Donna A. Santillan, PhD,¹ Daffney Gomendoza,¹ Lindsay Krotz,¹ Jeremy Sandgren,² Justin L. Grobe, PhD,² Mark K. Santillan, MD, PhD¹

Keywords: Preeclampsia, vasopressin, immune, Arginine vasopressin, AVP

Preeclampsia is a common disorder of pregnancy resulting in increased blood pressure and end organ effects. The pathogenesis of preeclampsia is multifactorial. Arginine vasopressin (AVP) is increased in preeclampsia, and the chronic infusion of AVP throughout gestation has previously been shown to be sufficient to produce a phenotype of preeclampsia in C57BL/6J mice representative of some of the cardiovascular and renal events seen in humans. Alterations in T-helper cell populations and their effector cytokines are also known to occur in preeclampsia. Therefore, we proposed that the increased secretion of AVP may be responsible for the immune changes that occur in preeclampsia. We also

hypothesized that known pharmacological AVP antagonist, vaptans, may be able to reverse the effects of AVP infusion. Using our previously published AVP infusion model for preeclampsia, we compared saline controls to mice and their offspring infused with AVP with or without an AVP receptor antagonist vaptan. AVP infusion throughout pregnancy in mice resulted in decreased anti-inflammatory cytokines IL-4 and IL-10 in the maternal kidneys and fetal kidneys, while it increased pro-inflammatory cytokines IFN- γ in maternal plasma and IL-17 in the placenta and amniotic fluid. Specifically blocking AVPR-2 increased production of IL-4 and IL-10 in maternal and fetal

¹Department of Obstetrics and Gynecology, Carver College of Medicine, University of Iowa Hospitals and Clinics, Iowa City, IA, 52242

²Department of Pharmacology, University of Iowa, Roy J. and Lucille A. Carver College of Medicine, Iowa City, Iowa 52242

Please cite this abstract as: Huber-Keener K, Scroggins SM, Santillan DA, Gomendoza D, Krotz L, Sandgren J, Grobe JL, Santillan MK. Impact of vasopressin receptors on regulation of immune response in preeclampsia. *Proc Obstet Gynecol.* 2018;8(1):Article 9 [2 p.]. Available from: <http://ir.uiowa.edu/pog/> Free full text article.

Corresponding author: Kathryn Huber-Keener Department of Obstetrics and Gynecology, University of Iowa, 200 Hawkins Drive, Iowa City, IA 42242, kathryn-huber-keener@uiowa.edu

Copyright: © 2018 Huber-Keener, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

kidneys, increasing the anti-inflammatory response. Interestingly, correction of IL-4 in the placenta required blockade of both AVPR-1a/2 with conivaptan. Proinflammatory marker IL-17 levels in the placenta were corrected by blocking AVPR-1a with relcovaptan, indicating that V1a may be responsible in the placenta for proinflammatory IL-17 production. Overall, the data seem to suggest that AVPR-1a plays a more dominant role in increasing pro-inflammatory markers,

while AVPR2 plays more of a role in decreasing anti-inflammatory markers in preeclampsia. These results also support the ability of AVP to induce the immunological changes seen in preeclampsia.

Presented at "Advances in Gynecologic Cancers," the University of Iowa Obstetrics and Gynecology Postgraduate conference, 10 November 2017, Marriott Coralville Hotel & Conference Center, Coralville, Iowa.